

Klein foams as families of real forms of Riemann surfaces

SABIR M. GUSEIN-ZADE AND SERGEY M. NATANZON

Klein foams are analogues of Riemann surfaces for surfaces with one-dimensional singularities. They first appeared in mathematical physics (string theory etc.). By definition a Klein foam is constructed from Klein surfaces by gluing segments on their boundaries. We show that, a Klein foam is equivalent to a family of real forms of a complex algebraic curve with some structures. This correspondence reduces investigations of Klein foams to investigations of real forms of Riemann surfaces. We use known properties of real forms of Riemann surfaces to describe some topological and analytic properties of Klein foams.

1. Introduction

A Klein foam (a seamed surface) is constructed from Klein surfaces by gluing segments on their boundaries. Klein foams first appeared in mathematical physics (string theory, A-models) [5, 9, 25] (see also [11]) and have applications in mathematics: [2, 3, 10] etc.

A Klein surface is an analog of a Riemann surface for surfaces with boundaries and non-orientable ones [4, 22]. One can consider a Klein foam as an analog of a Riemann surface for surfaces with one-dimensional singularities. Further on we consider only compact Riemann and Klein surfaces.

Following [7] we also require existence of a dianalytic map from a Klein foam to the complex disc. According to [7] such function exists if and only if the corresponding topological foam is strongly oriented. According to [21] this condition makes it possible to extend the 2D topological field theory with Klein surfaces [1] to Klein foams.

2010 Mathematics Subject Classification: 30F50, 57M20, 14H30, 14H37.

Partially supported by the grant RFBR–16-01-00409 (S.G.-Z.). The article was prepared within the framework of the Academic Fund Program at the National Research University Higher School of Economics (HSE) in 2017– 2018 (grant n. 17-01-0030) and by the Russian Academic Excellence Project "5-100". (S.N.).

In this paper we prove, that a Klein foam is a collection of real forms of a complex algebraic curve. Any Klein surface is the quotient S/τ , where $\tau : S \rightarrow S$ is an antiholomorphic involution of a Riemann surface S . In terms of algebraic geometry the pair (S, τ) is a complex algebraic curve with an involution of complex conjugation. Thus it is a real algebraic curve [4]. We will say also that τ is a real form of S . The fixed points of τ correspond to the boundary $\partial(S/\tau)$ of the quotient S/τ and to real points of the corresponding real algebraic curve. The fixed points form closed contours that are called *ovals* [23]. In what follows we consider only real algebraic curves with real points, that is with non-empty $\partial(S/\tau)$.

A general Riemann surface has not more than one antiholomorphic involution. However, there exist Riemann surfaces with several antiholomorphic involutions [13, 14, 17]. Here we prove that the Klein foams are in one-to-one correspondence with the equivalence classes of the families of real forms of Riemann surfaces, i.e. of the collections $\{\widehat{S}, G, (G_1, \widehat{\tau}_1), \dots, (G_r, \widehat{\tau}_r)\}$ consisting of:

- 1) a compact Riemann surface \widehat{S} ;
- 2) a finite subgroup G of the group $\text{Aut}(\widehat{S})$ of holomorphic automorphisms of \widehat{S} ;
- 3) real forms $\widehat{\tau}_1, \dots, \widehat{\tau}_r$ of \widehat{S} such that $\widehat{\tau}_i G \widehat{\tau}_i = G$;
- 4) subgroups $G_i \subset G$ ($i = 1, 2, \dots, r$) generating G and such that $\widehat{\tau}_i G_i \widehat{\tau}_i = G_i$.

A collection of real forms of a Riemann surface has some non-trivial “collective” properties. Initially these properties were observed in [12, 15, 16, 18]. These results were developed and refined in a long series of publications (see [6] and references therein). We use these properties of real forms to describe some non-trivial combinatorial properties of analytic structures of Klein foams. In particular, we give bounds for the number of non-isomorphic Klein surfaces in a Klein foam and for the total number of ovals in a Klein foam.

2. Klein foams

A Klein foam is a topological foam with an analytic structure. A topological foam is obtained from surfaces with boundaries by gluing some segments on the boundaries [2, 25]. For our goal we modify this definition considering instead of a surface with boundary a surface without boundary, but with

an involution such that the corresponding quotient is homeomorphic to the first one. This change of the definition does not change the notion itself.

A *generalized graph* is a one-dimensional space which consist of finitely many vertices and edges, where edges are either segments connecting **different** vertices or isolated circles without vertices on them. A pair of vertices may be connected by several edges.

A *normal topological foam* Ω is a triple (S, Δ, φ) , where

- $S = S(\Omega)$ is a closed (i.e. compact without boundary, but usually disconnected) oriented surface with a reversing the orientation involution with the fixed point set being a closed curve $L \neq \emptyset$, that is finite disconnected union of simple contours;
- $\Delta = \Delta(\Omega)$ is a generalized graph;
- $\varphi = \varphi_\Omega : L \rightarrow \Delta$ is *the gluing map*, that is, a map such that:
 - (a) $\text{Im } \varphi = \Delta$;
 - (b) on each connected component of L , φ is a homeomorphism on a circle in Δ ;
 - (c) for an edge l of Δ , any connected component of S contains at most one connected component of $\varphi^{-1}(l \setminus \partial l)$;
 - (d) (the normality condition) for $\check{\Omega} = S \cup_\varphi \Delta$ (the result of the gluing of S along Δ) and for each vertex v from the set Ω_b of vertices of the graph Δ , its punctured neighbourhood in $\check{\Omega}$ is connected.

A triple (S, Δ, φ) which satisfies all the properties above but (c) will be called a *topological pseudofoam*. The same terminology will be applied to analytic and Klein foams defined below.

Normal topological foams arise naturally in the theory of Hurwitz numbers [2, 3]. A foam $\Omega = (S, \Delta, \varphi)$ will be called *connected* if $\check{\Omega}$ is connected. In what follows topological foams are assumed to be normal and connected. Let Ω_b be the set of vertices of the graph Δ .

A *morphism* f of topological foams $\Omega' \rightarrow \Omega''$ ($\Omega' = (S', \Delta', \varphi')$, $\Omega'' = (S'', \Delta'', \varphi'')$) is a pair (f_S, f_Δ) of (continuous) maps $f_S : S' \rightarrow S''$ and $f_\Delta : \Delta' \rightarrow \Delta''$ such that f_S is an orientation preserving ramified covering commuting with the involutions on S' and S'' , $\varphi'' \circ f_S = f_\Delta \circ \varphi'$ and $f_\Delta|_{\Delta' \setminus \Omega'_b}$ is a local homeomorphism $\Delta' \setminus \Omega'_b \rightarrow \Delta'' \setminus \Omega''_b$.

An *analytic foam* is a topological foam $\Omega = (S, \Delta, \varphi)$, where S is a compact Riemann surface, the involution of which is antiholomorphic. A morphism f of analytic foams $\Omega' \rightarrow \Omega''$ ($\Omega' = (S', \Delta', \varphi')$, $\Omega'' = (S'', \Delta'', \varphi'')$) is

a morphism (f_S, f_Δ) of the corresponding topological foams such that f_S is complex analytic.

The simplest analytic foam is $\Omega_0 = (\overline{\mathbb{C}}, S^1, Id)$, where $\overline{\mathbb{C}} = \mathbb{C} \cup \infty$ is the Riemann sphere with the involution $z \mapsto \bar{z}$. An *analytic function* on an analytic foam Ω is a morphism of Ω to Ω_0 .

A *Klein foam* is an analytic foam $\Omega = (S, \Delta, \varphi)$ admitting an everywhere locally non-constant analytic function. This condition appeared first in [7] and is equivalent to the condition of being “strongly oriented” for the corresponding topological foam. The category of strongly oriented topological foams [21] is a subcategory of topological foams that allows the topological field theory to be extended to the Klein topological field theory [1].

A Klein (pseudo)foam Ω will be called *compressed* if the coincidence of $F(x)$ and $F(x')$ for two points x and x' of Δ_Ω and for all analytic functions F on Ω implies that $x = x'$. For any Klein foam Ω there exists a unique compressed Klein pseudofoam Ω' with a morphism $\Psi : \Omega \rightarrow \Omega'$ such that Ψ_S is an isomorphism. This follows from the following fact. In [7, Theorem 2.1], it was shown that there exists a connected Riemann surface \check{S} with an antiholomorphic involution $\check{\tau}$ and an analytic nowhere locally constant maps $f : \check{\Omega} \rightarrow \check{S}$, commuting with the involutions on S and on \check{S} , such that the corresponding morphism of the foams establishes an isomorphism between the fields of analytic functions on Ω and on \check{S} respectively. One can see that Ω' is obtained from Ω by gluing all the points x and x' of $\Delta(\Omega)$ with $f(x) = f(x')$. (In particular the graph $\Delta_{\Omega'}$ coincides with the set of the real points of \check{S} cut into edges by the ramification points of f .) We shall say that Ω' is the *compressing* of Ω .

Two Klein foams Ω_1 and Ω_2 will be called *weakly isomorphic* if their compressings are isomorphic.

3. Real forms

Definition 1. An *equipped family of real forms* of a Riemann surface \widehat{S} is a collection $\{\widehat{S}, G, (G_1, \widehat{\tau}_1), \dots, (G_r, \widehat{\tau}_r)\}$ consisting of:

- a compact Riemann surface \widehat{S} ;
- antiholomorphic involutions (i.e. real forms) $\{\widehat{\tau}_1, \dots, \widehat{\tau}_r\}$ of \widehat{S} ;
- a finite subgroup G of the group of holomorphic automorphisms of $\text{Aut}(S)$ such that $\widehat{\tau}_i G \widehat{\tau}_i = G$ for each $i = 1, 2, \dots, r$;
- subgroups $G_i \subset G$ such that $\widehat{\tau}_i G_i \widehat{\tau}_i = G_i$ for $i = 1, 2, \dots, r$ and G as a group is generated by G_1, \dots, G_r ;

- the inclusion $G_i \subset G$ generates a morphism of Klein surfaces $S/G_i \rightarrow S/G$ that is a homeomorphism on any oval of S/G .

Definition 2. Two equipped families of real forms of Riemann surfaces

$$\{\widehat{S}, G, (G_1, \widehat{\tau}_1), \dots, (G_r, \widehat{\tau}_r)\} \quad \text{and} \quad \{\widehat{S}, G', (G'_1, \widehat{\tau}'_1), \dots, (G'_r, \widehat{\tau}'_r)\}$$

are equivalent if there exists $h \in \text{Aut } \widehat{S}, h_i \in G, l_i \in G_i$, such that

$$hGh^{-1} = G', \quad hh_iG_ih_i^{-1}h^{-1} = G'_i, \quad hh_i l_i \widehat{\tau}_i l_i^{-1} h_i^{-1} h^{-1} = \widehat{\tau}'_i.$$

Two equipped families of real forms of Riemann surfaces

$$\{\widehat{S}, G, (G_1, \widehat{\tau}_1), \dots, (G_r, \widehat{\tau}_r)\} \quad \text{and} \quad \{\widehat{S}', G', (G'_1, \widehat{\tau}'_1), \dots, (G'_r, \widehat{\tau}'_r)\}$$

are equivalent if there exists an analytic isomorphism $H : \widehat{S} \rightarrow \widehat{S}'$ such that

$$\{\widehat{S}', HGH^{-1}, (HG_1H^{-1}, H\widehat{\tau}_1H^{-1}), \dots, (HG_rH^{-1}, H\widehat{\tau}_rH^{-1})\}$$

and

$$\{\widehat{S}', G'(G'_1, \widehat{\tau}'_1), \dots, (G'_r, \widehat{\tau}'_r)\}$$

are equivalent.

4. Main theorem

Theorem 1. *There exists a natural one-to-one correspondence between the classes of weakly isomorphic Klein foams and the equivalence classes of equipped families of real forms of Riemann surfaces.*

Proof. Consider a Klein foam $\Omega = (S, \Delta, \varphi)$ with S consisting of connected components $S_i, i = 1, \dots, r$. Let τ_i be the restriction of the (antiholomorphic) involution to S_i .

In [7, Theorem 2.1], it was shown that there exists a connected Riemann surface \check{S} with an antiholomorphic involution $\check{\tau}$ and analytic nowhere locally constant maps $f : \check{\Omega} \rightarrow \check{S}$, commuting with the involutions on S and on \check{S} , such that the corresponding morphism of the foams establishes an isomorphism between the fields of analytic functions on Ω and on \check{S} respectively. Consider the restrictions $f_i : S_i \rightarrow \check{S}$ of f to S_i .

Let \check{S}° be the surface \check{S} without all the critical values of the maps f_i and $S^\circ = f^{-1}(\check{S}^\circ)$. Consider an uniformization $U \rightarrow U/\check{\Gamma} = \check{S}^\circ$. (Excluding

trivial cases we assume that U is the upper half-plane in \mathbb{C} and Γ is a Fuchsian group.)

The restriction $f_i^\circ : S_i^\circ \rightarrow \check{S}^\circ$ of f_i to $S_i^\circ = S_i \cap S^\circ$ is a covering without ramification. Thus there exists a subgroup $\Gamma_i \subset \check{\Gamma}$, that uniformises $S_i^\circ = U/\Gamma_i$ and generates the map f_i° by the inclusion $\Gamma_i \subset \check{\Gamma}$. This subgroup Γ_i is well-defined up to conjugation in Γ . Let σ_i be a lifting to U of the involution τ_i to S_i° , such that $\sigma_i \Gamma_i \sigma_i = \Gamma_i$. This σ_i is well-defined up to conjugation in Γ_i .

The subgroup Γ_i has a finite index in Γ and there are only finitely many different subgroups of $\check{\Gamma}$ conjugate to Γ_i . Let us consider the intersection of all the subgroups Γ_i and all their conjugates:

$$\hat{\Gamma} = \bigcap_{j=1}^r \bigcap_{a \in \check{\Gamma}} a^{-1} \Gamma_j a \subset \check{\Gamma}.$$

Since all the involutions τ_i give one and the same involution $\check{\tau}$ on the surface \check{S} , one has $\sigma_i \sigma_j \in \check{\Gamma}$ for all i and j . Thus $\sigma_i a \sigma_j \in \check{\Gamma}$ for any $a \in \check{\Gamma}$ and

$$\begin{aligned} \sigma_i \hat{\Gamma} \sigma_i &= \sigma_i \left(\bigcap_{j=1}^r \bigcap_{a \in \check{\Gamma}} a^{-1} \Gamma_j a \right) \sigma_i \\ &= \bigcap_{j=1}^r \bigcap_{a \in \Gamma} \sigma_i a^{-1} \sigma_j \Gamma_j \sigma_j a \sigma_i = \bigcap_{j=1}^r \bigcap_{b \in \Gamma} b^{-1} \Gamma_j b = \hat{\Gamma}. \end{aligned}$$

Consider now the Riemann surface $\hat{S}^\circ = U/\hat{\Gamma}$. The involution σ_i generates an involution $\hat{\tau}_i^0$ on \hat{S}° . The inclusion $\hat{\Gamma} \subset \check{\Gamma}$ generates a covering $\Phi^\circ : \hat{S}^\circ \rightarrow \check{S}^\circ$. The inclusion $\check{S}^\circ \subset \check{S}$ generates a branching covering of Riemann surfaces $\hat{S} \rightarrow \check{S}$. The involution $\hat{\tau}_i^0$ generates an antiholomorphic involution $\hat{\tau}_i$ on \hat{S} . Moreover $\hat{\Gamma}$ is a normal subgroup of $\check{\Gamma}$ and thus the subgroup $\check{\Gamma}/\hat{\Gamma}$ in a natural way is isomorphic to a subgroup $G \subset \text{Aut}(\hat{S})$ of the group of biholomorphic automorphisms of \hat{S} .

The subgroups Γ_i from this construction also have an interpretation in terms of the Riemann surface \hat{S} . Let us consider the group $\text{Aut}^*(\hat{S})$ of biholomorphic and antiholomorphic automorphisms of \hat{S} . All the involutions $\hat{\tau}_i$ belong to $\text{Aut}^*(\hat{S})$. Moreover, the involution σ_i preserves the subgroup $\Gamma_i \subset \check{\Gamma}$. The homomorphism $\phi : \check{\Gamma}/\hat{\Gamma} \rightarrow \text{Aut}(\hat{S})$ maps the subgroup Γ_i to a subgroup $G_i \subset G$, such that $\hat{\tau}_i G_i \hat{\tau}_i = G_i$.

Thus we have constructed an equipped family of real forms of a compact Riemann surface corresponding to a Klein foam $\Omega = (S, \Delta, \varphi)$. This family

depends on the choice of the uniformizing subgroups Γ_i and of the pull-backs σ_i . This means that the family is defined up to equivalence.

Since the compressing of a real foam depends only on the maps f_i , the weak equivalence class of the foam Ω is determined only by the equivalence class of the corresponding equipped family of real forms of a Riemann surface. \square

Remark 1. One can see that there are finitely many Klein foams corresponding to an equivalence class of equipped families of real forms of a Riemann surface. They are determined by some additional combinatorial data.

5. Some topological and analytical properties of Klein foams

Consider a Klein foam

$$\Omega = (S, \Delta, \varphi), \quad \text{where} \quad S = \prod_{i=1}^r (S_i, \tau_i).$$

Denote by g_i and k_i the genus of S_i and the number of ovals of S_i (i.e. the number of the connected components of $\partial(S_i/\tau_i)$) respectively.

We say that (S_i, τ_i) and (S_j, τ_j) are *foam-equivalent* if the images under φ of the fixed point sets of τ_i and τ_j coincide. Otherwise we say that (S_i, τ_i) and (S_j, τ_j) are *foam-different*.

Recall that Klein surfaces (S', τ') and (S'', τ'') are isomorphic if there exists a biholomorphic map $h : S' \rightarrow S''$ such that $\tau'' = h\tau'h^{-1}$. We say that a real algebraic curve (S', τ') *covers* (S'', τ'') if there exists a holomorphic map $h : S' \rightarrow S''$ such that $\tau''h = h\tau'$. We say that Klein surfaces (S', τ') and (S'', τ'') are *weakly equivalent* if there exists a Klein surface (S^*, τ^*) covering both (S', τ') and (S'', τ'') .

Consider the equipped family of real forms

$$F = \{\hat{S}, G, (G_1, \hat{\tau}_1), \dots, (G_r, \hat{\tau}_r)\}$$

corresponding to Ω . The main topological invariants of the family are the genus \hat{g} of \hat{S} , the numbers of the ovals $\hat{k}_i = |\pi_0(\partial(\hat{S}/\hat{\tau}_i))|$ and the cardinalities $|G_i|$ of the subgroups. From the definitions it follows that $(S_i, \tau_i) = (\hat{S}/G_i, \hat{\tau}_i/G_i)$. Therefore

$$g_i \leq \frac{\hat{g} - 1}{|G_i|} + 1 \quad \text{and} \quad k_i \leq \hat{k}_i.$$

Let us describe some properties of foams related with their genres. These properties depend not only on the topological properties of a foam $\Omega = (S, \Delta, \varphi)$, but also on a holomorphic structure on it. Further on we assume that $\hat{g} > 1$. The topological classification of the equipped families of real forms for $\hat{g} \leq 1$ is more simple, but requires other methods.

Corollary 1. *Let $\Omega = (S, \Delta, \varphi)$ be a Klein foam such that S contains the union $\coprod_{i=1}^r S_i$ with all (S_i, τ_i) foam-different. Then $\sum_{i=1}^r k_i \leq 42(\hat{g} - 1)$.*

Proof. The involutions $\hat{\tau}_i$ corresponding to foam-different Klein surfaces are different. On the other hand, the sum of of the numbers of all the ovals of all the real forms of a Riemann surface of genus $\hat{g} > 1$ is at most $42(\hat{g} - 1)$ [15, 18]. Thus $\sum_{i=1}^r \hat{k}_i \leq 42(\hat{g} - 1)$. \square

Corollary 2. *Let $\Omega = (S, \Delta, \varphi)$ be a Klein foam such that S contains the union $\coprod_{i=1}^r S_i$ with all (S_i, τ_i) not weakly equivalent. Then $r \leq 2(\sqrt{\hat{g}} + 1)$ for any $\hat{g} > 1$ and $r \leq 4$, if \hat{g} is even.*

Proof. The involutions $\hat{\tau}_i$ corresponding to not weakly equivalent Klein surfaces are not conjugate in \hat{G} . On the other hand, the number of conjugation classes of involution on a Riemann surface of genus \hat{g} is at most $2(\sqrt{\hat{g}} + 1)$ for any $\hat{g} > 1$ [12] and at most 4 if \hat{g} is even [8]. \square

Corollary 3. *Let $\Omega = (S, \Delta, \varphi)$ be a Klein foam such that S contains the union $\coprod_{i=1}^r S_i$ with all (S_i, τ_i) not weakly equivalent. In addition assume that either $r = 3, 4$ or all the surfaces \hat{S}/G_i are orientable. Then $\sum_{i=1}^r k_i \leq 2\hat{g} - 2 + 2^{r-3}(9 - r) < 2\hat{g} + 30$.*

Proof. The corresponding bounds for the number of involutions on a Riemann surface are proved in [16, 18–20]. \square

References

- [1] A. Alexeevski and S. Natanzon, *Noncommutative two-dimensional topological field theory and Hurwitz numbers for real algebraic curves*, *Selecta Math. (N.S.)* **12** (2006), no. 3-4, 307–377 (first appeared in [arXiv:math/0202164](https://arxiv.org/abs/math/0202164)).

- [2] A. Alexeevski and S. Natanzon, *The algebra of bipartite graphs and Hurwitz numbers of seamed surfaces*, *Izv. Math.* **72** (2008), no. 4, 627–646.
- [3] A. Alexeevski and S. Natanzon, *Hurwitz numbers for regular coverings of surfaces by seamed surfaces and Cardy–Frobenius algebras of finite groups*, *Amer. Math. Soc. Transl., ser. 2*, **224** (2008), 1–25.
- [4] N. L. Alling and N. Greenleaf, *Foundation of the theory of Klein surfaces*, *Lect. Not. Math.* **219** (1971), 117pp.
- [5] J. C. Baez, *An introduction to spin foam models of BF theory and quantum gravity*, *Geometry and quantum physics* (Schladming, 1999), 25–93, *Lecture Notes in Phys.* **543**, Springer, Berlin, 2000.
- [6] E. Bujalance, F. J. Cirre, J. M. Gamboa, and G. Gromadski, *Symmetries of compact Riemann surfaces*, *Lecture Notes in Mathematics*, Springer-Verlag, Berlin–Heidelberg, **2007** (2010), 155pp.
- [7] A. F. Costa, S. M. Gusein-Zade, and S. M. Natanzon, *Klein foams*, *Indiana Univ. Math. J.* **60** (2011), no. 3, 985–995.
- [8] G. Gromadski and M. Izquierdo, *Real forms of a Riemann surface of even genus*, *Proc. Amer. Math. Soc.* **126** (1998), no. 12, 3475–3479.
- [9] M. Khovanov and L. Rozansky, *Topological Landau-Ginzburg models on the world-sheet foam*, *Adv. Theor. Math. Phys.* **11** (2007), no. 2, 233–259.
- [10] M. Mackaay and P. Vaz, *The foam and the matrix factorization sl_3 link homologies are equivalent*, *Algebr. Geom. Topol.* **8** (2008), no. 1, 309–342.
- [11] A. D. Mironov, A. Morozov, and S. M. Natanzon, *Hurwitz theory avatar of open-closed strings*, *The European Physical Journal C — Particles and Fields* **73** (2013), no. 2, 1–10.
- [12] S. M. Natanzon, *On the order of a finite group of homeomorphisms of a surface onto itself and the number of real forms of a complex algebraic curve*, *Dokl. Akad. Nauk SSSR* **242** (1978), no. 4, 765–768 (Russian); English translation in: *Soviet Math. Dokl.* **19** (1978), no. 5, 1195–1199.
- [13] S. M. Natanzon, *Automorphisms of the Riemann surface of an M -curve*, *Funktional. Anal. i Prilozhen.* **12** (1978), no. 3, 82–83 (Russian); English translation in: *Functional Anal. Appl.* **12** (1978), no. 3, 228–229.

- [14] S. M. Natanzon, *Automorphisms and real forms of a class of complex algebraic curves*, Funktsional. Anal.i Prilozhen. **13** (1979), no. 2, 89–90 (Russian); English translation in: Functional Anal. Appl. **13** (1979), 148–150.
- [15] S. M. Natanzon, *On the total number of ovals of real forms of a complex algebraic curve*, Uspekhi Mat. Nauk **35** (1980), no. 1, 207–208 (Russian); English translation in: Russian Math. Surveys **35** (1980), no. 1, 223–224.
- [16] S. M. Natanzon, *On the total number of ovals of four complex-isomorphic real algebraic curves*, Uspekhi Mat. Nauk **35** (1980), no. 4, 184.
- [17] S. M. Natanzon, *Lobachevskian geometry and automorphisms of complex M -curves*, Sel. Math. Sov. **1** (1981), no. 1, 81–99.
- [18] S. M. Natanzon, *Finite groups of homeomorphisms of surfaces and real forms of complex algebraic curves*, Trudy Moscow Math. Obshch. **51** (1988), 3–53 (Russian); English translation in: Trans. Moscow Math. Soc. (1989), 1–51.
- [19] S. M. Natanzon, *A Harnack-type theorem for a family of complex-isomorphic real algebraic curves*, Uspekhi Mat. Nauk **52** (1997), no. 6, 173–174 (Russian); English translation in: Russian Math. Surveys **52** (1997), no. 6, 1314–1315.
- [20] S. M. Natanzon, *Geometry and algebra of real forms of complex curves*, Math. Zeit. **243** (2003), 391–407.
- [21] S. M. Natanzon, *Cyclic foam topological field theory*, J. Geom. Phys. **60** (2010), no. 6-8, 874–883.
- [22] S. M. Natanzon, *Klein surfaces*, Russian Math. Surveys. **45** (1990), no. 6, 53–108.
- [23] S. M. Natanzon, *Moduli spaces of real algebraic curves*, Proc. Moscow Math. Soc. **37** (1978), 219–253.
- [24] S. M. Natanzon, *Moduli of Riemann surfaces, real algebraic curves and its superanalogs*, Translations of Math. Mon. **225** (2004), Amer. Math. Soc., 159pp.
- [25] L. Rozansky, *Topological A -models on seamed Riemann surfaces*, Adv. Theor. Math. Phys. **11** (2007), no. 4, 517–529.

FACULTY OF MECHANICS AND MATHEMATICS
MOSCOW STATE LOMONOSOV UNIVERSITY
MOSCOW, GSP-1, 119991, RUSSIA
E-mail address: `sabir@mccme.ru`

NATIONAL RESEARCH UNIVERSITY HIGHER SCHOOL OF ECONOMICS
20 MYASNITSKAYA ULITSA, MOSCOW 101000, RUSSIA
E-mail address: `natanzons@mail.ru`

