

GENERICALLY TRANSITIVE ACTIONS ON MULTIPLE FLAG VARIETIES

ROSTISLAV DEVYATOV

ABSTRACT. Let G be a semisimple algebraic group whose decomposition into a product of simple components does not contain simple groups of type A , and $P \subseteq G$ be a parabolic subgroup. Extending the results of Popov [7], we enumerate all triples (G, P, n) such that (a) there exists an open G -orbit on the *multiple flag variety* $G/P \times G/P \times \dots \times G/P$ (n factors), (b) the number of G -orbits on the multiple flag variety is finite.

INTRODUCTION

Let G be a semisimple connected algebraic group over an algebraically closed field of characteristic zero, and $P \subseteq G$ be a parabolic subgroup. One easily checks that the G -orbits on $G/P \times G/P$ are in bijection with P -orbits on G/P . The Bruhat decomposition of G implies that the number of P -orbits on G/P is finite and that these orbits are enumerated by a subset in the Weyl group W corresponding to G . In particular, there is an open G -orbit on $G/P \times G/P$. So we come to the following questions: for which G , P and $n \geq 3$ is there an open G -orbit on the *multiple flag variety* $(G/P)^n := G/P \times G/P \times \dots \times G/P$? For which G , P and n is the number of orbits finite?

Notice that if G is locally isomorphic to $G^{(1)} \times \dots \times G^{(k)}$, where $G^{(i)}$ are simple, then there exist parabolic subgroups $P^{(i)} \subseteq G^{(i)}$ such that $G/P \cong G^{(1)}/P^{(1)} \times \dots \times G^{(k)}/P^{(k)}$. Hence in the sequel we may assume that G is simple. Moreover, let $\pi: \tilde{G} \rightarrow G$ be a simply connected cover. Then π induces a bijection between parabolic subgroups $P \subseteq G$ and $\tilde{P} \subseteq \tilde{G}$, namely $\tilde{P} = \pi^{-1}(P)$, and an isomorphism $\tilde{G}/\tilde{P} \rightarrow G/P$. Also, \tilde{G}/\tilde{P} may be considered as G -variety since $\text{Ker } \pi$ acts trivially on it. In this sense the isomorphism is G -equivariant. Therefore we may consider only one simple group of each type.

The classification of multiple flag varieties with an open G -orbit for maximal subgroups P was given by Popov in [7]. We need some notation to formulate his result. Fix a maximal torus in G and an associated simple root system $\{\alpha_1, \dots, \alpha_l\}$ of the Lie algebra $\mathfrak{g} = \text{Lie } G$. We enumerate simple roots as in [3]. Let $P_i \subset G$ be the maximal parabolic subgroup corresponding to the simple root α_i .

Theorem 1. [7, Theorem 3] *Let G be a simple algebraic group. The diagonal G -action on the multiple flag variety $(G/P_i)^n$ is generically transitive if and only if $n \leq 2$ or (G, n, i) is an entry in the following table:*

<i>Type of G</i>	(n, i)
A_l	$n < \frac{(l+1)^2}{i(l+1-i)}$
$B_l, l \geq 3$	$n = 3, i = 1, l$
$C_l, l \geq 2$	$n = 3, i = 1, l$
$D_l, l \geq 4$	$n = 3, i = 1, l - 1, l$
E_6	$n = 3, 4, i = 1, 6$
E_7	$n = 3, i = 7$

In [7], the following question was posed: for which non-maximal parabolic subgroups $P \subset G$ is there an open G -orbit in $(G/P)^n$? We solve this problem for all simple groups except for those of type A_l .

Denote the intersection $P_{i_1} \cap \dots \cap P_{i_s}$ by P_{i_1, \dots, i_s} . It is easy to see that P_{i_1, \dots, i_s} is a parabolic subgroup and that every parabolic subgroup is conjugated to some P_{i_1, \dots, i_s} .

Theorem 2. *Let G be a simple algebraic group which is not locally isomorphic to SL_{l+1} , $P \subset G$ be a non-maximal parabolic subgroup and $n \geq 3$. Then the diagonal G -action on the multiple flag variety $(G/P_i)^n$ is generically transitive if and only if $n = 3$ and (G, P) is one of the following pairs:*

Type of G	P
$D_l, l \geq 5$ is odd	$P_{1, l-1}, P_{1, l}$
$D_l, l \geq 4$ is even	$P_{1, l-1}, P_{1, l}, P_{l-1, l}$

Now let us consider actions with a finite number of orbits. Recall that a G -variety X is called *spherical* if a Borel subgroup $B \subseteq G$ acts on X with an open orbit. It is well-known that the number of B -orbits on a spherical variety is finite, see [1], [9]. Equivalently, the number of G -orbits on $G/B \times X$ is finite if X is spherical. Therefore, if $P \subseteq G$ is a parabolic subgroup and X is a spherical G -variety, then the number of G -orbits on $G/P \times X$ is finite. The classification of all pairs of parabolic subgroups (P, Q) such that $G/P \times G/Q$ is spherical is given in [4] and [8]. According to this classification, if (G, P_i) is an entry in the table from Theorem 1, then $G/P_i \times G/P_i$ is spherical and hence the number of G -orbits on $G/P_i \times G/P_i \times G/P_i$ is finite. In the last section we prove that the number of G -orbits on $(G/P)^n$ is infinite if $n \geq 4$. We also check directly that if (G, P) is an entry in the table from Theorem 2, then the number of G -orbits on $G/P \times G/P \times G/P$ is infinite. Thus we come to the following result.

Theorem 3. *Let G be a simple algebraic group, $P \subset G$ be a parabolic subgroup and $n \geq 3$. The following properties are equivalent.*

- (1) *The number of G -orbits on $(G/P)^n$ is finite.*
- (2) *$n = 3$, P is maximal, and there is an open G -orbit on $G/P \times G/P \times G/P$.*
- (3) *$n = 3$, and $G/P \times G/P$ is spherical.*

Corollary 1. *Let $n \geq 3$. The number of G -orbits on $(G/P)^n$ is finite if and only if $n = 3$ and (G, P) is one of the pairs listed in the following table:*

Type of G	P
A_l	any maximal
$B_l, l \geq 2$	P_1, P_l
$C_l, l \geq 3$	P_1, P_l
$D_l, l \geq 4$	P_1, P_{l-1}, P_l
E_6	P_1, P_6
E_7	P_7

Let us mention a more general result for classical groups. Let $Q_{(1)}, \dots, Q_{(n)}$ be parabolic subgroups in G . We call the variety $G/Q_{(1)} \times \dots \times G/Q_{(n)}$ a *generalized multiple flag variety*. The classification of all generalized multiple flag varieties with a finite number of G -orbits is given in [5] for $G = SL_{l+1}$ and in [6] for $G = Sp_{2l}$.

Proofs of Theorems 2 and 3 use methods developed in [7]. The results concerning existence of an open orbit in a linear representation space in Section 2 may be of independent interest. In several cases for $G = SO_{2l}$ the existence of an open orbit on a multiple flag variety is checked directly.

Acknowledgements. I am grateful to my scientific advisor Ivan V. Arzhantsev for posing the problem and for paying attention to my work. Also I thank professor Vladimir L. Popov for useful comments and Dmitri A. Timashev for bringing my attention to the articles [4] and [8].

1. PRELIMINARIES

Let G be a connected simple algebraic group over an algebraically closed field \mathbb{K} of characteristic zero and $\mathfrak{g} = \text{Lie } G$. Fix a Borel subgroup $B \subset G$ and a maximal torus $T \subset B$. These data determine a root system Φ of \mathfrak{g} , a positive root subsystem Φ^+ and a system of simple roots $\Delta \subseteq \Phi^+$, $\Delta = \{\alpha_1, \dots, \alpha_l\}$. Choose a corresponding Chevalley basis $\{x_i, y_i, h_i\}$ of \mathfrak{g} . We have $[h, x_i] = \alpha(h)x_i$, $[h, y_i] = -\alpha(h)y_i$ for all $h \in \mathfrak{t} = \text{Lie } T$ and $h_i = [x_i, y_i]$.

Let $I = \{\alpha_{i_1}, \dots, \alpha_{i_s}\} \subseteq \Delta$ be a subset. The Lie algebra of the parabolic subgroup $P_I := P_{i_1, \dots, i_s}$ is

$$\mathfrak{p} = \mathfrak{b} \oplus \bigoplus_{\alpha \in \Phi_I} \mathfrak{g}_\alpha,$$

where $\mathfrak{b} = \text{Lie } B$ and $\Phi_I \subseteq \Phi^-$ denotes the set of the negative roots such that their decomposition into the sum of simple roots *does not* contain the roots α_i , $i \in I$. For example, $P_\Delta = B$ and $P_\emptyset = G$. It is known that [2, Theorem 30.1] if a parabolic group P contains B , then $P = P_I$ for some $I \subseteq \Delta$. Therefore any parabolic subgroup $P \subseteq G$ is conjugate to some P_I . If $P = P_I$ for some $I \subseteq \Delta$, we denote by P^- the parabolic subgroup whose Lie algebra is

$$\mathfrak{p}^- = \mathfrak{t} \oplus \bigoplus_{\alpha \in -\Phi_I \cup \Phi^-} \mathfrak{g}_\alpha.$$

Denote the weight lattice of T by $\mathfrak{X}(T)$. Let $\mathfrak{X}^+(T)$ be the subsemigroup of dominant weights with respect to B . Assume first that G is simply connected. Then \mathfrak{X}^+ is generated by the fundamental weights π_1, \dots, π_l . Given a dominant weight λ , denote the simple G -module with the highest weight λ by $V(\lambda)$. If G is not simply connected, we may consider a simply connected cover $p: \tilde{G} \rightarrow G$, the dominant weight lattice $\mathfrak{X}^+(p^{-1}(T))$ and the highest weight \tilde{G} -module $V(\lambda)$.

Let G be a simple group and $P = P_{i_1, \dots, i_s}$ be a parabolic subgroup. Notice that if there is an open G -orbit on $(G/P)^n$, then there exists an open G -orbit on $(G/P_i)^n$ for all $i \in \{i_1, \dots, i_s\}$. Indeed, since $P \subseteq P_i$, one has the surjective G -equivariant map $G/P \rightarrow G/P_i$, $gP \mapsto gP_i$. It induces the surjective G -equivariant map $\varphi: (G/P)^n \rightarrow (G/P_i)^n$, and the image of an open G -orbit on $(G/P)^n$ under φ is an open G -orbit on $(G/P_i)^n$. Similarly, if G acts on $(G/P)^n$ with an open orbit and $m < n$, then G acts on $(G/P)^m$ with an open orbit.

Theorem 1 leaves us very few cases of non-maximal parabolic groups to consider. Namely, if $n > 3$ and G is of type B_l , C_l or D_l , then G never acts on $(G/P)^n$ with an open orbit. If $n = 3$ and G is of type B_l or C_l , it suffices to consider $P = P_{1,l}$, and we show that there is no open orbit in this case. If $n = 3$ and G is of type D_l , an open orbit may exist only if $P = P_I$ where $I \subseteq \{\alpha_1, \alpha_{l-1}, \alpha_l\}$. So there are four cases to consider. We reduce the case $P_{1,l-1}$ to the case $P_{1,l}$. If G is of type E_6 , the only parabolic group we should consider is $P = P_{1,6}$. We show that there is no open orbit for $n = 3$. If G is of type E_7 , if there existed an open G -orbit on $(G/P)^n$ for $n \geq 3$, then the only maximal parabolic subgroup containing P would be P_7 , but in this case P should be maximal itself. If G is of type E_8 , F_4 or G_2 , an open orbit exists for no maximal parabolic subgroups for $n \geq 3$, so there are no cases to consider.

Given a group G acting on an irreducible variety X with an open orbit, according to [7] we denote the maximal n such that there is an open G -orbit on X^n by $\text{gtd}(G : X)$. If G acts on X^n with an open orbit, we say that the action $G : X$ is *generically n -transitive*.

We make use of the following fact proved by Popov.

Proposition 1. [7, Corollary 1 (ii) of Proposition 2] *Let G be a simple algebraic group, P be a parabolic subgroup, P^- be an opposite parabolic subgroup, $L = P \cap P^-$ be the corresponding Levi subgroup and \mathfrak{u}^- be the Lie algebra of the unipotent radical of P^- . If P is conjugate to P^- , then $\text{gtd}(G : G/P) = 2 + \text{gtd}(L : \mathfrak{u}^-)$.*

We suppose that the group SO_l acts in the l -dimensional space and preserves the bilinear form whose matrix with respect to a standard basis is

$$Q = \begin{pmatrix} & & & & 1 \\ & 0 & & & \\ & & 1 & & \\ & & & 1 & \\ \dots & & & & 0 \\ 1 & \dots & & & \end{pmatrix}.$$

We denote the l -dimensional projective space by \mathbf{P}^l and the Grassmannian of k -dimensional subspaces in \mathbb{K}^l by $\text{Gr}(k, l)$.

2. EXISTENCE OF AN OPEN ORBIT

2.1. Groups of type B_l . By Theorem 1, it is sufficient to consider the case $P = P_{1,l}$. The Dynkin diagram B_l has no automorphisms, hence P is conjugate to P^- . So we may apply Proposition 1, and it suffices to check that $\text{gtd}(L : \mathfrak{u}^-) = 0$, i. e. L acts on \mathfrak{u}^- with no open orbit.

Let $G = SO_{2l+1}$. Then $L = \mathbb{K}^* \times GL_{l-1}$ and the L -module \mathfrak{u}^- can be decomposed into the direct sum $V_1 \oplus V_2 \oplus V_3 \oplus V_4 \oplus V_5$. Here V_1 is a GL_{l-1} -module $(\mathbb{K}^{l-1})^*$ dual to the tautological one and its \mathbb{K}^* -weight is 1, V_2 is a trivial one-dimensional GL_{l-1} -module of weight 1, V_3 is a tautological GL_{l-1} -module \mathbb{K}^{l-1} of weight 0, V_4 is a GL_{l-1} -module \mathbb{K}^{l-1} of weight 1, V_5 is a GL_{l-1} -module $\Lambda^2 \mathbb{K}^{l-1}$ of weight 0. According to this decomposition, we denote components of a vector $u \in \mathfrak{u}^-$ by u_1, u_2, u_3, u_4, u_5 .

Notice that there exists a GL_{l-1} -invariant pairing between V_1 and V_3 . Its \mathbb{K}^* -weight is 1. Also there exists a GL_{l-1} -invariant pairing between V_1 and V_4 , whose \mathbb{K}^* -weight is 2. Therefore the rational function

$$\frac{(u_1, u_3)^2}{(u_1, u_4)}$$

is a non-constant invariant for $L : \mathfrak{u}^-$, and the action of G on G/P is not generically 3-transitive.

2.2. Groups of type C_l . This case is completely similar to the previous one, and again the only thing we should do is to prove that there is no open L -orbit on \mathfrak{u}^- , where $L = P \cap P^-$ is a Levi subgroup of $P = P_{1,l}$ and \mathfrak{u}^- is the Lie algebra of the unipotent radical of P^- .

Let $G = Sp_{2l}$. Then $L = \mathbb{K}^* \times GL_{l-1}$ and the L -module \mathfrak{u}^- can be written as $V_1 \oplus V_2 \oplus V_3 \oplus V_4$. Here V_1 is a GL_{l-1} -module $(\mathbb{K}^{l-1})^*$ and its \mathbb{K}^* -weight is 1, V_2 is a GL_{l-1} -module \mathbb{K}^{l-1} of weight 1, V_3 is a GL_{l-1} -module $S^2 \mathbb{K}^{l-1}$ of weight 0, V_4 is a trivial GL_{l-1} -module of weight 2. According to this decomposition, we denote components of a vector $u \in \mathfrak{u}^-$ by u_1, u_2, u_3, u_4 .

We see that there exists a GL_{l-1} -invariant pairing between V_1 and V_2 with \mathbb{K}^* -weight is 2. Therefore we have the rational invariant

$$\frac{(u_1, u_2)}{u_4}$$

for $L : \mathfrak{u}^-$, and the action of G on G/P is not generically 3-transitive.

2.3. Groups of type D_l . This time we should consider the following four cases of parabolic subgroups: $P = P_{1,l-1}, P_{1,l}, P_{l-1,l}, P_{1,l-1,l}$. One easily checks that P and P^- are conjugate except for the cases $P = P_{1,l}$, l odd, and $P = P_{1,l-1}$, l odd.

Let $G = SO_{2l}$. There exists a diagram automorphism of G that interchanges α_{l-1} and α_l . It preserves the maximal torus and the Borel subgroup and interchanges $P_{1,l-1}$ and $P_{1,l}$. Therefore, the actions $G : G/P_{1,l-1}$ and $G : G/P_{1,l}$ are either generically 3-transitive or not generically 3-transitive simultaneously.

suppose that u_1 and u_2 are canonical, the only non-zero coordinates of u_3 are the first one and the second one, they are equal, and $(u_2)_1(u_3)_1 = 1$.

Now recall that $L = GL_{l-1} \times \mathbb{K}^*$, the \mathbb{K}^* -weights of V_1 , V_2 and V_3 are 0, 1 and -1 , respectively. Therefore, after applying a suitable element of \mathbb{K}^* , we have $u_1 = S$, $u_2 = (1, 0, \dots, 0)^T$ and $u_3 = (1, 1, 0, \dots, 0)^T$, and the G -action on G/P is generically 3-transitive.

2.3.2. $P = P_{1,l}$. In this case, Proposition 1 applies if and only if l is even.

Let l be even. It is sufficient to prove that there is an open L -orbit on \mathfrak{u}^- .

Again $L = \mathbb{K}^* \times GL_{l-1}$, and the L -module V can be decomposed into three summands, $V = V_1 \oplus V_2 \oplus V_3$, but this time V_1 is a GL_{l-1} -module $\Lambda^2 \mathbb{K}^{l-1}$ and its \mathbb{K}^* -weight is 0, V_2 is a GL_{l-1} -module \mathbb{K}^{l-1} of weight 1, V_3 is a GL_{l-1} -module $(\mathbb{K}^{l-1})^*$ of weight 1. We denote components of an element $u \in \mathfrak{u}^-$ by u_1, u_2, u_3 .

Recall the notation we have introduced for the GL_{l-1} -module V' . Also this time denote $Y_w = V_3 \setminus \text{Ker } w$. Define $W_2 = \cup_{w \in Z} (w \times X_w \times Y_w)$ and $W = \{u \in W_2 : \langle u_2, u_3 \rangle \neq 0\}$. Here $\langle \cdot, \cdot \rangle$ denotes the GL_{l-1} -invariant pairing between V_2 and V_3 . Its \mathbb{K}^* -weight is 2, but the condition $\langle u_2, u_3 \rangle \neq 0$ is not affected by \mathbb{K}^* -action, so W is L -invariant. We are going to prove that L acts transitively on W .

Again we may suppose that $u_1 = R$ and the only non-zero coordinate of u_2 is the first one. Notice that $(u_3)_1 \neq 0$ since $\langle u_2, u_3 \rangle \neq 0$. This time V_2 and V_3 are dual GL_{l-1} -modules, so by applying a suitable element of $(GL_{l-1})_{u_1, \langle u_2 \rangle}$ the coordinates $(u_2)_1$ and $(u_3)_1$ can be made equal.

Consider the vector $v = ((u_3)_2, (u_3)_3, \dots, (u_3)_{l-1})$. It cannot be zero since $u_3 \notin \text{Ker } u_1$. Since Sp_{l-2} acts transitively on $(\mathbb{K}^{l-2})^* \setminus 0$, there exists an element $g \in (GL_{l-1})_{u_1, u_2}$, $g = g_1 \oplus g_2$, $g_1 = 1$, $g_2 \in Sp_{l-2}$ such that $g_2 v = ((u_3)_1, 0, \dots, 0)^T$. In other words, we may suppose that u_1 and u_2 are canonical, the only non-zero coordinates of u_3 are the first one and the second one, and $(u_2)_1 = (u_3)_1 = (u_3)_2$.

This time the \mathbb{K}^* -weights of V_2 and V_3 are both 1, so with the help of the \mathbb{K}^* -action we can satisfy the equality $(u_2)_1 = (u_3)_1 = (u_3)_2 = 1$. Thus, L acts transitively on W , and the G -action on G/P is generically 3-transitive.

Let l be odd. Proposition 1 does not apply, and we have to find $\text{gtd}(G : G/P)$ directly.

Consider the tautological SO_{2l} -module \mathbb{K}^{2l} , and let e_1, \dots, e_{2l} be its standard basis. Let $X' \subset \text{Gr}(l, 2l)$ be the set of all isotropic subspaces of dimension l in \mathbb{K}^{2l} . One easily checks that X' is a disjoint union of two SO_{2l} -orbits, and the group O_{2l} interchanges them. If two subspaces belong to the same SO_{2l} -orbit, then their intersection is non-zero.

Denote the orbit $SO_{2l} \langle e_1, \dots, e_l \rangle \subset X'$ by X . Then X is an irreducible subvariety in $\text{Gr}(l, 2l)$.

For each $s \in X$ let $Y_s \subset \mathbf{P}^{2l-1}$ be the set of all lines contained in s . Clearly, $W = \cup_{s \in X} (s \times Y_s)$ is a closed G -invariant subset in $\text{Gr}(l, 2l) \times \mathbf{P}^{2l-1}$. One easily checks that $G/P = W$.

Let us prove that there exists an open G -orbit on $W \times W \times W$. We impose some conditions on the point $(s_1, a_1, s_2, a_2, s_3, a_3) \in W \times W \times W$ and so define an open subset $Y \subseteq W \times W \times W$. Then we define a point $p \in \text{Gr}(l, 2l) \times \mathbf{P}^{2l-1} \times \text{Gr}(l, 2l) \times \mathbf{P}^{2l-1} \times \text{Gr}(l, 2l) \times \mathbf{P}^{2l-1}$ and prove that (a) each point $y \in Y$ belongs to the same G -orbit that p does, and (b) p belongs to Y . Condition (b) guarantees that Y is not empty.

Let $Y \subseteq W \times W \times W$ be the set of all tuples $(s_1, a_1, s_2, a_2, s_3, a_3)$ such that:

- (1) $s_1 \cap s_2 \cap s_3 = 0$.
- (2) $s_1 + s_2 + s_3 = \mathbb{K}^{2l}$.
- (3) $\dim s_1 \cap s_2 = \dim s_2 \cap s_3 = \dim s_1 \cap s_3 = 1$.
- (4) $\dim(a_1 + a_2 + a_3) = 3$.
- (5) The intersection of the subspaces $s = (s_1 \cap s_2) + (s_2 \cap s_3) + (s_1 \cap s_3)$ and $a = a_1 + a_2 + a_3$ is zero.
- (6) $a_i + s_j + s_k = \mathbb{K}^{2l}$, where $i = 1, 2, 3, j \neq i, k \neq i, j < k$.
- (7) The lines a_i and a_j are not orthogonal for all $i \neq j$.

Notice that if conditions (1)–(3) hold, the sum of subspaces $s_1 \cap s_2$, $s_2 \cap s_3$ and $s_1 \cap s_3$ is direct.

following bases:

$$\begin{aligned} U_1 &= \langle q_1, \dots, q_k \rangle \\ U_2 &= \langle q_{k+1}, \dots, q_{2k} \rangle \\ U_3 &= \langle q_1 - q_{k+1}, \dots, q_{k/2} - q_{k+k/2}, q_{k/2+1} + q_{k+k/2+1}, \dots, q_k + q_{2k} \rangle. \end{aligned}$$

This completes the proof of the lemma. \square

Consider the following basis of \mathbb{K}^{2l} : $g_1, g_2, g_3, q_1, \dots, q_{2l-6}, g_4, g_5, g_6$, where q_i are defined above in the proof of the lemma. Notice that the matrix of the bilinear form in this basis is Q . Define the operator $B: \mathbb{K}^{2l} \rightarrow \mathbb{K}^{2l}$ that maps this basis to the standard one. We know that the matrix of the bilinear form is Q in both bases, so $B \in O_{2l}$.

Let us check that $\det B = 1$. Assume that $\det B = -1$. Since $s_1 = \langle g_1, g_2, g_3, q_1, \dots, q_{l-3} \rangle$, $Bs_1 = \langle e_1, \dots, e_l \rangle \in X$. Since $s_1 \in X$, there exists an operator $C \in SO_{2l}$ such that $CBs_1 = s_1$. Thus, $CB \in (O_{2l})_{s_1}$, $\det CB = -1$, and the O_{2l} -orbit X' cannot be a union of two distinct SO_{2l} -orbits, a contradiction.

Bases of the subspaces Ba_i and Bs_i can be written in terms of e_i using matrices that do not depend on a_i and s_i . Namely, they are the same matrices that we need to write bases of a_i and s_i using g_i and q_i , and the latter do not depend on a_i and s_i . Denote the 6-tuple $(Bs_1, Ba_1, Bs_2, Ba_2, Bs_3, Ba_3)$ by p . It suffices to prove that $p \in Y$. Conditions (1)–(7) hold by the construction of g_i and q_i , but we should check that $(Bs_1, Bs_2, Bs_3) \in X \times X \times X$. It is sufficient to find elements of SO_{2l} that map s_1 to s_2 and s_1 to s_3 . Since $s_i = (s_i \cap S) \oplus (s_i \cap S^\perp)$, we find them as direct sums of elements of $SO(S)$ and $SO(S^\perp)$. The elements of $SO(S)$ are already found, they are cyclic permutations of f_1, f_2, f_3 and f_4, f_5, f_6 . To interchange $s_1 \cap S^\perp$ and $s_2 \cap S^\perp$, consider the map that permutes all the pairs of vectors $g_i \leftrightarrow g_{2l+1-i}$, $i = 1, \dots, l-3$. It is orthogonal and its determinant is 1 since $l-3$ is even. Finally, the operator with the following matrix in the basis g_i maps $s_1 \cap S^\perp$ to $s_3 \cap S^\perp$.

$$\begin{pmatrix} I_{l-3} & 0 \\ D & I_{l-3} \end{pmatrix}$$

where

$$D = \begin{pmatrix} -1 & & & & & & \\ & \ddots & & & & & \\ & & -1 & & & & \\ & & & 1 & & & \\ & & & & \ddots & & \\ 0 & & & & & & 1 \end{pmatrix}.$$

Therefore, SO_{2l} acts transitively on Y , and $\text{gtd}(G: G/P) = 3$.

2.3.3. $P = P_{1,l-1,l}$. The subgroups P and P^- are conjugate for all l . It is sufficient to find $\text{gtd}(L: \mathfrak{u}^-)$, where $L = (\mathbb{K}^*)^2 \times GL_{l-2}$ and the L -module \mathfrak{u}^- is isomorphic to the direct sum of 7 simple modules that we denote by V_1, \dots, V_7 . Namely, V_1 is a GL_{l-1} -module $(\mathbb{K}^{l-2})^*$ and its $(\mathbb{K}^*)^2$ -weight is $(1, 0)$, V_2 is a trivial GL_{l-2} -module of weight $(1, 1)$, V_3 is a GL_{l-2} -module \mathbb{K}^{l-2} of weight $(0, 1)$, V_4 is a trivial GL_{l-2} -module of weight $(1, -1)$, V_5 is a GL_{l-2} -module \mathbb{K}^{l-2} of weight $(0, -1)$, V_6 is a GL_{l-2} -module \mathbb{K}^{l-2} of weight $(1, 0)$, V_7 is a GL_{l-2} -module $\Lambda^2 \mathbb{K}^{l-2}$ of weight $(0, 0)$. Denote the components of $u \in \mathfrak{u}^-$ by u_1, \dots, u_7 .

There exists a GL_{l-2} -invariant pairing between V_1 and V_3 whose $(\mathbb{K}^*)^2$ -weight is $(1, 1)$. The following function is a rational L -invariant:

$$\frac{(u_1, u_3)}{u_4}.$$

Thus, the G -action on G/P is not generically 3-transitive.

2.4. Groups of type E_6 . The only parabolic subgroup to consider is $P = P_{1,6}$. The set $\{1, 6\}$ of Dynkin diagram vertices is invariant under all automorphisms of the Dynkin diagram. Hence the Weyl group element of the maximal length interchanges P and P^- . We have to find $\text{gtd}(L : \mathfrak{u}^-)$.

The Levi subgroup L is locally isomorphic to $(\mathbb{K}^*)^2 \times SO_8$, and the L -module \mathfrak{u}^- is isomorphic to $V_1 \oplus V_2 \oplus V_3$. Here V_1 is an SO_8 -module with the lowest weight $-\pi_1$, i. e. a tautological SO_8 -module, V_2 is an SO_8 -module with the lowest weight $-\pi_3$, V_3 is an SO_8 -module with the lowest weight $-\pi_4$. Denote the components of $u \in \mathfrak{u}^-$ by u_1, u_2, u_3 .

Since V_1 is a tautological SO_8 -module, there exists an SO_8 -invariant symmetric bilinear form on it that we denote by (u_1, u_1) . There exist diagram automorphisms of SO_8 that transform the tautological SO_8 -module to SO_8 -modules isomorphic to V_2 and V_3 . So there exist an SO_8 -invariant on V_2 that we denote by (u_2, u_2) and an SO_8 -invariant on V_3 that we denote by (u_3, u_3) . These bilinear forms are not necessarily $(\mathbb{K}^*)^2$ -invariant, in general their $(\mathbb{K}^*)^2$ -weights are three pairs of integers. There is a linear combination of these pairs that is equal to zero. Hence, there exists a non-trivial rational L -invariant of the form

$$(u_1, u_1)^a (u_2, u_2)^b (u_3, u_3)^c,$$

where $a, b, c \in \mathbb{Z}$, and the G -action on $G/P \times G/P \times G/P$ is not generically transitive.

3. FINITE NUMBER OF ORBITS

Proposition 2. *Let G be a simple algebraic group and P be a proper parabolic subgroup. If $n \geq 4$, the number of G -orbits on $(G/P)^n$ is infinite.*

Proof. Let $P = P_{i_1, \dots, i_s}$. Consider the dominant weight $\lambda = \pi_{i_1} + \dots + \pi_{i_s}$. Then G/P is isomorphic to the projectivization of the orbit of the highest weight vector $v_\lambda \in V(\lambda)$. In the sequel we shortly write $i = i_1$. It is easy to check that $y_i^2 v_\lambda = 0$. Denote the unipotent subgroup $\exp(\mathfrak{t}y_i)$ by U_i . We see that $U_i v_\lambda$ is an affine line not containing zero. The closure of its image in the projectivization $\mathbf{P}(V(\lambda))$ is a projective line $\mathbf{P}^1 \subseteq G/P \subseteq \mathbf{P}(V(\lambda))$. Choose $n \geq 4$ points $(x_1, \dots, x_n) \in \mathbf{P}^1 \times \dots \times \mathbf{P}^1 \subseteq G/P \times \dots \times G/P$. The double ratio of the first four of these points does not change under G -action. Hence, two n -tuples with different double ratios cannot belong to the same orbit, and the number of orbits is infinite. \square

Now we prove that in the cases $P = P_{1,l}$ and $P = P_{l-1,l}$ the number of orbits on $G/P \times G/P \times G/P$ is infinite.

We suppose that $G = SO_{2l}$. Let \mathbb{K}^{2l} be the tautological SO_{2l} -module and let e_1, \dots, e_{2l} be the standard basis. Let $X' \subset \text{Gr}(l, 2l)$ be the set of all isotropic subspaces of dimension l in \mathbb{K}^{2l} . It is known that G/P_l is isomorphic to a connected component of X' . In the sequel we suppose that $G/P_l = X \subseteq X'$. For each $s \in X$ let $Y_s \subset \mathbf{P}^{2l-1}$ be the set of all lines contained in s . One easily checks that the closed subset $Y = \cup_{s \in X} (s \times Y_s) \subset \text{Gr}(l, 2l) \times \mathbf{P}^{2l-1}$ is isomorphic to $SO_{2l}/P_{1,l}$.

Similarly, if $s \in X$, denote by $Z_s \subset \text{Gr}(l-1, 2l)$ the set of all subspaces of dimension $l-1$ in s . Let Z be the closed subset $\cup_{s \in X} (s \times Z_s) \subset \text{Gr}(l, 2l) \times \text{Gr}(l-1, 2l)$. One easily checks that it is isomorphic to $SO_{2l}/P_{l-1,l}$.

First, let $l = 3$. Consider the following isotropic subspaces: $S_1 = \langle e_1, e_2, e_4 \rangle$, $S_2 = \langle e_2, e_3, e_6 \rangle$, $S_3 = \langle e_1, e_3, e_5 \rangle$. They belong to the same SO_6 -orbit, so we may suppose that $S_1, S_2, S_3 \in X$. Choose a line $T_1 \subset S_1$ such that $T_1 \subset \langle e_1, e_2 \rangle$. Also choose lines $T_2 \subset S_2$ and $T_3 \subset S_3$ such that $T_2 \subset \langle e_2, e_3 \rangle$ and $T_3 \subset \langle e_1, e_3 \rangle$. Impose one more restriction, namely, the sum $T_2 + T_3$ should be direct and should not be equal to $\langle e_1, e_2 \rangle$. Consider the point $((S_1, T_1), (S_2, T_2), (S_3, T_3)) \in G/P_{1,l} \times G/P_{1,l} \times G/P_{1,l}$. There are four subspaces of $\langle e_1, e_2 \rangle$: $\langle e_1 \rangle = S_1 \cap S_3$, $\langle e_2 \rangle = S_2 \cap S_3$, T_1 and $T_4 = (T_2 \oplus T_3) \cap \langle e_1, e_2 \rangle$. Thus, we have defined four lines in \mathbb{K}^6 in terms of intersections and sums of S_i and T_i . If we apply an element $g \in G$ to these four lines, we will obtain four lines defined in the same way using gS_i and gT_i instead of S_i and T_i . The double ratio of these four lines in their sum of dimension two is not changed under G -action. Since T_1 is chosen arbitrarily, this double ratio can be any number and the number of orbits is infinite.

Consider the same subspaces S_i and T_i and set $U_1 = T_1 \oplus \langle e_4 \rangle$, $U_2 = T_2 \oplus \langle e_6 \rangle$ and $U_3 = T_3 \oplus \langle e_5 \rangle$. The point $((S_1, U_1), (S_2, U_2), (S_3, U_3))$ belongs to $Z \times Z \times Z$. Note that $\langle e_1, e_2, e_3 \rangle = (S_1 \cap S_2) \oplus (S_2 \cap S_3) \oplus (S_1 \cap S_3)$ and $T_i = U_i \cap \langle e_1, e_2, e_3 \rangle$. Again we have a subspace of dimension two and four lines in it defined in terms of intersections and sums of S_i and U_i . The existence of SO_6 -invariant double ratio in this case yields that the number of orbits is infinite.

Let $l > 3$. Construct the subspaces S_i , T_i and U_i as above, using the last three basis vectors instead of e_4, e_5, e_6 . Let $S'_i = S_i \oplus \langle e_4, \dots, e_l \rangle$ and $U'_i = U_i \oplus \langle e_4, \dots, e_l \rangle$. The points $((S'_1, T_1), (S'_2, T_2), (S'_3, T_3))$ and $((S'_1, U'_1), (S'_2, U'_2), (S'_3, U'_3))$ belong to $Y \times Y \times Y$ and $Z \times Z \times Z$, respectively. Consider also the subspace $V = (S'_1 \cap S'_2 \cap S'_3)^\perp$. The restriction of the bilinear form to this subspace is degenerate, its kernel is $S'_1 \cap S'_2 \cap S'_3 = \langle e_4, \dots, e_l \rangle$. The quotient is a space of dimension 6 with a bilinear form. The quotient morphism restricted to $\langle e_1, e_2, e_3, e_{2l-2}, e_{2l-1}, e_{2l} \rangle$ is an isomorphism, so we have subspaces S_i , T_i , U_i in the 6-dimensional space. This is exactly the same situation as we had above for the group SO_6 , and it enables us to define double ratios for the points of $G/P_{1,l} \times G/P_{1,l} \times G/P_{1,l}$ and $G/P_{l-1,l} \times G/P_{l-1,l} \times G/P_{l-1,l}$ under consideration. Therefore the number of SO_{2l} -orbits on these multiple flag varieties is infinite. This finishes the proof of Theorem 3.

REFERENCES

- [1] M. Brion, *Quelques propriétés des espaces homogènes sphériques*, Manuscripta Math. **55** (1986), no. 2, 191–198.
- [2] J. Humphreys, *Linear algebraic groups*. GTM **21** Springer-Verlag, New York-Heidelberg, 1975.
- [3] J. Humphreys, *Introduction to Lie Algebras and Representation Theory*. GTM **9**, Springer-Verlag, New York-Berlin, 1978.
- [4] P. Littleman, *On spherical double cones*, J. Algebra **166** (1994), no. 1, 142–157.
- [5] P. Magyar, J. Weynman, A. Zelevinsky, *Multiple flag varieties of finite type*, Adv. Math. **141** (1999), no. 1, 97–118.
- [6] P. Magyar, J. Weynman, A. Zelevinsky, *Symplectic multiple flag varieties of finite type*, J. Algebra **230** (2000), no. 1, 245–265.
- [7] V.L. Popov, *Generically multiple transitive algebraic group actions*, Algebraic groups and homogeneous spaces, 481–523, Tata Inst. Fund. Res. Stud. Math., Tata Inst. Fund. Res., Mumbai, 2007.
- [8] J. Stembridge, *Multiplicity-free products and restrictions of Weyl characters.*, Representation Theory **7** (2003), 404–439.
- [9] E.B. Vinberg, *Complexity of actions of reductive groups*, Funkt. Anal. i Pril. **20** (1986), no. 1, 1–13; English transl.: Funct. Anal. Appl. **20** (1986), no 1, 1–11.

CHAIR OF HIGHER ALGEBRA, FACULTY OF MECHANICS AND MATHEMATICS, LOMONOSOV MOSCOW STATE UNIVERSITY, LENINSKIE GORY 1, GSP-1, MOSCOW, 119991, RUSSIA

E-mail address: deviatov@mccme.ru