

## SINGULARITIES OF DIVISORS ON FLAG VARIETIES VIA HWANG’S PRODUCT THEOREM

EVGENY SMIRNOV

ABSTRACT. We give an alternative proof of a recent result by B. Pasquier stating that for a generalized flag variety  $X = G/P$  and an effective  $\mathbb{Q}$ -divisor  $D$  stable with respect to a Borel subgroup the pair  $(X, D)$  is Kawamata log terminal if and only if  $\lfloor D \rfloor = 0$ .

### 1. Introduction

Let  $G$  be a connected reductive algebraic group over  $\mathbb{C}$ . Recall that a horospherical  $G$ -variety  $X$  is a normal  $G$ -variety with an open  $G$ -orbit isomorphic to a torus fibration  $G/H$  over a flag variety  $G/P$ , where  $P$  is a parabolic subgroup in  $G$  and  $P/H$  is a torus. In [7] Boris Pasquier shows that for a horospherical variety  $X$  and an effective  $\mathbb{Q}$ -divisor  $D$  stable with respect to a Borel subgroup the pair  $(X, D)$  is Kawamata log terminal if and only if  $\lfloor D \rfloor = 0$ .

An essential part of the proof is the case when  $X$  itself is a flag variety  $G/P$ . In this case Pasquier uses a Bott–Samelson resolution to provide an explicit log resolution of the pair  $(G/P, D)$ , and to check that this pair is Kawamata log terminal using rather heavy combinatorics related to root systems. Further on, he uses the latter resolution to provide a log resolution of a general horospherical pair, and he uses Kawamata log terminality of  $(G/P, D)$  to establish the same result in general.

The main purpose of this note is to prove a similar result for a variety  $G/P$  avoiding explicit log resolutions, and instead using the Product Theorem for log canonical thresholds due to Jun-Muk Hwang, see [5]. In particular, we will not assume that the  $\mathbb{Q}$ -divisor  $D$  is stable under the action of the Borel subgroup. Note that while this approach does not allow one to get rid of Bott–Samelson resolutions because they are needed for the case of general horospherical varieties, it does allow to avoid the computations from [7, §5] and to replace them by easier computations of Proposition 4.4 below.

---

Received September 19, 2016; Accepted February 13, 2017.

2010 *Mathematics Subject Classification*. **Please supply MSC numbers.**

*Key words and phrases*. **Please supply Keywords.**

The study was partially supported by the Russian Academic Excellence Project “5–100”, Simons–IUM fellowship (§1–3), and RSF grant, project 14-21-00053 dated 11.08.14 (§4).

The plan of the paper is as follows. In §2 we recall definitions and properties of log canonical thresholds. In §3 we recall the basic facts about geometry of flag varieties and give a precise statement of our main result, which is Theorem 3.2. In §4 we introduce more notation and prove the main theorem.

I am grateful to I. Cheltsov, Yu. Prokhorov and especially C. Shramov for very useful discussions and to M. Brion and B. Pasquier for valuable comments.

## 2. Log canonical thresholds

In this section we recall definitions and some properties of log canonical thresholds. We refer a reader to [6, §8] for (much) more details.

Let  $X$  be a smooth complex algebraic variety,  $D$  be an effective  $\mathbb{Q}$ -divisor. Choose a point  $x \in X$ .

**Definition 2.1.** If  $D$  is a Cartier divisor locally defined by the equation  $f = 0$ , then the log canonical threshold  $\text{lct}_x(D)$  of  $D$  near  $x$  is defined by

$$\text{lct}_x(D) = \sup \left\{ c > 0 \mid \frac{1}{|f|^c} \in L_{loc}^2 \right\}.$$

In particular, if  $x$  is not contained in the support of  $D$ , we put  $\text{lct}_x(D) = +\infty$ .

If  $D$  is an arbitrary  $\mathbb{Q}$ -divisor, then we define  $\text{lct}_x(D) = \text{lct}_x(rD)/r$  for sufficiently divisible integer  $r$ . The log canonical threshold  $\text{lct}(X, D)$  of  $D$  is defined as the infimum of  $\text{lct}_x(D)$  over all  $x \in X$ .

**Definition 2.2.** The pair  $(X, D)$  is said to be *Kawamata log terminal* if the inequality  $\text{lct}(X, D) > 1$  holds, and *log canonical* if the inequality  $\text{lct}(X, D) \geq 1$  holds.

Let  $L \in \text{Pic}(X)$  be a line bundle such that the linear system  $|L|$  is non-empty. We define  $\text{lct}(X, L)$  as the infimum  $\inf_{\Delta \in |L|} \text{lct}(X, \Delta)$ . The following theorem is taken from [5].

**Theorem 2.3** (see [5, §2]). *Let  $f: X \rightarrow Y$  be a smooth projective morphism between two smooth projective varieties,  $y \in Y$ , and  $X_y = f^{-1}(y)$  be the fiber over  $y$ . Let  $D$  be an effective divisor on  $X$ , and let  $L$  be the restriction of  $\mathcal{O}_X(D)$  to  $X_y$ . Then one of the following holds:*

- (i) *either  $\text{lct}_x(D) \geq \text{lct}(X_y, L)$  for each  $x \in X_y$ ,*
- (ii) *or  $\text{lct}_{x_1}(D) = \text{lct}_{x_2}(D)$  for any two points  $x_1, x_2 \in X_y$ .*

One can define the global log canonical threshold

$$\text{lct}(X) = \inf \{ \text{lct}(X, \Delta) \mid \Delta \sim_{\mathbb{Q}} -K_X \text{ is an effective } \mathbb{Q}\text{-divisor} \},$$

where  $-K_X$  is the anticanonical class of  $X$ . This definition makes sense if some positive multiple of  $-K_X$  is effective; for example, this holds for Fano varieties and for spherical varieties (see [1, §4]). For more properties of  $\text{lct}(X)$  see [3]; for its relation to the  $\alpha$ -invariant of Tian see [3, Appendix A].

### 3. Flag varieties

Let  $G$  be a connected reductive algebraic group. We fix a Borel subgroup  $B$  in  $G$  and a maximal torus  $T \subset B$ . Denote by  $R$  the root system of  $G$ , and by  $S \subset R$  the set of simple roots in  $R$ , where the positive roots are the roots of  $(B, T)$ . The Weyl group of  $R$  will be denoted by  $W$ ; let  $\ell: W \rightarrow \mathbb{Z}_{\geq 0}$  be the length function on  $W$ .

Let  $P \supset B$  be a parabolic subgroup in  $G$ . Then  $G/P$  is a (generalized) partial flag variety. Denote by  $I$  the set of simple roots of the Levi subgroup of  $P$ ; in particular, for  $P = B$  we have  $I = \emptyset$  and for  $P$  maximal the set  $I$  is obtained from  $S$  by removing exactly one simple root. For example, if  $G = \mathrm{SL}_n(\mathbb{C})$ , then for the maximal parabolic subgroup  $P$  corresponding to  $S \setminus \{\alpha_k\}$ ,  $1 \leq k \leq n-1$ , the homogeneous space  $G/P$  is the Grassmannian  $\mathrm{Gr}(k, n)$ .

For a subset  $I \subset S$  of the set of simple roots, let  $W_P \subset W$  be the subgroup of  $W$  generated by the simple reflections  $s_\alpha$ , where  $\alpha \in I$ . In each left coset from  $W/W_P$  there exists a unique element of minimal length. Denote the set of such elements by  $W^P$ ; we will identify it with  $W/W_P$ . It is well-known (see, for instance, [2, §1.2]) that the partial flag variety  $G/P$  admits a Schubert decomposition into orbits of  $B$ , and the orbits are indexed by the elements of  $W^P$ :

$$G/P = \bigsqcup_{w \in W^P} BwP/P.$$

Moreover, the dimension of the cell  $BwP/P$  equals the length of  $w$ . The closures of these cells are called *Schubert varieties*; we denote them by  $Y_w = \overline{BwP/P}$ .

Denote by  $w_0$  and  $w_0^P$  the longest elements in  $W$  and  $W_P$ , respectively. Then the length  $\ell(w_0 w_0^P)$  is the dimension of  $G/P$ . We shall also need the *opposite Schubert varieties*

$$Y^w = \overline{w_0 B w_0 w P / P} = w_0 Y_{w_0 w}.$$

If  $w \in W^P$ , then  $w_0 w w_0^P$  also belongs to  $W^P$  (i.e., is the shortest representative in its left coset  $w_0 w W_P = w_0 w w_0^P W_P$ ); in this case  $\dim Y_w = \ell(w)$  and

$$\dim Y^w = \ell(w_0 w_0^P) - \ell(w).$$

From the definition of  $Y_w$  and  $Y^w$  we readily see that the cohomology classes  $[Y_w]$  and  $[Y^{w_0 w w_0^P}]$  in  $H^*(G/P, \mathbb{Z})$  are equal.

Irreducible  $B$ -stable divisors of  $G/P$  are the Schubert varieties of codimension 1. Denote them by

$$D_\alpha = \overline{B w_0 s_\alpha w_0^P P / P} = w_0 Y^{s_\alpha}.$$

The following proposition is a standard fact on Schubert varieties (cf. [2, §1.4]).

- Proposition 3.1.** (i) *The divisors  $D_\alpha$  for  $\alpha \notin I$  freely generate  $\text{Pic}(G/P)$ , so one has  $\text{rk Pic}(G/P) = |S \setminus I|$ . In particular, for  $P$  maximal one has  $\text{Pic}(G/P) \cong \mathbb{Z}$ , and there is a unique  $B$ -stable prime divisor.*
- (ii) *The classes of Schubert varieties  $[Y_w] \in H^\bullet(G/P, \mathbb{Z})$  freely generate  $H^\bullet(G/P, \mathbb{Z})$  as an abelian group. The elements of this basis are Poincaré dual to the classes of the corresponding opposite Schubert varieties: if  $w, v \in W^P$  and  $\ell(w) = \ell(v)$ , then*

$$[Y^w] \smile [Y_v] = \delta_{wv} \quad \text{for each } w, v \in W.$$

*In particular, the classes of one-dimensional Schubert varieties, that is, of  $B$ -stable curves  $\overline{Bs_\alpha P/P} \subset G/P$ , are dual to the classes of divisors:*

$$D_\alpha \smile [\overline{Bs_\beta P/P}] = \delta_{\alpha\beta} \quad \text{for each } \alpha, \beta \in S \setminus I.$$

The purpose of our paper is to give a new proof of the following result.

**Theorem 3.2** (see [7, Theorem 3.1]). *Let  $D \sim \sum a_\alpha D_\alpha$ , where  $a_\alpha$  are non-negative rational numbers, be an effective non-zero  $\mathbb{Q}$ -divisor. Then*

$$\text{lct}(G/P, D) \geq \frac{1}{\max a_\alpha}.$$

*In particular, the pair  $(G/P, D)$  is Kawamata log terminal provided that all  $a_\alpha$  are less than 1.*

*Remark 3.3.* One can show that every effective divisor on  $G/P$  is linearly equivalent to an effective  $B$ -stable divisor. In other words, the classes of divisors  $D_\alpha$  in the  $\mathbb{Q}$ -vector space  $\text{Pic}(G/P) \otimes \mathbb{Q}$  span the cone of effective divisors. Thus the assumption of Theorem 3.2 requiring that  $a_\alpha$  are non-negative is implied by effectiveness of  $D$ ; we keep it just to make the assertion more transparent.

*Remark 3.4.* If the  $\mathbb{Q}$ -divisor  $D$  of Theorem 3.2 is  $B$ -stable, then in addition to the inequality given by Theorem 3.2 we have an obvious opposite inequality, because  $D_\alpha$  is an effective divisor. Therefore, in this case we recover the equality given by [7, Theorem 3.1].

A by-product of Theorem 3.2 is the following assertion on global log canonical thresholds of complete flag varieties that is well known to experts.

**Corollary 3.5.** *One has  $\text{lct}(G/B) = 1/2$ .*

*Proof.* One has  $-K_{G/B} \sim \sum_{\alpha \in S} 2D_\alpha$ . Thus  $\text{lct}(G/B) \geq 1/2$  by Theorem 3.2. The opposite inequality is implied by the fact that the divisor  $D_\alpha$  is effective.  $\square$

#### 4. Proof of the main theorem

In this section we prove Theorem 3.2.

Fix a simple root  $\alpha \in S \setminus I$ . Let  $J = I \cup \{\alpha\}$ , and let  $P'$  be the parabolic subgroup corresponding to  $J$ ; then  $P \subset P'$ . There is a  $G$ -equivariant fibration  $\pi_\alpha: G/P \rightarrow G/P'$ .

Let  $X_\alpha$  be a fiber of this fibration. Consider the Dynkin diagram of  $G$ ; its vertices correspond to simple roots from  $S$ . Let  $\bar{J}$  be the connected component containing  $\alpha$  of the subgraph spanned by the vertices of  $J$ . This component is the Dynkin diagram of a connected simple algebraic group  $\bar{G}$ . Let  $\bar{P}_\alpha$  be a maximal parabolic subgroup of  $\bar{G}$  with the set of roots  $\bar{J} \setminus \{\alpha\}$ . Then  $X_\alpha$  is isomorphic to the  $\bar{G}$ -homogeneous space  $\bar{G}/\bar{P}_\alpha \cong P'/P$ .

**Example 4.1.** Let  $G = \mathrm{SL}_n(\mathbb{C})$ . Its Dynkin diagram is  $A_{n-1}$ ; denote its simple roots by  $\alpha_1, \dots, \alpha_{n-1}$ . Put  $I = S \setminus \{\alpha_{d_1}, \dots, \alpha_{d_r}\}$ , where  $1 \leq d_1 < \dots < d_r \leq n-1$ . We also formally set  $d_0 = 0$  and  $d_{r+1} = n$ . Then  $G/P$  is a partial flag variety

$$\mathrm{Fl}(d_1, \dots, d_r) \cong \{U_1 \subset \dots \subset U_r \subset \mathbb{C}^n \mid \dim U_j = d_j\}.$$

Let  $\alpha = \alpha_{d_s}$ , where  $1 \leq s \leq r$ , and  $J = I \cup \{\alpha\}$ . Then  $G/P'$  is an  $(r-1)$ -step flag variety  $\mathrm{Fl}(d_1, \dots, \widehat{d_s}, \dots, d_r)$ , and the map  $\pi_\alpha: G/P \rightarrow G/P'$  is given by forgetting the  $s$ -th component of each flag. The fibers of this projection are isomorphic to the Grassmannian  $\mathrm{Gr}(d_s - d_{s-1}, d_{s+1} - d_{s-1})$ .

According to Proposition 3.1(i), since  $\bar{P}_\alpha \subset \bar{G}$  is a maximal parabolic subgroup, one has  $\mathrm{Pic} X_\alpha \cong \mathbb{Z}$ . Let  $H_\alpha$  be the ample generator of  $\mathrm{Pic} X_\alpha$ .

**Theorem 4.2** ([5, Theorem 2]; see also [4]). *Let  $k$  be a positive integer. Then one has  $\mathrm{lct}(X_\alpha, kH_\alpha) = 1/k$ .*

*Remark 4.3.* The assertion of [5, Theorem 2] is that the inequality  $\mathrm{lct}(X_\alpha, kH_\alpha) \geq 1/k$  holds. This is equivalent to Theorem 4.2 since the linear system  $|H_\alpha|$  is always non-empty by Proposition 3.1(i).

The following computation will be the central point of our proof of Theorem 3.2.

**Proposition 4.4.** *For each  $\beta \in S \setminus I$  one has*

$$D_\alpha|_{X_\beta} \sim \begin{cases} H_\alpha & \text{if } \alpha = \beta; \\ 0 & \text{otherwise.} \end{cases}$$

*Proof.* As we discussed above, the fiber  $X_\beta$  can be identified with the variety  $P'/P \subset G/P$ . It is a flag variety with the Picard group of rank one; since the classes of  $B$ -stable curves are dual to the classes of ( $B$ -stable) divisors (see Proposition 3.1(ii)),  $X_\beta$  contains a unique  $B$ -stable curve. This curve has the form  $\overline{Bs_\beta P/P} \subset P'/P \cong X_\beta$ . Its class in  $H^\bullet(X_\beta, \mathbb{Z})$  is Poincaré dual to the ample generator  $H_\beta$  of  $\mathrm{Pic}(X_\beta, \mathbb{Z})$ .

At the same time, as it was stated in Proposition 3.1(ii), the intersection of  $\overline{Bs_\beta P/P}$  with  $D_\alpha$  equals the class of a point if  $\alpha = \beta$  and zero otherwise.  $\square$

*Proof of Theorem 3.2.* Replacing  $D$  by its appropriate multiple, we may assume that it is a Cartier divisor, not just a  $\mathbb{Q}$ -divisor. Put  $a = \max a_\alpha$ . Suppose that  $\mathrm{lct}(X, D) < 1/a$ .

Pick a point  $x \in X$  such that  $\text{lct}_x(D) < 1/a$ . Choose an index  $\alpha$  from  $S \setminus I$ , and let  $[D|_{X_\alpha}]$  be the class of the restriction of  $\mathcal{O}_X(D)$  to  $X_\alpha$  in  $\text{Pic}(X_\alpha)$ . According to Proposition 4.4, one has  $[D|_{X_\alpha}] \sim a_\alpha H_\alpha$ . Theorem 4.2 implies that

$$\text{lct}(X_\alpha, [D|_{X_\alpha}]) \geq \frac{1}{a_\alpha} \geq \frac{1}{a};$$

this trivially includes the case when  $a_\alpha = 0$ . Without loss of generality we can suppose that the fiber  $X_\alpha$  passes through the point  $x$ . The above means that the alternative (i) in Theorem 2.3 never holds.

Let  $\alpha_1, \dots, \alpha_r$  be the simple roots from  $S \setminus I$ . Let  $\tilde{X}_1$  be the fiber of  $\pi_{\alpha_1}$  passing through the point  $x$ . For each  $i = 2, \dots, r$  let  $\tilde{X}_i$  be the union of all fibers of  $\pi_{\alpha_i}$  passing through the points of  $\tilde{X}_{i-1}$ . In particular, one has  $\tilde{X}_r = X$ .

First apply Theorem 2.3 to the point  $x$  and the fibration  $\pi_{\alpha_1}$ . It implies that for each  $x_1 \in \tilde{X}_1$  we have  $\text{lct}_{x_1}(D) < 1/a$ . Now apply it to each point  $x_1 \in \tilde{X}_1$  and the fibration  $\pi_{\alpha_2}$ . We see that for each  $x_2 \in \tilde{X}_2$  the inequality  $\text{lct}_{x_2}(D) < 1/a$  holds. Proceeding by induction, we obtain the same inequality for every point in  $\tilde{X}_r = X$ . In particular, each point of  $X$  is contained in the support of  $D$ , which is a contradiction.  $\square$

## References

- [1] M. Brion, *Curves and divisors in spherical varieties*, In Algebraic groups and Lie groups, volume 9 of Austral. Math. Soc. Lect. Ser., pages 21–34. Cambridge Univ. Press, Cambridge, 1997.
- [2] ———, *Lectures on the geometry of flag varieties*, In Topics in cohomological studies of algebraic varieties, Trends Math., pages 33–85. Birkhäuser, Basel, 2005.
- [3] I. A. Chel'tsov and K. A. Shramov, *Log-canonical thresholds for nonsingular Fano threefolds*, Uspekhi Mat. Nauk **63** (2008), no. 5, 73–180.
- [4] J.-M. Hwang, *Log canonical thresholds of divisors on Grassmannians*, Math. Ann. **334** (2006), no. 2, 413–418.
- [5] ———, *Log canonical thresholds of divisors on Fano manifolds of Picard number 1*, Compos. Math. **143** (2007), no. 1, 89–94.
- [6] J. Kollár, *Singularities of pairs*, In Algebraic geometry—Santa Cruz 1995, volume 62 of Proc. Sympos. Pure Math., pages 221–287. Amer. Math. Soc., Providence, RI, 1997.
- [7] B. Pasquier, *Klt singularities of horospherical pairs*, arXiv:1509.06502 (2015), to appear in Annales de l'Institut Fourier, 2016.

EVGENY SMIRNOV  
 FACULTY OF MATHEMATICS AND  
 LABORATORY OF ALGEBRAIC GEOMETRY AND ITS APPLICATIONS  
 NATIONAL RESEARCH UNIVERSITY HIGHER SCHOOL OF ECONOMICS  
 USACHEVA STR., 6, MOSCOW 119048, RUSSIA  
 AND  
 INDEPENDENT UNIVERSITY OF MOSCOW  
 BOLSHOI VLASSIEVSKII PER., 11, MOSCOW 119002, RUSSIA  
*E-mail address:* `esmirnov@hse.ru`