




**NINTH INTERNATIONAL CONFERENCE ON
VORTEX MATTER IN NANOSTRUCTURED
SUPERCONDUCTORS**

**Rhodes (Greece)
12-17 September, 2015**

ABSTRACT BOOK



**NANOSCALE SUPERCONDUCTIVITY,
FLUXONICS AND PHOTONICS:
ADDRESSING GRAND CHALLENGES**
KU LEUVEN METHUSALEM GROUP

EDITED BY :

**Johan Vanacken
Jeroen Scheerder
Matias Timmermans
Monique Van Meerbeek**

Quantum Phase Slip in a Superconductor

K. Yu. Arutyunov^{1,2}

¹ *Moscow Institute of Electronics and Mathematics at
National Research University Higher School of Economics, 101000, Moscow, Russia*

² *Kapitza Institute of Physical Problems, Moscow, 119334, Russia*

The topic of quantum fluctuations in quasi-1D superconductors, also called quantum phase slips (QPS), has recently attracted the significant attention. It has been shown that the phenomenon is capable to suppress the zero resistivity of ultra-narrow superconducting nanowires at low temperatures $T \ll T_c$ and quench persistent currents in tiny nanorings. The coherent QPS effect enables fabrication of the new generation of quantum logic devices – qubits. It has been predicted that a superconducting nanowire in the regime of QPS is dual to a Josephson junction. In particular case of an extremely narrow superconducting nanowire imbedded in high-impedance environment the duality leads to an intuitively controversial result: the superconductor should enter an insulating state.

Here we experimentally demonstrate that the I-V characteristic of such a wire indeed shows Coulomb blockade which disappears with application of a critical magnetic field and/or above the critical temperature proving that the effect is related to superconductivity. Such system can be considered as a junctionless single electron transistor (with charge $2e$), where the QPS provide the dynamic equivalent of weak links in conventional devices containing static (in space and time) tunnel junctions. Application of external RF radiation can be synchronized with the internal Bloch oscillations of charge. The phenomenon is dual to the well-known Shapiro effect: the voltage steps for a Josephson junction are substituted by the current steps for a QPS wire: the proof-of-principle demonstration of the long-awaited metrological application - the quantum standard of electric current.

We will also discuss our latest results demonstrating finite noise below the ‘residual’ critical current and temperature due to the same QPS effect. In ultra-narrow nanowires the quantum fluctuations of the amplitude of the order parameter result in smearing of the energy gap edge, which has been measured in tunneling experiments.

Presenting Author: K. Yu. Arutyunov

Email: karutyunov@hse.ru