

**TRIANGULATED ENDOFUNCTORS OF THE DERIVED
CATEGORY OF COHERENT SHEAVES WHICH DO NOT ADMIT
DG LIFTINGS**

VADIM VOLOGODSKY

ABSTRACT. In [RV], Rizzardo and Van den Bergh constructed an example of a triangulated functor between the derived categories of coherent sheaves on smooth projective varieties over a field k of characteristic 0 which is not of the Fourier-Mukai type. The purpose of this note is to show that if $\text{char } k = p$ then there are very simple examples of such functors. Namely, for a smooth projective Y over \mathbb{Z}_p with the special fiber $i : X \hookrightarrow Y$, we consider the functor $Li^* \circ i_* : D^b(X) \rightarrow D^b(X)$ from the derived categories of coherent sheaves on X to itself. We show that if Y is a flag variety which is not isomorphic to \mathbb{P}^1 then $Li^* \circ i_*$ is not of the Fourier-Mukai type. Note that by a theorem of Toen ([T], Theorem 8.15) the latter assertion is equivalent to saying that $Li^* \circ i_*$ does not admit a lifting to a \mathbb{F}_p -linear DG quasi-functor $D_{dg}^b(X) \rightarrow D_{dg}^b(X)$, where $D_{dg}^b(X)$ is a (unique) DG enhancement of $D^b(X)$. However, essentially by definition, $Li^* \circ i_*$ lifts to a \mathbb{Z}_p -linear DG quasi-functor.

Given smooth proper schemes X_1, X_2 over a field k and an object $E \in D^b(X_1 \times X_2)$ of the bounded derived category of coherent sheaves on $X_1 \times X_2$ define a triangulated functor

$$(0.1) \quad \Phi_E : D^b(X_1) \rightarrow D^b(X_2)$$

sending a bounded complex M of coherent sheaves on X_1 to $Rp_{2*}(E \otimes^L p_1^* M)$, where $p_i : X_1 \times X_2 \rightarrow X_i$ are the projections. Recall that a triangulated functor $D^b(X_1) \rightarrow D^b(X_2)$ is said to be of the Fourier-Mukai type if it is isomorphic to Φ_E for some E .

Let Y be a smooth projective scheme over $\text{Spec } \mathbb{Z}_p$ and let X be its special fiber, $i : X \hookrightarrow Y$ the closed embedding. Consider the triangulated functor $G : D^b(X) \rightarrow D^b(X)$ given by the formula

$$G = Li^* \circ i_*$$

We shall see that in general G is not of the Fourier-Mukai type.

Theorem 1. *Let Z a smooth projective scheme over $\text{Spec } \mathbb{Z}_p$, $Y = Z \times Z$, $X = Y \times_{\text{Spec } \mathbb{Z}_p} \text{Spec } \mathbb{F}_p$. Assume that*

- (1) *The Frobenius morphism $Fr : \overline{Z} \rightarrow \overline{Z}$, where $\overline{Z} = Z \times \text{Spec } \mathbb{F}_p$, does not lift modulo p^2 .*
- (2) *$H^1(X, T_X) = 0$, where T_X is the tangent sheaf on X .*

Then $G = Li^ \circ i_* : D^b(X) \rightarrow D^b(X)$ is not of the Fourier-Mukai type.*

For example, let GL_n be the general linear group over $\text{Spec } \mathbb{Z}_p$, $B \subset GL_n$ a Borel subgroup. Then, by Theorem 6 from [BTLM], for any $n > 2$, the flag variety $Z = GL_n/B$ satisfies the first assumption of the Theorem *i.e.*, the Frobenius $Fr : \overline{Z} \rightarrow \overline{Z}$ does not lift on $Z \times \text{Spec } \mathbb{Z}/p^2\mathbb{Z}$. By ([KLT], Theorem 2), we have that $H^1(\overline{Z}, T_{\overline{Z}}) =$

$H^1(\overline{\mathbb{Z}}, \mathcal{O}_{\overline{\mathbb{Z}}}) = 0$. It follows that $H^1(X, T_X) = 0$. Hence, by the Theorem, for $n > 2$, $G : D^b(X) \rightarrow D^b(X)$ is not of the Fourier-Mukai type.

Proof. Assume the contrary and let $E \in D^b(X \times X)$ be the Fourier-Mukai kernel. By definition, for every $M \in D^b(X)$ we have a functorial isomorphism

$$(0.2) \quad G(M) \xrightarrow{\sim} Rp_{2*}(E \overset{L}{\otimes} p_1^*M).$$

By the projection formula ([H], Chapter II, Prop. 5.6) we have that

$$i_* \circ Li^* \circ i_*(M) \xrightarrow{\sim} i_*(M) \overset{L}{\otimes} i_*(\mathcal{O}_X) \xrightarrow{\sim} i_*(M) \otimes (\mathcal{O}_Y \xrightarrow{p} \mathcal{O}_Y) \xrightarrow{\sim} i_*(M) \oplus i_*(M)[1]$$

In particular, if M is a coherent sheaf then $\underline{H}^i(G(M)) \simeq M$ for $i = 0, -1$ and $\underline{H}^i(G(M)) = 0$ otherwise. Applying this observation and formula (0.2) to skyscraper sheaves, $M = \delta_x$, $x \in X(\overline{\mathbb{F}}_p)$, we conclude that the coherent sheaves $\underline{H}^i(E)$ are set theoretically supported on the diagonal $\Delta_X \subset X \times X$. Applying the same formulas to $M = \mathcal{O}_X$ we see that $p_{2*}(\underline{H}^i(E)) = \mathcal{O}_X$ for $i = 0, -1$ and $p_{2*}(\underline{H}^i(E)) = 0$ otherwise. In fact, every coherent sheaf F on $X \times X$ which is set theoretically supported on the diagonal and such that $p_{2*}F = \mathcal{O}_X$ is isomorphic to \mathcal{O}_{Δ_X} . It follows that $\underline{H}^0(E) = \underline{H}^{-1}(E) = \mathcal{O}_{\Delta_X}$. In the other words, E fits into an exact triangle in $D^b(X \times X)$

$$(0.3) \quad \mathcal{O}_{\Delta_X}[1] \xrightarrow{\alpha} E \longrightarrow \mathcal{O}_{\Delta_X} \xrightarrow{\beta} \mathcal{O}_{\Delta_X}[2]$$

for some $\beta \in Ext_{\mathcal{O}_{X \times X}}^2(\mathcal{O}_{\Delta_X}, \mathcal{O}_{\Delta_X})$. We wish to show that the second assumption in the Theorem implies that $\beta = 0$, while the first one implies that $\beta \neq 0$. For every $M \in D^b(X)$, (0.3) gives rise to an exact triangle

$$(0.4) \quad M[1] \xrightarrow{\alpha_M} G(M) \longrightarrow M \xrightarrow{\beta_M} M[2]$$

Our main tool is the following result.

Lemma 0.1. *For a coherent sheaf M the following conditions are equivalent.*

- (1) $\beta_M = 0$.
- (2) $G(M) \xrightarrow{\sim} M \oplus M[1]$.
- (3) *There exists a morphism $\lambda : G(M) \rightarrow M[1]$ such that $\lambda \circ \alpha_M$ is an isomorphism.*
- (4) *M admits a lift modulo p^2 i.e., there is a coherent sheaf \tilde{M} on Y flat over $\mathbb{Z}/p^2\mathbb{Z}$ such that $i^*\tilde{M} \simeq M$.*

Proof. The equivalence of (1), (2) and (3) is immediate. Let us check that (3) is equivalent to (4). A morphism $\lambda : G(M) \rightarrow M[1]$ gives rise by adjunction a morphism $\gamma : i_*M \rightarrow i_*M[1]$. Note that $\tilde{M} = \text{cone } \gamma[1]$ is a coherent sheaf on Y which is an extension of i_*M by itself. It suffices to prove that $\lambda \circ \alpha_M : M[1] \rightarrow M[1]$ is an isomorphism if and only if \tilde{M} is flat over $\mathbb{Z}/p^2\mathbb{Z}$. Indeed, from the exact triangle

$$Li^*i_*M \rightarrow Li^*(\tilde{M}) \rightarrow Li^*i_*M \rightarrow Li^*i_*M[1]$$

we get a long exact sequence of the cohomology sheaves

$$0 \rightarrow M \rightarrow L_1i^*(\tilde{M}) \rightarrow M \xrightarrow{\lambda \circ \alpha_M} M \rightarrow i^*(\tilde{M}) \rightarrow M \rightarrow 0$$

Thus $\lambda \circ \alpha_M$ is an isomorphism if and only if in the exact sequence

$$0 \rightarrow i_*M \rightarrow \tilde{M} \rightarrow i_*M \rightarrow 0$$

the image of second map is the kernel of the multiplication by p on \tilde{M} and also the image of this map. The latter is equivalent to flatness of \tilde{M} over $\mathbb{Z}/p^2\mathbb{Z}$. \square

We have a spectral sequence converging to $Ext_{\mathcal{O}_{X \times X}}^*(\mathcal{O}_{\Delta_X}, \mathcal{O}_{\Delta_X})$ whose second page is $H^*(X, \mathcal{E}xt_{\mathcal{O}_{X \times X}}^*(\mathcal{O}_{\Delta_X}, \mathcal{O}_{\Delta_X}))$. In particular, we have a homomorphism

$$Ext_{\mathcal{O}_{X \times X}}^2(\mathcal{O}_{\Delta_X}, \mathcal{O}_{\Delta_X}) \rightarrow H^0(X, \mathcal{E}xt_{\mathcal{O}_{X \times X}}^2(\mathcal{O}_{\Delta_X}, \mathcal{O}_{\Delta_X})) \xrightarrow{\sim} H^0(X, \wedge^2 T_X).$$

Let us check that the image μ of β under this map is 0. To do this we apply the Lemma to skyscraper sheaves δ_x , where x runs over closed points of X . On the one hand, the evaluation of the bivector field μ at x is equal to the class of β_{δ_x} in $Ext_{\mathcal{O}_X}^2(\delta_x, \delta_x) \xrightarrow{\sim} \wedge^2 T_{x, X}$. On the other hand, by the Lemma, $\beta_{\delta_x} = 0$ since δ_x is liftable modulo p^2 . Next, the assumption that $H^1(X, T_X) = 0$ implies that β lies in the image of the map

$$(0.5) \quad v : H^2(X, \mathcal{O}_X) \xrightarrow{\sim} H^2(X, \mathcal{E}xt_{\mathcal{O}_{X \times X}}^0(\mathcal{O}_{\Delta_X}, \mathcal{O}_{\Delta_X})) \rightarrow Ext_{\mathcal{O}_{X \times X}}^2(\mathcal{O}_{\Delta_X}, \mathcal{O}_{\Delta_X}).$$

The map (0.5) has a left inverse $u : Ext_{\mathcal{O}_{X \times X}}^2(\mathcal{O}_{\Delta_X}, \mathcal{O}_{\Delta_X}) \rightarrow H^2(X, \mathcal{O}_X)$ which takes β to $\beta_{\mathcal{O}_X}$. But, by the Lemma, the later class is equal to 0 since \mathcal{O}_X is liftable modulo p^2 . It follows that β is 0.

On the other hand, let $\Gamma \subset X = \overline{Z} \times \overline{Z}$ be the graph of the Frobenius morphism $Fr : \overline{Z} \rightarrow \overline{Z}$ and \mathcal{O}_Γ the structure sheaf of Γ viewed as a coherent sheaf on X . Then, by our first assumption, the sheaf \mathcal{O}_Γ is not liftable modulo p^2 . Hence, by the Lemma, $\beta_{\mathcal{O}_\Gamma}$ is not 0. This contradiction completes the proof. \square

Acknowledgements. I would like to thank Alberto Canonaco and Paolo Stellari: their interest prompted writing this note. Also, I am grateful to Alexander Samokhin for stimulating discussions and references.

REFERENCES

- [BTLM] A. Buch, J. F. Thomsen, N. Lauritzen, and V. Mehta, *The Frobenius morphism on a toric variety*, Tohoku Math. J. (2), Volume 49, Number 3 (1997), 355-366.
- [CV] D. Calaque, M. Van den Bergh, *Hochschild cohomology and Atiyah classes*, Adv. Math. 224 (2010), no. 5, 1839–1889.
- [H] R. Hartshorne, *Residues and duality*, LNM 20 (1966).
- [KLT] S. Kumar, N. Lauritzen, J.F.Thomsen, *Frobenius splitting of cotangent bundles of flag varieties*, Invent. Math. 136 (1999), pp. 603-62
- [RV] A. Rizzardo, M. Van den Bergh, *An example of a non-Fourier?Mukai functor between derived categories of coherent sheaves*, arXiv:1410.4039.
- [T] B. Toën, *The homotopy theory of dg-categories and derived Morita theory*, Invent. Math. 167 (2007),

NATIONAL RESEARCH UNIVERSITY “HIGHER SCHOOL OF ECONOMICS” AND THE UNIVERSITY OF OREGON

E-mail address: vvologod@uoregon.edu