

Remarks on mass transportation minimizing expectation of a minimum of affine functions

Alexander V. Kolesnikov and Nikolay Lysenko *

Abstract

We study the Monge–Kantorovich problem with one-dimensional marginals μ and ν and the cost function $c = \min\{l_1, \dots, l_n\}$ that equals the minimum of a finite number n of affine functions l_i satisfying certain non-degeneracy assumptions. We prove that the problem is equivalent to a finite-dimensional extremal problem. More precisely, it is shown that the solution is concentrated on the union of n products $I_i \times J_i$, where $\{I_i\}$ and $\{J_i\}$ are partitions of the real line into unions of disjoint connected sets. The families of sets $\{I_i\}$ and $\{J_i\}$ have the following properties: 1) $c = l_i$ on $I_i \times J_i$, 2) $\{I_i\}, \{J_i\}$ is a couple of partitions solving an auxiliary n -dimensional extremal problem. The result is partially generalized to the case of more than two marginals.

Keywords: Monge–Kantorovich problem, concave cost functions.

1 Introduction

Suppose we are given a couple of probability distributions μ, ν on the real line that are assumed to be atomless and a Borel function $c: \mathbb{R}^2 \rightarrow \mathbb{R}$. Denote by $\Pi(\mu, \nu)$ the set of Borel probability measures on $\mathbb{R} \times \mathbb{R}$ with marginals μ, ν . Recall (see, e.g., [1], [2], and [8]) that a measure $\pi \in \Pi(\mu, \nu)$ is a solution to the Monge–Kantorovich problem if it gives the minimum to the functional

$$\pi \mapsto \int c d\pi$$

on $\Pi(\mu, \nu)$:

$$\int c(x, y) d\pi \rightarrow \min, \pi \in \Pi(\mu, \nu). \quad (1)$$

It is a classical and well-known fact that for a broad class of convex functions, such as, for instance, $c = h(|x - y|)$ with a strictly convex function h , any solution to (1) is concentrated on the graph of a non-decreasing function. The assumption of convexity of c is standard for many core results of the transportation theory. The case of the quadratic cost function $c = |x - y|^2$ is of particular interest.

In general, the Monge–Kantorovich problem with a concave cost c is harder. Remarkably, solutions to (1) with concave c have a completely different structure as compared to solutions for convex costs. For instance, the corresponding optimal transportation mapping need not exist

* This study was supported by the RFBR project 17-01-00662 and the DFG project RO 1195/12-1. The article was prepared within the framework of the Academic Fund Program at the National Research University Higher School of Economics (HSE) in 2017–2018 (Grant No 17-01-0102) and by the Russian Academic Excellence Project “5-100”.

even for a strictly concave cost c . The case of $c = h(|x - y|)$ with a strictly concave function h has been studied in [4], where a general result on the existence of optimal transportation mappings has been established (see more recent developments in [7]). An exact solution in the one-dimensional case for $c = h(|x - y|)$ has been obtained in [5]. An algorithm to solve the discrete transportation problem with a concave cost has been proposed in [3].

We study problem (1) for the cost

$$c = \min\{l_1, l_2, \dots, l_n\},$$

$$l_i = a_i x + b_i y + c_i.$$

This problem, yet quite specific, is of particular interest, since the minima of affine functions are dense in the set of all concave functions. It turns out that solutions have a nice and relatively simple structure provided that the functions l_i satisfy certain non-degeneracy assumption. In particular, problem (1) can be reduced to a finite-dimensional optimization problem. The corresponding optimal transportation problem admits a non-unique solution. Some of our results are generalized for the case of $m \geq 2$ one-dimensional marginals. In particular, we find a complete characterization of the solution for the cost function

$$c = \min\{x_1, x_2, \dots, x_m\}$$

(the minimum of coordinate functions).

We emphasize that nowadays the multi-marginal transportation problem (in particular, with one-dimensional marginals) is attracting attention of many researchers (see the recent survey [6] about general results and particular examples). Both the concave and the multi-marginal transportation problems have potential applications in economics (see [5] and [6]).

2 Results

Definition 2.1. We say that a couple of distinct affine functions l_1, l_2 satisfies the non-degeneracy assumption **(A)** if the set

$$\Gamma_{1,2} = \{l_1 = l_2\} \neq \emptyset$$

is not parallel to one of the axes.

Example 2.2. The assumption **(A)** is violated for $c(x, y) = \min(x, x + y)$. Since $c(x, y) = x + \min(0, y)$, the corresponding problem is degenerate and every $\pi \in \Pi(\mu, \nu)$ is optimal.

Definition 2.3. Let l_1, l_2 be a couple of affine functions satisfying **(A)** and let $M = (x_0, y_0)$ be a point that belongs to $\Gamma_{1,2} = \{l_1 = l_2\}$. Let Q_{M,l_1,l_2} be that one of the sets

$$\begin{aligned} & \left\{x \leq x_0, y \geq y_0\right\} \cup \left\{x \geq x_0, y \leq y_0\right\}, \\ & \left\{x \leq x_0, y \leq y_0\right\} \cup \left\{x \geq x_0, y \geq y_0\right\} \end{aligned}$$

which does not contain $\Gamma_{1,2}$. More precisely, Q_{M,l_1,l_2} is defined by the following condition:

$$\Gamma_{1,2} \cap Q_{M,l_1,l_2} = M.$$

Definition 2.4. Given a Borel cost function $c: \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R} \cup \{\infty\}$, we say that a subset $S \subset \mathbb{R} \times \mathbb{R}$ is c -cyclically monotone (or simply cyclically monotone) if, for every non-empty sequence of its elements $(x_1, y_1), \dots, (x_n, y_n)$, the following inequality holds:

$$c(x_1, y_1) + c(x_2, y_2) + \dots + c(x_n, y_n) \leq c(x_1, y_n) + c(x_2, y_1) + \dots + c(x_n, y_{n-1}). \quad (2)$$

It is known that for a broad class of cost functions any solution π to (1) satisfies $\pi(S) = 1$ for some cyclically monotone set S .

The following lemma is a version of the so-called “no-crossing rule” (see [5]).

Lemma 2.5. Let l_1, l_2 be affine functions satisfying **(A)** and let π be a solution to the Monge–Kantorovich problem (1) with

$$c = \min(l_1, l_2).$$

Then there exists a point $M \in \{l_1 = l_2\}$ such that the support of π is contained in Q_{M, l_1, l_2} .

Proof. It is clear that shifting the coordinates $x \rightarrow x - x_0, y \rightarrow y - y_0$ we can deal from the very beginning with linear functions l_1, l_2 . In particular, the origin belongs to $\Gamma_{1,2}$. In addition, since the marginals are fixed, the assertion is invariant with respect to subtracting a linear function l : in place of c one can deal with

$$c - l = \min(l_1 - l, l_2 - l).$$

Passing to this situation if necessary, we can deal with the case $c = \min(ax, by)$ for some $a \neq 0, b \neq 0$. Multiplying by a constant, we reduce the problem to the case $c = \min(x, by)$. Let $b > 0$ ($b < 0$ can be considered similarly).

Note that every two-points set $\{(x_1, y_1), (x_2, y_2)\}$ satisfying

$$\max(x_1, by_1) < \min(x_2, by_2)$$

is not cyclically monotone. Indeed, this can be easily verified by direct computations:

$$\min(x_1, by_1) + \min(x_2, by_2) > x_1 + \min(x_2, by_1) \geq \min(x_1, by_2) + \min(x_2, by_1).$$

Now let us find the smallest number s such that

$$\mu(-\infty, s) = \nu(s/b, +\infty). \quad (3)$$

We claim that the measure π is concentrated on the set

$$Q_{(s, bs), x, by} = \left\{ x \leq s, y \geq s/b \right\} \cup \left\{ x \geq s, y \leq s/b \right\}.$$

Assume the contrary and find a point (x_1, y_1) from the support of π such that, say,

$$x_1 < s, y_1 < s/b.$$

Then (3) implies that there exists another point (x_2, y_2) from the support of π such that

$$x_2 > s, y_2 > s/b.$$

But this means that $\max(x_1, by_2) < s < \min(x_2, by_1)$, hence c -monotonicity is violated. \square

Definition 2.6. We say that we are given a (μ, ν) -partition of order n if

1. The x -axis and y -axis are represented as unions of n non-empty disjoint connected sets

$$\begin{aligned}\mathbb{R} \times \{0\} &= I_1 \cup I_2 \dots \cup I_n, \\ \{0\} \times \mathbb{R} &= J_1 \cup J_2 \dots \cup J_n.\end{aligned}$$

2.

$$\mu(I_i) = \nu(J_i) > 0, \quad \forall i \in \{1, 2, \dots, n\}.$$

Let us proceed to our first main result.

Theorem 2.7. *Let l_1, \dots, l_n be n affine functions such that every two of them satisfy assumption **(A)**, and, moreover, every set $A_i = \{c = l_i\}$, $1 \leq i \leq n$ has a non-empty interior, where*

$$c = \min_{1 \leq i \leq n} (l_1, l_2, \dots, l_n)$$

is the corresponding cost function.

Then, for every solution π to the Monge–Kantorovich problem (1), there exists a (μ, ν) -partition of order $k \leq n$ such that

- 1) π is concentrated on $\cup_{i=1}^k I_i \times J_i$,
- 2) for every $1 \leq i \leq k$ either $\pi(I_i \times J_i) = 0$ or (after a suitable renumeration of the functions l_i)

$$c = l_i$$

on $I_i \times J_i$.

Proof. Let us fix i and consider the set

$$\Omega_i = \{c = l_i\}.$$

Assume that $\pi(\Omega_i) > 0$. Then for every $j \neq i$ set

$$h_{ij} = \min(l_i, l_j)$$

and consider the restrictions $\pi_{ij} = \pi|_{\Omega_{ij}}$ to the set

$$\Omega_{ij} = \{c = h\}.$$

We claim that π_{ij} is optimal for the cost function h_{ij} and the projections $\pi_{ij} \circ Pr_X^{-1}$, $\pi_{ij} \circ Pr_Y^{-1}$ onto the axes. Indeed, assuming the contrary consider another measure

$$\tilde{\pi}_{ij} = \pi|_{\Omega_{ij}^c} + \hat{\pi}_{ij},$$

where $\hat{\pi}_{ij}$ is optimal for h_{ij} and $\pi_{ij} \circ Pr_X^{-1}$, $\pi_{ij} \circ Pr_Y^{-1}$. Using that $c \leq h_{ij}$, $c = h_{ij}$ on Ω_{ij} and $\hat{\pi}_{ij}$ is optimal, we obtain

$$\int cd\tilde{\pi}_{ij} = \int cd\pi|_{\Omega_{ij}^c} + \int cd\hat{\pi}_{ij} \leq \int cd\pi|_{\Omega_{ij}^c} + \int h_{ij}d\hat{\pi}_{ij} < \int cd\pi|_{\Omega_{ij}^c} + \int h_{ij}d\pi_{ij} = \int cd\pi.$$

This contradicts the optimality of π .

Applying Lemma 2.5 we obtain that the supports of $\pi|_{\Omega_i} = \pi_{ij}|_{\Omega_i}$ and $\pi|_{\Omega_j} = \pi_{ij}|_{\Omega_j}$ are contained in the sets $L_{ij}^1 \times M_{ij}^1$, $L_{ij}^2 \times M_{ij}^2$, respectively, where L_{ij}^1 and $L_{ij}^2 = \mathbb{R} \setminus L_{ij}^1$ are disjoint and connected (the same is true for M_{ij}^1, M_{ij}^2). The i th intervals of the desired (μ, ν) -partition are defined as follows:

$$I_i = \cap_{j \neq i} L_{ij}^1, \quad J_i = \cap_{j \neq i} M_{ij}^1.$$

By construction $l_i = c$ on $I_i \times J_i$, hence $I_i \times J_i \subset \Omega_i$, and the support of $\pi|_{\Omega_i}$ is contained in $I_i \times J_i$. The proof is complete. \square

Remark 2.8. Let π be a solution to the Monge–Kantorovich problem with the cost function c satisfying assumptions of Theorem 2.7 and let $\{I_i, J_i\}$ be the corresponding (μ, ν) -partition. Then every measure with marginals μ, ν concentrated on $\cup_{i=1}^k I_i \times J_i$ solves the same Monge–Kantorovich problem.

Moreover, if such a measure is concentrated on the graph of a mapping T , then T is the corresponding optimal transportation.

Theorem 2.7 shows, in particular, that the transportation problem is reduced to a finite-dimensional problem of finding an optimal partition with the constraint $c = l_i$ on $I_i \times J_i$. In Theorem 2.9 below we present yet another equivalent finite-dimensional problem, where we relax the latter constraint on partitions and replace integrals over minima by minima of certain (easy computable) integrals. This viewpoint might be useful for computational purposes.

Theorem 2.9. Let l_1, \dots, l_n be affine functions of the form

$$l_j = a_j x + b_j y + c_j$$

satisfying the assumptions of Theorem 2.7 and let

$$c = \min_{1 \leq i \leq n} (l_1, l_2, \dots, l_n).$$

For every (μ, ν) -partition \mathcal{P} , we define a functional J in the following way:

$$J(\mathcal{P}) = \sum_i \min_j \left(a_j \int_{I_i} x d\mu + b_j \int_{J_i} y d\nu + c_j \mu(I_i) \right).$$

Then the minimal value of the functional J over all (μ, ν) -partitions of order not greater than n coincides with the minimum $K(\mu, \nu)$ of the Kantorovich functional for the cost function c .

Remark 2.10. Note that

$$J(\mathcal{P}) = \sum_i \min_j \int_{I_i \times J_i} l_j d\pi. \quad (4)$$

for every π with marginals μ, ν and with the support in $\cup_{i=1}^k I_i \times J_i$. In particular,

$$J(\mathcal{P}) = \sum_i \min_j \frac{1}{\mu(I_i)} \int_{I_i \times J_i} l_j d\mu|_{I_i} \times \nu|_{J_i}.$$

Proof. It follows from representation (4) that

$$J(\mathcal{P}) \geq \sum_i \int_{I_i \times J_i} \min_j l_j d\pi = \int c d\pi$$

for every partition \mathcal{P} and every measure π with marginals μ, ν . Thus,

$$J(\mathcal{P}) \geq K(\mu, \nu).$$

On the other hand, given a solution π to the Monge–Kantorovich problem, one can consider the particular partition \mathcal{P}_0 with the properties established in Theorem 2.7. Using that $l_i = c$ on $I_i \times J_i$ for every i , one obtains

$$J(\mathcal{P}_0) = \sum_i \min_j \int_{I_i \times J_i} l_j d\pi = \sum_i \int_{I_i \times J_i} l_i d\pi = \sum_i \int_{I_i \times J_i} c d\pi = \int c d\pi = K(\mu, \nu).$$

The proof is complete. □

Example 2.11. Let μ and ν be the same Lebesgue measure on $[0, 1]$. Consider the set Π of couples of partitions $(\mathcal{P}_x, \mathcal{P}_y)$ of $[0, 1]$ of the form

$$I_i = [t_{i-1}, t_i), \quad J_i = [s_{i-1}, s_i)$$

with the property

$$s_i - s_{i-1} = t_i - t_{i-1}.$$

Then according to Theorem 2.9 the value of the Kantorovich functional equals

$$\min_{(\mathcal{P}_x, \mathcal{P}_y) \in \Pi} \sum_i (t_i - t_{i-1}) \min_j \left(a_j \frac{t_i + t_{i-1}}{2} + b_j \frac{s_i + s_{i-1}}{2} + c_j \right).$$

Finally, let us make some remarks about the multi-marginal case. We give below a generalization of our main result for the case when the number of affine functions coincides with the number of marginals. This covers, in particular, the cost function

$$c(x_1, x_2, \dots, x_n) = \min\{x_1, x_2, \dots, x_n\}.$$

We omit the proofs because they are completely similar to the case of two marginals.

Remark 2.12. From the description of solutions to the Monge–Kantorovich problem for this cost function one can conclude that the straightforward generalization of Theorem 2.7 fails at least in the following respect: the projections of $\text{supp}(\pi) \cap \{c = l_i\}$ can have intersections for different i .

Definition 2.13. We say that an n -tuple of distinct affine functions of n arguments l_1, \dots, l_n satisfies the non-degeneracy assumption if the set

$$\Gamma_{1, \dots, n} := \{l_1 = l_2 = \dots = l_n\}$$

is a straight line spanned by an n -dimensional vector that has no zero components.

Definition 2.14. Let l_1, \dots, l_n be an n -tuple of distinct affine functions satisfying the non-degeneracy assumption. Suppose that $M = (x_1^0, \dots, x_n^0)$ is a point from

$$\Gamma_{l_1, \dots, l_n} := \{l_1 = l_2 = \dots = l_n\}.$$

Let \mathcal{S} be the set $\{\leq, \geq\}^n$, i.e., the set of sequences of n symbols " \leq " or " \geq ". We shall agree that " \leq " coincides with " \geq " and " \geq " coincides with " \leq ".

Finally, for any $s \in \mathcal{S}$, let us define Q_s as follows:

$$Q_s = \bigcup_{i \in \{1, \dots, n\}} \{x_1 s[1] x_1^0, x_2 s[2] x_2^0, \dots, x_i s[i] x_i^0, \dots, x_n s[n] x_n^0\}.$$

Definition 2.15. Take a directing vector v of Γ_{l_1, \dots, l_n} with $v_1 > 0$. Define $t \in \mathcal{S}$ by the following rule: $t[i] = \geq$ if $v_i > 0$ and $t[i] = \leq$ if $v_i < 0$. The set Q_t is further referred to as Q_{M, l_1, \dots, l_n} .

Lemma 2.16. Let l_1, \dots, l_n be an n -tuple of distinct affine functions satisfying the non-degeneracy assumption and let π be a solution to the Monge–Kantorovich problem with n marginals and the cost function

$$c = \min(l_1, \dots, l_n).$$

Assume, in addition, that every set $\{c = l_i\}$ has a non-empty interior. Then there exists a point $M \in \Gamma_{1, \dots, n}$ such that the support of π is contained in Q_{M, l_1, \dots, l_n} .

Applying Lemma 2.16, we obtain the following result.

Theorem 2.17. *Let μ_i , $1 \leq i \leq n$, be atomless probability measures on the real line. Define*

$$s = \sup \left\{ x : \sum_{i=1}^n \mu_i(-\infty, x] \leq 1 \right\}.$$

Then the measure $\pi \in \Pi(\mu_1, \dots, \mu_n)$ solves the Monge–Kantorovich problem with marginals μ_i and the cost function

$$c = \min(x_1, \dots, x_n)$$

if and only if every point $x \in \text{supp}(\pi) \subset \mathbb{R}^n$ satisfies the following conditions:

- 1) if $x_i \leq s$, then $x_j \geq s$ for every $j \neq i$,*
- 2) if $x_i \geq s$, then there exists $j \neq i$ such that $x_j \leq s$.*

References

- [1] Bogachev V.I., Kolesnikov A.V., The Monge–Kantorovich problem: achievements, connections, and perspectives. *Uspehi Matem. Nauk* 67 (5) (2012), 3–110 (in Russian); English transl.: *Russian Math. Surveys* 67 (5) (2012), 785–890.
- [2] Bogachev V.I., *Measure theory*. V. 1, 2, Springer, Berlin, 2007.
- [3] Delon J., Salomon J., Sobolevskii A., Local matching indicators for transport problems with concave costs, *SIAM J. Disc. Math.* 26 (2) (2012), 801–827.
- [4] Gangbo W., McCann R.J., The geometry of optimal transportation, *Acta Math.* 177 (1996), 113–161.
- [5] McCann R.J., Exact solutions to the transportation problem on the line, *R. Soc. Lond. Proc. Ser. A Math. Phys. Eng. Sci.* 455 (1999), 1341–1380.
- [6] Pass B., Multi-marginal optimal transport: theory and applications, *ESAIM: Math. Model. Numer. Anal.* 49 (2015), 1771–1790.
- [7] Pegon P., Piazzoli D., Santambrogio F., Full characterization of optimal transport plans for concave costs, *Disc. Cont. Dyn. Syst. A* 35(12) (2015), 6113–6132.
- [8] Villani C., *Topics in optimal transportation*, Amer. Math. Soc., Rhode Island, 2003.