

Generalized model of interacting integrable tops

A. Grekov ^{◇‡‡‡} I. Sechin ^{◇‡‡} A. Zotov ^{◇‡§‡‡}

[◇] – Steklov Mathematical Institute of Russian Academy of Sciences,
Gubkina str. 8, Moscow, 119991, Russia

[‡] – ITEP, B. Cheremushkinskaya str. 25, Moscow, 117218, Russia

[‡] – Center for Advanced Studies, Skolkovo Institute of Science and Technology,
Nobel str. 1, Moscow, 143026, Russia

[§] - National Research University Higher School of Economics,
Usacheva str. 6, Moscow, 119048, Russia

[‡] – Moscow Institute of Physics and Technology,
Institutskii per. 9, Dolgoprudny, Moscow region, 141700, Russia

E-mails: grekovandrew@mail.ru, shnbuz@gmail.com, zotov@mi-ras.ru

Abstract

We introduce a family of classical integrable systems describing dynamics of M interacting \mathfrak{gl}_N integrable tops. It extends the previously known model of interacting elliptic tops. Our construction is based on the GL_N R -matrix satisfying the associative Yang-Baxter equation. The obtained systems can be considered as extensions of the spin type Calogero-Moser models with (the classical analogues of) anisotropic spin exchange operators given in terms of the R -matrix data. In $N = 1$ case the spin Calogero-Moser model is reproduced. Explicit expressions for \mathfrak{gl}_{NM} -valued Lax pair with spectral parameter and its classical dynamical r -matrix are obtained. Possible applications are briefly discussed.

Contents

| | | |
|----------|--|-----------|
| 1 | Introduction | 2 |
| 2 | Lax equations | 7 |
| 2.1 | <i>R</i> -matrix properties | 7 |
| 2.2 | Lax pair and equations of motion | 9 |
| 2.3 | Hamiltonian description | 12 |
| 2.4 | Interacting tops | 14 |
| 3 | Classical <i>r</i>-matrix | 16 |
| 4 | Examples | 17 |
| 4.1 | Elliptic models | 17 |
| 4.2 | Trigonometric models | 19 |
| 4.3 | Rational models | 21 |
| 5 | Discussion | 22 |
| 6 | Appendix | 23 |
| 6.1 | A: Definitions and identities | 23 |
| 6.2 | B: Spin \mathfrak{gl}_M Calogero-Moser model | 24 |
| 6.3 | C: Integrable \mathfrak{gl}_N tops | 26 |
| | References | 28 |

1 Introduction

In this paper we describe the classical integrable \mathfrak{gl}_{NM} model given by the Hamiltonian of the following form:

$$\mathcal{H} = \sum_{i=1}^M \frac{p_i^2}{2} + \sum_{i=1}^M H^{\text{top}}(\mathcal{S}^{ii}) + \frac{1}{2} \sum_{i,j:i \neq j}^M \mathcal{U}(\mathcal{S}^{ij}, \mathcal{S}^{ji}, q_i - q_j), \quad (1.1)$$

where p_i and q_j are the canonical variables:

$$\{p_i, q_j\} = \delta_{ij}, \quad \{p_i, p_j\} = \{q_i, q_j\} = 0, \quad i, j = 1 \dots M. \quad (1.2)$$

For all $i, j = 1 \dots M$ \mathcal{S}^{ij} are $N \times N$ matrices of "classical spin" variables, i.e.

$$\mathcal{S}^{ij} = \sum_{a,b=1}^N \mathcal{S}_{ab}^{ij} e_{ab} \in \text{Mat}(N, \mathbb{C}), \quad (1.3)$$

where $\{e_{ab}, a, b = 1 \dots N\}$ is the standard basis in $\text{Mat}(N, \mathbb{C})$. They are naturally arranged into $NM \times NM$ block-matrix \mathcal{S} :

$$\mathcal{S} = \sum_{i,j=1}^M E_{ij} \otimes \mathcal{S}^{ij} = \sum_{i,j=1}^M \sum_{a,b=1}^N \mathcal{S}_{ab}^{ij} E_{ij} \otimes e_{ab} \in \text{Mat}(NM, \mathbb{C}), \quad (1.4)$$

where $\{E_{ij}, i, j = 1 \dots M\}$ is the standard basis in $\text{Mat}(M, \mathbb{C})$. The Poisson structure is given by the Poisson-Lie brackets on \mathfrak{gl}_{NM}^* Lie coalgebra:

$$\{\mathcal{S}_{ab}^{ij}, \mathcal{S}_{cd}^{kl}\} = \mathcal{S}_{cb}^{kj} \delta^{il} \delta_{ad} - \mathcal{S}_{ad}^{il} \delta^{kj} \delta_{bc}. \quad (1.5)$$

Integrable tops. In order to clarify the structure of the Hamiltonian (1.1) consider the case $M = 1$. Then the last term in (1.1) is absent, and we are left with a free particle (with momenta p_1) and the Hamiltonian $H^{\text{top}}(\mathcal{S}^{11})$ of integrable top of Euler-Arnold type [3]. Here we deal with the models admitting the Lax pairs with spectral parameter on elliptic curves [32]. The general form for equations of motion (for the top like models) is

$$\dot{S} = [S, J(S)], \quad (1.6)$$

where $S \in \text{Mat}(N, \mathbb{C})$ is the matrix of dynamical variables, while the inverse inertia tensor J is a linear map

$$J(S) = \sum_{i,j,k,l=1}^N J_{ijkl} e_{ij} S_{lk} \in \text{Mat}(N, \mathbb{C}) \quad (1.7)$$

In the general case the model (1.6) is not integrable. It is integrable for some special $J(S)$ only. More precisely, here we consider special tops, which were described in [19, 36], [17], [1, 20] for elliptic, trigonometric and rational cases respectively. All of them can be written [20, 22] in the R -matrix form based on a quantum GL_N R -matrix (in the fundamental representation) satisfying the associative Yang-Baxter equation [12, 28]:

$$R_{12}^h(q_{12}) R_{23}^\eta(q_{23}) = R_{13}^\eta(q_{13}) R_{12}^{h-\eta}(q_{12}) + R_{23}^{\eta-h}(q_{23}) R_{13}^h(q_{13}), \quad q_{ab} = q_a - q_b. \quad (1.8)$$

Having solution of (1.8) with some additional properties (see the next Section) the inverse inertia tensor comes from the term $m_{12}(z)$ in the classical limit expansion:

$$R_{12}^h(z) = \frac{1}{\hbar} 1_N \otimes 1_N + r_{12}(z) + \hbar m_{12}(z) + O(\hbar^2). \quad (1.9)$$

Namely, for

$$m_{12}(z) = \sum_{i,j,k,l=1}^N m_{ijkl}(z) e_{ij} \otimes e_{kl} \quad (1.10)$$

the components of J are

$$J_{ijkl} = m_{ijkl}(0), \quad (1.11)$$

that is

$$J(S) = \text{tr}_2(m_{12}(0) S_2), \quad S_2 = 1_N \otimes S. \quad (1.12)$$

The Hamiltonian of the model is of the form:

$$H^{\text{top}}(S) = \frac{1}{2} \text{tr}(S J(S)) = \frac{1}{2} \text{tr}_{12}(m_{12}(0) S_1 S_2), \quad S_1 = S \otimes 1_N. \quad (1.13)$$

This expression enters (1.1). The phase space of the model is a coadjoint orbit

$$\mathcal{M}^{\text{top}} = \mathcal{O}_N \quad (1.14)$$

of GL_N Lie group, i.e. the space spanned by S_{ij} with some fixed eigenvalues of matrix S (or the Casimir functions $C_k = \text{tr} S^k$). Its dimension depends on the eigenvalues. The minimal orbit $\mathcal{O}_N^{\text{min}}$ corresponds to $N - 1$ coincident eigenvalues, i.e the matrix S (up to a matrix proportional to identity matrix) is of rank one:

$$\dim \mathcal{O}_N^{\text{min}} = 2(N - 1). \quad (1.15)$$

The Lax pair is given in the Appendix C.

Spin Calogero-Moser model. In the case $N = 1$ the second term in (1.1) is trivial, and the last one boils down to the spin Calogero-Moser model [13, 4]:

$$H^{\text{spin}} = \sum_{i=1}^M \frac{p_i^2}{2} - \sum_{i>j}^M S_{ij} S_{ji} E_2(q_i - q_j), \quad (1.16)$$

where $E_2(q)$ is the second Eisenstein function (A.4). Some details of the spin Calogero-Moser model are given in the Appendix B. Let us only remark here that the model (1.16) is integrable through the Lax representation and the classical r -matrix structure on the constraints

$$S_{ii} = \nu \text{ for all } i = 1 \dots M \quad (1.17)$$

supplemented by some gauge fixation conditions generated by the coadjoint action of the Cartan subgroup $\mathfrak{H}_M \subset \text{GL}_M$. That is the phase space of the model is given by

$$\mathcal{M}^{\text{spin}} = T^* \mathfrak{h}_M \times \mathcal{O}_M // \mathfrak{H}_M, \quad (1.18)$$

where $\mathfrak{h}_M = \text{Lie}(\mathfrak{H}_M)$ is the Lie algebra of \mathfrak{H}_M , and \mathcal{O}_M is an orbit of the coadjoint action of GL_M . The first factor in (1.18) describes the many-body degrees of freedom (1.2), and the second factor describes the "classical spin" variables. In the general case the spin variables can be parameterized by the set of canonically conjugated variables:

$$S_{ij} = \sum_{a=1}^N \xi_a^i \eta_a^j, \quad (1.19)$$

$$\{\xi_a^i, \eta_b^j\} = \delta_{ab} \delta_{ij}, \quad i, j = 1 \dots M, \quad a, b = 1 \dots N.$$

The Poisson structure (B.7) is reproduced in this way. Using these notations it is easy to see that

$$S_{ij} S_{ji} = \sum_{a,b=1}^N \xi_a^i \eta_a^j \xi_b^j \eta_b^i = \text{tr}(\mathcal{S}^{ii} \mathcal{S}^{jj}), \quad (1.20)$$

and the potential in the Hamiltonian (1.16) takes the form

$$\mathcal{V}^{\text{spin}}(\mathcal{S}^{ii}, \mathcal{S}^{jj}, q_{ij}) = -\text{tr}(\mathcal{S}^{ii} \mathcal{S}^{jj}) E_2(q_i - q_j). \quad (1.21)$$

Below we construct anisotropic (in $\text{Mat}(N, \mathbb{C})$ space) generalizations of (1.21).

In the special case, when the matrix of spin variables S is of rank 1 (it is the minimal \mathcal{O}_M^{\min} orbit (1.15))

$$S_{ij} = \xi_i \eta_j \quad (1.22)$$

the reduction with respect to the action of \mathfrak{H}_M leads to the spinless Calogero-Moser (CM) model [7, 18] since the second factor in (1.18) become trivial. Indeed, plugging (1.22) into (1.16) and using (1.17) we get

$$H^{\text{spin}} = \sum_{i=1}^M \frac{p_i^2}{2} - \nu^2 \sum_{i>j}^M E_2(q_i - q_j). \quad (1.23)$$

The spinless Calogero-Moser models are gauge equivalent to the special top with the minimal orbit (1.15). See [19, 17, 1] for details.

Interacting tops. Turning back to the \mathfrak{gl}_{NM} model (1.1) consider the special case when the matrix \mathcal{S} is of rank 1:

$$\mathcal{S}_{ab}^{ij} = \xi_a^i \eta_b^j. \quad (1.24)$$

We will see that in this case the last term in (1.1) is rewritten in the form

$$\mathcal{U}(\mathcal{S}^{ij}, \mathcal{S}^{ji}, q_i - q_j) = \mathcal{V}(\mathcal{S}^{ii}, \mathcal{S}^{jj}, q_i - q_j), \quad (1.25)$$

and the Hamiltonian (1.1) acquires the form

$$\mathcal{H}^{\text{tops}} = \sum_{i=1}^M \frac{p_i^2}{2} + \sum_{i=1}^M H^{\text{top}}(\mathcal{S}^{ii}) + \frac{1}{2} \sum_{i,j:i \neq j}^M \mathcal{V}(\mathcal{S}^{ii}, \mathcal{S}^{jj}, q_i - q_j). \quad (1.26)$$

It describes mechanics of M interacting integrable \mathfrak{gl}_N tops. The Hamiltonian of (1.26) type was introduced by A.P. Polychronakos [29] from his study of matrix models. Then the elliptic version of model (1.1) and (1.26) was described as \mathfrak{gl}_{NM} Hitchin system [35, 25, 36] (see some details in Section 4), and (1.1) was also generalized for arbitrary complex Lie group [23].

Similarly to the spin Calogero-Moser model the general model (1.1) requires additional constraints (cf. (1.17))

$$\text{tr}(\mathcal{S}^{ii}) = \nu \text{ for all } i = 1 \dots M. \quad (1.27)$$

They should be supplied with some gauge fixation conditions generated by the coadjoint action of $\mathfrak{H}'_{NM} \subset \mathfrak{H}_{NM}$ – subgroup of the Cartan subgroup $\mathfrak{H}_{NM} \subset \text{GL}_{NM}$ with elements of the form $\sum_{i=1}^M h_i E_{ii} \otimes 1_N$. Together with (1.27) the gauge fixation conditions are the second class constraints, and one can perform the Dirac reduction procedure to compute the final Poisson structure starting from the linear one (1.5). The phase space of the general model (1.1) is of the form:

$$\mathcal{M}^{\text{gen}} = T^* \mathfrak{h}'_{NM} \times \mathcal{O}_{NM} // \mathfrak{H}'_{NM}, \quad \mathfrak{h}'_{NM} = \text{Lie}(\mathfrak{H}'_{NM}). \quad (1.28)$$


For the interacting tops case (1.24)-(1.26) the orbit \mathcal{O}_{NM} becomes \mathcal{O}_{NM}^{\min} . Then the phase space

$$\mathcal{M}^{\text{tops}} = T^* \mathfrak{h}'_{NM} \times \mathcal{O}_{NM}^{\min} // \mathfrak{H}'_{NM} \quad (1.29)$$

has dimension $2NM$, while its "spin part" is of dimension

$$\dim \left(\mathcal{O}_{NM}^{\min} // \mathfrak{H}'_{NM} \right) = 2NM - 2M. \quad (1.30)$$

A brief summary of the described models is given in the following scheme:



Purpose of the paper is to describe a family of the models (1.1) and (1.26) in terms of R -matrices satisfying the associative Yang-Baxter equation (1.8). We give explicit formulae for $NM \times NM$ Lax pair with spectral parameter (see the next Section) and compute the Hamiltonians (1.1) and (1.26). As a result we obtain the potentials

$$\mathcal{U}(\mathcal{S}^{ij}, \mathcal{S}^{ji}, q_i - q_j) = \text{tr}_{12} \left(\partial_{q_i} r_{21}(q_{ij}) P_{12} \mathcal{S}_1^{ij} \mathcal{S}_2^{ji} \right) \quad (1.32)$$

for the general model (1.1) and

$$\mathcal{V}(\mathcal{S}^{ii}, \mathcal{S}^{jj}, q_i - q_j) = \text{tr}_{12} \left(\partial_{q_i} r_{12}(q_{ij}) \mathcal{S}_1^{ii} \mathcal{S}_2^{jj} \right) \quad (1.33)$$

for the model of interacting tops (1.26). Notice that in the simplest case related to the rational Yang's XXX R -matrix

$$R_{12}^z(q_{ij}) = \frac{1_N \otimes 1_N}{z} + \frac{P_{12}}{q_i - q_j} \quad (1.34)$$

we get just the spin Calogero-Moser model written in terms of matrix variables:

$$\mathcal{V} = - \frac{\text{tr}(\mathcal{S}^{ii} \mathcal{S}^{jj})}{(q_i - q_j)^2}. \quad (1.35)$$

Next, we proceed to the classical (dynamical) r -matrix. It is similar to the one for the spin Calogero-Moser case [4] but this time its matrix elements are R -matrices themselves. The classical exchange relations are verified directly. This guarantees the Poisson commutativity of the Hamiltonians generated by the Lax matrix.

The answers (1.32) and (1.33) depend on the classical r -matrix, which appears from the quantum one in the limit (1.9). The quantum R -matrix enters the higher Hamiltonians. It should satisfy a set of properties which we discuss in the next Section. The most general R -matrix satisfying all the required properties is the elliptic Baxter-Balavin's one. In this case

the the integrable models are known. They were first described by Polychronakos in [29] and later reproduced as Hitchin type systems on the bundles with nontrivial characteristic classes in [35, 25].

The family of the obtained models includes new integrable systems in the trigonometric and rational cases. While the quantization of the potential \mathcal{V} from (1.35) is given by isotropic spin exchange operator $\hat{\mathcal{V}} = -P_{ij}/(q_i - q_j)^2$, the obtained general answer (1.32)-(1.33) leads to the anisotropic potentials. An example of such anisotropic extension to the spin (trigonometric) Calogero-Moser-Sutherland model was first suggested by Hikami and Wadati [16] at quantum level. From the point of view of (1.33) their answer corresponds to the \mathfrak{gl}_2 XXZ r -matrix. At the same time the set of trigonometric R -matrices satisfying the required properties is much larger [2, 30], and all these R -matrices can be used for construction of the integrable tops [17]. The results of the present paper are also valid for all these cases. An example based on the \mathfrak{gl}_2 7-th vertex deformation of the XXZ R -matrix is given Section 4. Similarly, in the rational case the admissible R -matrices includes not only the Yang's R -matrix (1.34) but also its deformations such as 11-vertex R -matrix [8] and its higher rank versions [20]. An example related to 11-vertex R -matrix is given in Section 4.

Possible applications of the described models are discussed in the end. Namely, we argue that the obtained models can be used for construction of higher Hamiltonians for the anisotropic generalizations of the Haldane-Shastry-Inozemtsev long-range spin chains. The latter is important for the proof of integrability of these chains, which still remains an open problem.

2 Lax equations

In this Section we construct the $NM \times NM$ Lax pair $\mathcal{L}(z), \mathcal{M}(z)$ satisfying the Lax equations

$$\dot{\mathcal{L}}(z) = [\mathcal{L}(z), \mathcal{M}(z)] \quad (2.1)$$

for the model (1.1). Our construction is based on GL_N R -matrix – solution of the associative Yang-Baxter equation (1.8). Besides (1.8) the R -matrix should also satisfy a set of properties.

2.1 R -matrix properties.

We consider R -matrices satisfying (1.8) and (1.9). Let us also impose the following set of conditions for GL_N R -matrices under consideration:

Expansion near $z = 0$:

$$R_{12}^h(z) = \frac{1}{z} P_{12} + R_{12}^{h,(0)} + z R_{12}^{h,(1)} + O(z^2), \quad (2.2)$$

Also,

$$R_{12}^{z,(0)} = \frac{1}{z} 1_N \otimes 1_N + r_{12}^{(0)} + O(z), \quad r_{12}(z) = \frac{1}{z} P_{12} + r_{12}^{(0)} + z r_{12}^{(1)} + O(z^2). \quad (2.3)$$

Skew-symmetry:¹

$$R_{12}^h(z) = -R_{21}^{-h}(-z) = -P_{12}R_{12}^{-h}(-z)P_{12}, \quad P_{12} = \sum_{i,j=1}^N E_{ij} \otimes E_{ji}. \quad (2.4)$$

Unitarity:

$$R_{12}^h(z)R_{21}^h(-z) = f^h(z) 1_N \otimes 1_N, \quad f^h(z) = \wp(\hbar) - \wp(z). \quad (2.5)$$

We are also going to use the Fourier symmetry:

$$R_{12}^h(z)P_{12} = R_{12}^z(\hbar) \quad (2.6)$$

It is not necessary but convenient property. The following relations on the coefficients of expansions (1.9) and (2.2) follow from the skew-symmetry:

$$\begin{aligned} r_{12}(z) &= -r_{21}(-z), & m_{12}(z) &= m_{21}(-z), \\ R_{12}^{h,(0)} &= -R_{21}^{-h,(0)}, & r_{12}^{(0)} &= -r_{21}^{(0)}. \end{aligned} \quad (2.7)$$

Similarly, from the Fourier symmetry we have (see details in [37]):

$$\begin{aligned} R_{12}^{z,(0)} &= r_{12}(z)P_{12}, \\ R_{12}^{z,(1)} &= m_{12}(z)P_{12}, \\ r_{12}^{(0)} &= r_{12}^{(0)}P_{12}. \end{aligned} \quad (2.8)$$

In what follows we use special notation for the R -matrix derivative:

$$F_{12}^z(q) = \partial_q R_{12}^z(q) \quad (2.9)$$

It is the R -matrix analogue of the function (A.5) entering the M -matrix of the spin Calogero-Moser model (B.3) likewise R -matrix itself is a matrix analogue of the Kronecker function (A.1) due to similarity of (A.6) and (1.8). See [21]. Then from the classical limit (1.9) we have

$$F_{12}^0(q) = \partial_q R_{12}^z(q) |_{z=0} = \partial_q r_{12}(q). \quad (2.10)$$

The latter is the R -matrix analogue of the function $-E_2(q)$ (A.12) entering the Calogero-Moser potential. Notice also that $F_{12}^0(q) = F_{21}^0(-q)$ due to (2.7). From (2.9) and (2.2) the local expansion near $q = 0$ is as follows

$$F_{12}^z(q) = -\frac{1}{q^2} P_{12} + R_{12}^{z,(1)} + O(q) \quad (2.11)$$

and, therefore,

$$F_{12}^0(q) = -\frac{1}{q^2} P_{12} + R_{12}^{z,(1)} |_{z=0} + O(q) \stackrel{(2.8)}{=} -\frac{1}{q^2} P_{12} + m_{12}(0)P_{12} + O(q). \quad (2.12)$$

¹ P_{12} entering (2.4) is the permutation operator, $(P_{12})^2 = 1_N \otimes 1_N$.

On the other hand

$$F_{12}^0(q) \stackrel{(2.10)}{=} \partial_q r_{12}(q) \stackrel{(2.3)}{=} -\frac{1}{q^2} P_{12} + r_{12}^{(1)} + O(q). \quad (2.13)$$

From (2.12) and (2.13) we conclude that

$$r_{12}^{(1)} = m_{12}(0)P_{12}. \quad (2.14)$$

In the elliptic case the set of properties is fulfilled by the Baxter-Belavin [5] R -matrix (C.14). A family of trigonometric R -matrices include the XXZ 6-vertex one, its 7-vertex deformation [8] and GL_N generalizations [2, 30]. See a brief review and applications to integrable tops in [17]. The rational R -matrices possessing the properties are the XXX Yang's R -matrix, its 11-vertex deformation [8] and higher rank analogues obtained from the elliptic case by special limiting procedure [34]. The final answer for such R -matrix was obtained in [20] through the gauge equivalence between the relativistic top with minimal orbit and the rational Ruijsenaars-Schneider model.

2.2 Lax pair and equations of motion

Using coefficients of the expansion of the GL_N R -matrix near $z = 0$ we define $NM \times NM$ Lax pair

$$L(z) = \sum_{i,j=1}^M E_{ij} \otimes L^{ij}(z), \quad L^{ij}(z) \in \text{Mat}_N \quad L(z) \in \text{Mat}_{NM}, \quad (2.15)$$

$$L^{ij}(z) = \delta_{ij} \left(p_i 1_N + \text{tr}_2(\mathcal{S}_2^{ii} R_{12}^{z,(0)} P_{12}) \right) + (1 - \delta_{ij}) \text{tr}_2(\mathcal{S}_2^{ij} R_{12}^z(q_{ij}) P_{12}). \quad (2.16)$$

and similarly for $M^{ij}(z) \in \text{Mat}_N$

$$M^{ij}(z) = \delta_{ij} \text{tr}_2(\mathcal{S}_2^{ii} R_{12}^{z,(1)} P_{12}) + (1 - \delta_{ij}) \text{tr}_2(\mathcal{S}_2^{ij} F_{12}^z(q_{ij}) P_{12}). \quad (2.17)$$

where the entries are defined from (2.2) and (2.9). The tensor notations are similar to those used in (C.3)-(C.6).

Proposition 2.1 *Consider an R -matrix satisfying the associative Yang-Baxter equation (1.8), the classical limit (1.9) and the set of properties from the previous paragraph. Then the Lax equation (2.1) holds true for the Lax pair (2.15)-(2.17) on the constraints*

$$\text{tr}(\mathcal{S}^{ii}) = \text{const}, \quad \forall i \quad (2.18)$$

(cf. (4.4)) and provides the following equations of motion for off-diagonal $N \times N$ blocks of \mathcal{S} :

$$\begin{aligned} \dot{\mathcal{S}}^{ij} = & \sum_{k:k \neq i,j}^M \left(\mathcal{S}^{ik} \text{tr}_2(\mathcal{S}_2^{kj} F_{12}^0(q_{kj}) P_{12}) - \text{tr}_2(\mathcal{S}_2^{ik} F_{12}^0(q_{ik}) P_{12}) \mathcal{S}^{kj} \right) + \\ & + \mathcal{S}^{ii} \text{tr}_2(\mathcal{S}_2^{ij} F_{12}^0(q_{ij}) P_{12}) - \text{tr}_2(\mathcal{S}_2^{ii} m_{12}(0)) \mathcal{S}^{ij} - \\ & - \text{tr}_2(\mathcal{S}_2^{ij} F_{12}^0(q_{ij}) P_{12}) \mathcal{S}^{jj} + \mathcal{S}^{ij} \text{tr}_2(\mathcal{S}_2^{jj} m_{12}(0)), \end{aligned} \quad (2.19)$$

for diagonal $N \times N$ blocks of \mathcal{S} :

$$\dot{\mathcal{S}}^{ii} = [\mathcal{S}^{ii}, \text{tr}_2(m_{12}(0)\mathcal{S}_2^{ii})] + \sum_{k:k \neq i}^M \left(\mathcal{S}^{ik} \text{tr}_2(\mathcal{S}_2^{ki} F_{21}^0(q_{ik})P_{12}) - \text{tr}_2(\mathcal{S}_2^{ik} F_{12}^0(q_{ik})P_{12})\mathcal{S}^{ki} \right), \quad (2.20)$$

and for momenta:

$$\dot{p}_i = - \sum_{k:k \neq i}^M \text{tr}_{23} \left(\partial_{q_i} F_{32}^0(q_{ik})P_{23} \mathcal{S}_2^{ik} \mathcal{S}_3^{ki} \right). \quad (2.21)$$

Proof: We imply $p_i = \dot{q}_i$ in the formulae above. This follows from the Hamiltonian description, which is given in the next paragraph.

1. Let us begin with the non-diagonal blocks. Consider the one numbered ij ($i \neq j$). The l.h.s. of the Lax equations reads

$$\text{l.h.s.} = \dot{\mathcal{L}}^{ij}(z) = \text{tr}_2(\dot{\mathcal{S}}_2^{ij} R_{12}^z(q_{ij})P_{12}) + \text{tr}_2(\mathcal{S}_2^{ij} F_{12}^z(q_{ij})P_{12})(\dot{q}_i - \dot{q}_j). \quad (2.22)$$

The r.h.s. of the Lax equation is as follows:

$$\text{r.h.s.} = \mathcal{L}^{ij} \mathcal{M}^{jj} - \mathcal{M}^{ii} \mathcal{L}^{ij} + \mathcal{L}^{ii} \mathcal{M}^{ij} - \mathcal{M}^{ij} \mathcal{L}^{jj} + \sum_{k:k \neq i,j}^M \left(\mathcal{L}^{ik} \mathcal{M}^{kj} - \mathcal{M}^{ik} \mathcal{L}^{kj} \right). \quad (2.23)$$

The last sum is computed using identity

$$R_{12}^z(x)F_{23}^z(y) - F_{12}^z(x)R_{23}^z(y) = F_{23}^0(y)R_{13}^z(x+y) - R_{13}^z(x+y)F_{12}^0(x), \quad (2.24)$$

which follows from (1.8). It is the R -matrix analogue of (A.7). In its turn (A.7) is the key tool underlying ansatz for the Lax pairs with spectral parameter [18]. For $k \neq i, j$ we have

$$\begin{aligned} & \mathcal{L}^{ik} \mathcal{M}^{kj} - \mathcal{M}^{ik} \mathcal{L}^{kj} = \\ & = \text{tr}_{23}(R_{12}^z(q_{ik})P_{12}\mathcal{S}_2^{ik}F_{13}^z(q_{kj})P_{13}\mathcal{S}_3^{kj}) - \text{tr}_{23}(F_{12}^z(q_{ik})P_{12}\mathcal{S}_2^{ik}R_{13}^z(q_{kj})P_{13}\mathcal{S}_3^{kj}) = \\ & = \text{tr}_{23} \left(\left(R_{12}^z(q_{ik})F_{23}^z(q_{kj}) - F_{12}^z(q_{ik})R_{23}^z(q_{kj}) \right) P_{12}P_{13} \mathcal{S}_2^{ik} \mathcal{S}_3^{kj} \right) \stackrel{(2.24)}{=} \\ & = \text{tr}_{23} \left(\left(F_{23}^0(q_{kj})R_{13}^z(q_{ij}) - R_{13}^z(q_{ij})F_{12}^0(q_{ik}) \right) P_{12}P_{13} \mathcal{S}_2^{ik} \mathcal{S}_3^{kj} \right) = \\ & = \text{tr}_{23} \left(R_{12}^z(q_{ij})P_{12} \left(\mathcal{S}_2^{ik} \mathcal{S}_3^{kj} F_{23}^0(q_{kj})P_{23} - F_{23}^0(q_{ik})P_{23} \mathcal{S}_3^{ik} \mathcal{S}_2^{kj} \right) \right). \end{aligned} \quad (2.25)$$

This expression provides the upper line in the equations of motion (2.19). To proceed we need degenerations of the identity (2.24) when $y \rightarrow 0$. It comes from the expansions (2.2), (2.11) and (2.13):

$$R_{12}^z(x)R_{23}^{z,(1)} - F_{12}^z(x)R_{23}^{z,(0)} = r_{23}^{(1)}R_{13}^z(x) - R_{13}^z(x)F_{12}^0(x) - \frac{1}{2}P_{23}\partial_x^2 R_{13}^z(x). \quad (2.26)$$

In the same way in the limit $x \rightarrow 0$ (2.24) takes the form

$$R_{12}^{z,(0)}F_{23}^z(y) - R_{12}^{z,(1)}R_{23}^z(y) = F_{23}^0(y)R_{13}^z(y) - R_{13}^z(y)r_{12}^{(1)} + \frac{1}{2}\partial_y^2 R_{13}^z(y)P_{12}. \quad (2.27)$$

Similarly to the ordinary (spin) Calogero-Moser case the terms linear in momenta in the r.h.s. (2.23) $(p_i - p_j)\mathcal{M}^{ij}$ are cancelled out by the last term in the l.h.s. of (2.22). Consider the first and the fourth terms from (2.23) without momenta. Using evaluations similar to (2.25) we get

$$\begin{aligned}
& \mathcal{L}^{ij}\mathcal{M}^{jj} - \mathcal{M}^{ij}(\mathcal{L}^{jj} - p_j 1_N) = \\
& = \text{tr}_{23} \left(\left(R_{12}^z(q_{ij})R_{23}^{z,(1)} - F_{12}^z(q_{ij})R_{23}^{z,(0)} \right) P_{12}P_{13} \mathcal{S}_2^{ij} \mathcal{S}_3^{jj} \right) \stackrel{(2.26)}{=} \\
& = \text{tr}_{23} \left(\left(r_{23}^{(1)} R_{13}^z(q_{ij}) - R_{13}^z(q_{ij})F_{12}^0(q_{ij}) - \frac{1}{2} P_{23} \partial_{q_i}^2 R_{13}^z(q_{ij}) \right) P_{12}P_{13} \mathcal{S}_2^{ij} \mathcal{S}_3^{jj} \right) = \quad (2.28) \\
& = \text{tr}_{23}(R_{12}^z(q_{ij})P_{12} \mathcal{S}_2^{ij} \mathcal{S}_3^{jj} m_{23}(0)) - \text{tr}_{23}(R_{12}^z(q_{ij})P_{12}F_{23}^0(q_{ij})P_{23} \mathcal{S}_3^{ij} \mathcal{S}_2^{jj}) - \\
& \quad - \frac{1}{2} \text{tr}_{23}(\partial_{q_i}^2 R_{12}^z(q_{ij})P_{12} \mathcal{S}_2^{ij} \mathcal{S}_3^{jj}),
\end{aligned}$$

where the relation (2.14) was also used (for the first term in the answer). The first and the second terms in the obtained answer provide the last line in the equations of motion (2.19), while the last term in (2.19) is the "unwanted term".

In the same way, using (2.27) one gets

$$\begin{aligned}
& (\mathcal{L}^{ii} - p_i 1_N)\mathcal{M}^{ij} - \mathcal{M}^{ii}\mathcal{L}^{ij} = \\
& = \text{tr}_{23}(R_{12}^z(q_{ij})P_{12} \mathcal{S}_2^{ii} \mathcal{S}_3^{ij} F_{23}^0(q_{ij})P_{23}) - \text{tr}_{23}(R_{12}^z(q_{ij})P_{12}m_{23}(0)\mathcal{S}_3^{ii} \mathcal{S}_2^{ij}) + \quad (2.29) \\
& \quad + \frac{1}{2} \text{tr}_{23}(\partial_{q_i}^2 R_{12}^z(q_{ij})P_{12} \mathcal{S}_3^{ii} \mathcal{S}_2^{ij}).
\end{aligned}$$

Again, the first two terms provide an input to equations of motion – the second line in (2.19). The last term is the "unwanted term". It is cancelled by the one from (2.28) after taking the trace over the third component and imposing the constraints (2.18).

2. Consider a diagonal $N \times N$ block (numbered ii) of the Lax equation. The l.h.s. of the Lax equations is

$$\text{l.h.s.} = \dot{\mathcal{L}}^{ii}(z) = \dot{p}_i 1_N + \text{tr}_2(\dot{\mathcal{S}}_2^{ii} R_{12}^{z,(0)} P_{12}) \stackrel{(2.8)}{=} \dot{p}_i 1_N + \text{tr}_2(\dot{\mathcal{S}}_2^{ii} r_{12}(z)). \quad (2.30)$$

The r.h.s. of the Lax equation is as follows:

$$\text{r.h.s.} = [\mathcal{L}^{ii}, \mathcal{M}^{ii}] + \sum_{k:k \neq i}^M \left(\mathcal{L}^{ik} \mathcal{M}^{ki} - \mathcal{M}^{ik} \mathcal{L}^{ki} \right). \quad (2.31)$$

The commutator term in (2.31) provides the commutator term in the equations of motion (2.20) since it is the input from the internal ii -th top's dynamics, and this was derived in [22]. See (C.2)-(C.4). In order to simplify expression in the sum we need the following degeneration of (1.8):

$$R_{12}^z(x)R_{23}^z(y) = R_{13}^z(x+y)r_{12}(x) + r_{23}(y)R_{13}^z(x+y) - \partial_z R_{13}^z(x+y), \quad (2.32)$$

It corresponds to $\hbar = \eta = z$. In the scalar case it is the identity (A.9). In the limit $x = q = -y$ from (2.32) we get

$$R_{12}^z(q)R_{23}^z(-q) = R_{13}^{z,(0)}r_{12}(q) - r_{32}(q)R_{13}^{z,(0)} - \partial_z R_{13}^{z,(0)} + F_{32}^0(q)P_{13}, \quad (2.33)$$

or, using (2.8)

$$R_{12}^z(q)R_{23}^z(-q) = (r_{13}(z)r_{32}(q) - r_{32}(q)r_{13}(z))P_{13} - F_{13}^0(z)P_{13} + F_{32}^0(q)P_{13}. \quad (2.34)$$

By differentiating (2.34) with respect to q we obtain

$$R_{12}^z(q)F_{23}^z(-q) - F_{12}^z(q)R_{23}^z(-q) = [F_{32}^0(q), r_{13}(z)]P_{13} - \partial_q F_{32}^0(q)P_{13}. \quad (2.35)$$

For $k \neq i$ consider

$$\begin{aligned} & \mathcal{L}^{ik} \mathcal{M}^{ki} - \mathcal{M}^{ik} \mathcal{L}^{ki} = \\ & = \text{tr}_{23} \left(\left(R_{12}^z(q_{ik})F_{23}^z(q_{ki}) - F_{12}^z(q_{ik})R_{23}^z(q_{ki}) \right) P_{12}P_{13} \mathcal{S}_2^{ik} \mathcal{S}_3^{ki} \right) \stackrel{(2.35)}{=} \\ & = \text{tr}_{23} \left(\left([F_{32}^0(q_{ik}), r_{13}(z)]P_{13} - \partial_{q_i} F_{32}^0(q_{ik})P_{13} \right) P_{12}P_{13} \mathcal{S}_2^{ik} \mathcal{S}_3^{ki} \right). \end{aligned} \quad (2.36)$$

The commutator term in the obtained expression yields the sum term in the equations of motion (2.20), while the last term in (2.36) provides equations of motion (2.21). Indeed,

$$\text{tr}_{23} \left(\left(\partial_{q_i} F_{32}^0(q_{ik})P_{13} \right) P_{12}P_{13} \mathcal{S}_2^{ik} \mathcal{S}_3^{ki} \right) = 1_N \text{tr}_{23} \left(\partial_{q_i} F_{32}^0(q_{ik})P_{23} \mathcal{S}_2^{ik} \mathcal{S}_3^{ki} \right), \quad (2.37)$$

and the momenta is the scalar component in the l.h.s. (2.30). ■

2.3 Hamiltonian description

The Hamiltonian function. Let us compute the Hamiltonian for the model (2.15)-(2.21). It comes from the generating function

$$\frac{1}{2N} \text{tr}(\mathcal{L}^2(z)) = \frac{1}{2N} \sum_{i=1}^M \text{tr} \left(\mathcal{L}^{ii}(z) \right)^2 + \frac{1}{2N} \sum_{i \neq j}^M \text{tr} \left(\mathcal{L}^{ij}(z) \mathcal{L}^{ji}(z) \right). \quad (2.38)$$

Consider

$$\text{tr} \left(\mathcal{L}^{ii}(z) \right)^2 = Np_i^2 + 2p_i \text{tr}_{12} \left(r_{12}(z) \mathcal{S}_2^{ii} \right) + \text{tr}_{123} \left(r_{12}(z) r_{13}(z) \mathcal{S}_2^{ii} \mathcal{S}_3^{ii} \right). \quad (2.39)$$

As before, the numbered tensor components are $\text{Mat}(N, \mathbb{C})$ -valued. In order to simplify (2.39) we use the identity (see [21])

$$r_{12}(z)r_{13}(z+w) - r_{23}(w)r_{12}(z) + r_{13}(z+w)r_{23}(w) = m_{12}(z) + m_{23}(w) + m_{13}(z+w), \quad (2.40)$$

which can be treated as a half of the classical Yang-Baxter equation². In the limit $w \rightarrow 0$ (2.40) yields

$$r_{12}(z)r_{13}(z) = r_{23}^{(0)}r_{12}(z) - r_{13}(z)r_{23}^{(0)} - F_{13}^0(z)P_{23} + m_{12}(z) + m_{23}(0) + m_{13}(z). \quad (2.41)$$

Also, we are going to use the following R -matrix property:

$$\text{tr}_1 R_{12}^q(z) = \text{tr}_2 R_{12}^q(z) = \tilde{\phi}(z, q) 1_N, \quad (2.42)$$

²The difference of two such equations gives the classical Yang-Baxter equation for the classical r -matrix.

where $\tilde{\phi}(z, q)$ is the Kronecker function (A.1) but with possibly different normalization factor and normalization of arguments. The property (2.42) holds true in the elliptic case (C.15) as well as for its trigonometric and rational degenerations. From (2.42), expansion (1.9) and (A.10) we also have similar properties for $\text{tr}_1 r_{12}(z) = \tilde{E}_1(z)$ and $\text{tr}_1 m_{12}(z)$ – they are scalar operators:

$$\text{tr}_1 R_{12}^q(z) = q^{-1} 1_N + \text{tr}_1 r_{12}(z) + q \text{tr}_1 m_{12}(z) + O(q^2). \quad (2.43)$$

Return now to (2.39). On the constraints (2.18) the second term is equal to $2p_i \tilde{E}_1(z) \text{const.}$ After summation over i it provides the Hamiltonian proportional to $\sum_{i=1}^M p_i$. Plugging (2.41) into the last term of (2.39) we get

$$\begin{aligned} & \text{tr}_{123} \left(r_{12}(z) r_{13}(z) \mathcal{S}_2^{ii} \mathcal{S}_3^{ii} \right) = \\ & = \text{tr}_{123} \left(\left(r_{23}^{(0)} r_{12}(z) - r_{13}(z) r_{23}^{(0)} - F_{13}^0(z) P_{23} + m_{23}(0) + m_{12}(z) + m_{13}(z) \right) \mathcal{S}_2^{ii} \mathcal{S}_3^{ii} \right). \end{aligned} \quad (2.44)$$

Due to (2.42) the first two terms are cancelled out after taking the trace over the component 1. By the same reason the last two terms in (2.44) provide $2\text{tr}_1(m_{12}(z))\text{tr}_{23}(\mathcal{S}_2^{ii}\mathcal{S}_3^{ii})$. These are constants on the constraints (2.18). The rest of the terms are

$$\text{tr}_{123} \left(\left(-F_{13}(z) P_{23} + m_{23}(0) \right) \mathcal{S}_2^{ii} \mathcal{S}_3^{ii} \right) \stackrel{(2.42)}{=} \tilde{E}_2(z) \text{tr} \left(\mathcal{S}^{ii} \right)^2 + N \text{tr}_{23} \left(m_{23}(0) \mathcal{S}_2^{ii} \mathcal{S}_3^{ii} \right), \quad (2.45)$$

where $\tilde{E}_2(z) 1_N = -\text{tr}_1(F_{13}^0(z)) = -\partial_z \text{tr}_1(r_{13}(z)) = -\partial_z \tilde{E}_1(z) 1_N$. It is a scalar function coming from (2.43) and similar to $E_2(z)$ (A.4). The factor N in the last term comes from tr_1 . The first term in (2.45) is a part of the Casimir function $\text{tr} \mathcal{S}^2$, and the second one is $H^{\text{top}}(\mathcal{S}^{ii})$ from (1.1):

$$H^{\text{top}}(\mathcal{S}^{ii}) = \frac{1}{2} \text{tr}_{12} \left(m_{12}(0) \mathcal{S}_1^{ii} \mathcal{S}_2^{ii} \right). \quad (2.46)$$

Next, consider

$$\begin{aligned} & \text{tr} \left(\mathcal{L}^{ij}(z) \mathcal{L}^{ji}(z) \right) = \text{tr}_{123} \left(R_{12}^z(q_{ij}) P_{12} R_{13}^z(q_{ji}) P_{13} \mathcal{S}_2^{ij} \mathcal{S}_3^{ji} \right) = \\ & = \text{tr}_{123} \left(R_{12}^z(q_{ij}) R_{23}^z(q_{ji}) P_{12} P_{13} \mathcal{S}_2^{ij} \mathcal{S}_3^{ji} \right) \stackrel{(2.34)}{=} \\ & = \text{tr}_{123} \left(\left([r_{13}(z), r_{32}(q_{ij})] - F_{13}^0(z) + F_{32}^0(q_{ij}) \right) P_{23} \mathcal{S}_2^{ij} \mathcal{S}_3^{ji} \right). \end{aligned} \quad (2.47)$$

Again, the commutator term vanishes after taking the trace over the first tensor component. Therefore,

$$\begin{aligned} & \text{tr} \left(\mathcal{L}^{ij}(z) \mathcal{L}^{ji}(z) \right) = \text{tr}_{123} \left(\left(-F_{13}^0(z) + F_{32}^0(q_{ij}) \right) P_{23} \mathcal{S}_2^{ij} \mathcal{S}_3^{ji} \right) = \\ & = \tilde{E}_2(z) \text{tr} \left(\mathcal{S}^{ij} \mathcal{S}^{ji} \right) + N \text{tr}_{12} \left(F_{21}^0(q_{ij}) P_{12} \mathcal{S}_1^{ij} \mathcal{S}_2^{ji} \right). \end{aligned} \quad (2.48)$$

Finally, for the potential term from (1.1) we have

$$\mathcal{U}(\mathcal{S}^{ij}, \mathcal{S}^{ji}, q_{ij}) = \text{tr}_{12} \left(F_{21}^0(q_{ij}) P_{12} \mathcal{S}_1^{ij} \mathcal{S}_2^{ji} \right) \quad (2.49)$$

and the Hamiltonian (1.1) is of the form:

$$\mathcal{H} = \sum_{i=1}^M \frac{p_i^2}{2} + \frac{1}{2} \sum_{i=1}^M \text{tr}_{12} \left(m_{12}(0) \mathcal{S}_1^{ii} \mathcal{S}_2^{ii} \right) + \sum_{i < j}^M \text{tr}_{12} \left(F_{21}^0(q_{ij}) P_{12} \mathcal{S}_1^{ij} \mathcal{S}_2^{ji} \right). \quad (2.50)$$

In $M = 1$ case \mathcal{H} reproduce the Hamiltonian of the integrable top, while in the $M = 1$ case we obtain the spin Calogero-Moser Hamiltonian (1.16) up to terms containing S_{ii} – they are constant in this case (1.17).

Poisson brackets. The Poisson structure (before reduction (1.28)) consists of the canonical brackets for positions and momenta

$$\{p_i, q_j\} = \delta_{ij}, \quad \{p_i, p_j\} = \{q_i, q_j\} = 0, \quad i = 1 \dots M \quad (2.51)$$

and the linear Poisson-Lie brackets for the \mathcal{S} variables. They are of the form (B.7) but for $\text{Mat}(NM, \mathbb{C})$ case instead of $\text{Mat}(M, \mathbb{C})$ in (B.7). It is convenient to write down these brackets in terms of $\text{Mat}(N, \mathbb{C})$ -valued blocks \mathcal{S}^{ij} . For $i, j, k, l = 1 \dots M$ and $a, b, c, d = 1 \dots N$:

$$\{\mathcal{S}_{ab}^{ij}, \mathcal{S}_{cd}^{kl}\} = \mathcal{S}_{cb}^{kj} \delta^{il} \delta_{ad} - \mathcal{S}_{ad}^{il} \delta^{kj} \delta_{bc} \quad (2.52)$$

or

$$\{\mathcal{S}_1^{ij}, \mathcal{S}_2^{kl}\} = P_{12} \mathcal{S}_1^{kj} \delta^{il} - \mathcal{S}_1^{il} P_{12} \delta^{kj}, \quad (2.53)$$

where P_{12} as before the permutation operator in $\text{Mat}(N, \mathbb{C})^{\otimes 2}$. For the diagonal blocks we have

$$\{\mathcal{S}_1^{ii}, \mathcal{S}_2^{kk}\} = [P_{12}, \mathcal{S}_1^{ii}] \delta^{ik}. \quad (2.54)$$

It is verified directly that

Proposition 2.2 *The Poisson structure (2.51), (2.53) and the Hamiltonian (2.50) provides equations of (2.19)-(2.21), i.e. for the l.h.s. of the Lax equation (2.1) we have*

$$\dot{\mathcal{L}}(z) = \{\mathcal{H}, \mathcal{L}(z)\}. \quad (2.55)$$

2.4 Interacting tops

Suppose the matrix \mathcal{S} is of rank one, i.e. (1.24) is fulfilled. Consider the potential

$$\text{tr}_{12} \left(F_{21}^0(q_{ij}) P_{12} \mathcal{S}_1^{ij} \mathcal{S}_2^{ji} \right) = \sum_{a,b,c,d=1}^N (F_{12}^0(q_{ji}) P_{12})_{ab,cd} \mathcal{S}_{ba}^{ij} \mathcal{S}_{dc}^{ji}. \quad (2.56)$$

The right multiplication of an element $T_{12} = \sum_{i,j,k,l=1}^N T_{ijkl} E_{ij} \otimes E_{kl} \in \text{Mat}(N, \mathbb{C})^{\otimes 2}$ by permutation operator P_{12} yields $T_{ijkl} \rightarrow T_{ilkj}$, i.e.

$$\text{tr}_{23} \left(F_{32}^0(q_{ij}) P_{23} \mathcal{S}_2^{ij} \mathcal{S}_3^{ji} \right) = \sum_{a,b,c,d=1}^N (F_{12}^0(q_{ji}))_{ad,cb} \mathcal{S}_{ba}^{ij} \mathcal{S}_{dc}^{ji}. \quad (2.57)$$

In the rank 1 case we have

$$\mathcal{S}_{ba}^{ij} \mathcal{S}_{dc}^{ji} = \xi_b^i \eta_a^j \xi_d^j \eta_c^i = \mathcal{S}_{bc}^{ii} \mathcal{S}_{da}^{jj}. \quad (2.58)$$

Therefore,

$$\mathrm{tr}_{23} \left(F_{32}^0(q_{ij}) P_{23} \mathcal{S}_2^{ij} \mathcal{S}_3^{ji} \right) = \mathrm{tr}_{12} \left(F_{12}^0(q_{ji}) \mathcal{S}_1^{jj} \mathcal{S}_2^{ii} \right) = \mathrm{tr}_{12} \left(F_{12}^0(q_{ij}) \mathcal{S}_1^{ii} \mathcal{S}_2^{jj} \right). \quad (2.59)$$

The Hamiltonian of interacting tops model acquires the form:

$$\mathcal{H}^{\mathrm{tops}} = \sum_{i=1}^M \frac{p_i^2}{2} + \frac{1}{2} \sum_{i=1}^M \mathrm{tr}_{12} \left(m_{12}(0) \mathcal{S}_1^{ii} \mathcal{S}_2^{ii} \right) + \sum_{i < j}^M \mathrm{tr}_{12} \left(F_{12}^0(q_{ij}) \mathcal{S}_1^{ii} \mathcal{S}_2^{jj} \right). \quad (2.60)$$

From the Poisson brackets (2.51), (2.54) we get the corresponding equations of motion:

$$\dot{\mathcal{S}}^{ii} = [\mathcal{S}^{ii}, \mathrm{tr}_2(m_{12}(0) \mathcal{S}_2^{ii})] + \sum_{k:k \neq i}^M [\mathcal{S}^{ii}, \mathrm{tr}_2(F_{12}^0(q_{ik}) \mathcal{S}_2^{kk})], \quad (2.61)$$

$$\dot{p}_i = - \sum_{k:k \neq i}^M \mathrm{tr}_{12} \left(\partial_{q_i} F_{12}^0(q_{ik}) \mathcal{S}_1^{ii} \mathcal{S}_2^{kk} \right). \quad (2.62)$$

In this model we are left with M matrix variables $\mathcal{S}^{ii} \in \mathrm{Mat}(N, \mathbb{C})$ of rank one. It is notable that the spin part of the phase space (1.29) is isomorphic to a product of M minimal coadjoint orbits (1.15):

$$\mathcal{O}_{NM}^{\min} // \mathfrak{H}'_{NM} \cong \underbrace{O_N^{\min} \times \dots \times O_N^{\min}}_{M \text{ times}}. \quad (2.63)$$

Notice that the orbits O_N^{\min} come from the constraints conditions (2.18). Hence it appears that

1. For the model of interacting tops the constraints (2.18) play the role of fixation of the Casimir functions for M copies of \mathfrak{gl}_N^* (of rank one). Consequently, equations of motion (2.61) are not changed after reduction. For the $N = 1$ case (the spin Calogero-Moser model) we get $\dot{\mathcal{S}}_{ii} = 0$ since the r.h.s. of (2.61) consists of commutators.

2. The model of interacting tops is formulated in terms of M $\mathrm{Mat}(N, \mathbb{C})$ -valued variables of rank one, describing the minimal coadjoint orbits. The integrability condition is that all Casimir functions $\mathrm{tr}(\mathcal{S}^{ii})$ are equal to each other³.

3. The spin part of the phase space for the model of interacting tops coincides with the phase space of GL_N classical spin chain on M sites with the spins described by minimal coadjoint orbits at each site.

Let us also remark that the top like models with matrix-valued variables were studied in [22, 37] and [6]. In contrast to these papers here we deal with the models, where the matrix variables have their own internal dynamics.

³More precisely, we can not confirm that the model is not integrable in the case $\mathrm{tr}(\mathcal{S}^{ii}) \neq \mathrm{tr}(\mathcal{S}^{jj})$, but the presented Lax pair does not work in this case.

3 Classical r -matrix

In this Section we describe the classical r -matrix structure for the Lax matrix (2.16). Since $\mathcal{L} \in \text{Mat}(NM, \mathbb{C})$ then the corresponding classical \mathfrak{gl}_{NM} r -matrix $\mathbf{r} \in \text{Mat}(NM, \mathbb{C})^{\otimes 2}$. Recall that for the Lax matrix we use the matrix basis (2.15), in which $\mathcal{L} \in \text{Mat}(M, \mathbb{C}) \otimes \text{Mat}(N, \mathbb{C})$. Let the $\text{Mat}(M, \mathbb{C})$ -valued tensor components be numbered by primed numbers, and the $\text{Mat}(N, \mathbb{C})$ -valued components – without primes (as before). Introduce the following r -matrix:

$$\mathbf{r}_{1'2'12}(z, w) = \sum_{i=1}^M \overset{1'}{E}_{ii} \otimes \overset{2'}{E}_{ii} \otimes r_{12}(z-w) + \sum_{i \neq j}^M \overset{1'}{E}_{ij} \otimes \overset{2'}{E}_{ji} \otimes R_{12}^{z-w}(q_{ij}) P_{12}, \quad (3.1)$$

so that $\mathbf{r}_{1'2'12} \in \text{Mat}(M, \mathbb{C})^{\otimes 2} \otimes \text{Mat}(N, \mathbb{C})^{\otimes 2}$. In the case $M = 1$ we come to a non-dynamical r -matrix describing the top model, while in the $N = 1$ we reproduce the dynamical r -matrix of the spin Calogero-Moser model (B.9). r -matrices of these type are known in \mathfrak{gl}_{NM} case and can be extended for arbitrary complex semisimple Lie algebras [10, 11, 24]. In the elliptic case (3.1) is known in the quantum case as well [25]. At the same time (3.1) includes the cases, which have not been described yet. For instance, the new cases correspond to the rational $R_{12}^z(q)$ -matrix from [20]. Similarly to the Lax equations the construction of the r -matrix (3.1) is based on the associative Yang-Baxter equation (1.8) and its degenerations.

Proposition 3.1 *Consider an R -matrix satisfying the associative Yang-Baxter equation (1.8), the classical limit (1.9) and the set of properties from the Section 2.1. Then for the Lax pair (2.15)-(2.16) the following classical exchange relation holds true:*

$$\begin{aligned} \{\mathcal{L}_{1'1}(z), \mathcal{L}_{2'2}(w)\} &= [\mathcal{L}_{1'1}(z), \mathbf{r}_{1'2'12}(z, w)] - [\mathcal{L}_{2'2}(w), \mathbf{r}_{2'1'21}(w, z)] - \\ &\quad - \sum_{k=1}^M \text{tr}(\mathcal{S}^{kk}) \partial_{q_k} \mathbf{r}_{1'2'12}(z, w), \end{aligned} \quad (3.2)$$

where

$$\mathcal{L}_{1'1}(z) = \sum_{i,j=1}^M E_{ij} \otimes 1_M \otimes \mathcal{L}^{ij}(z) \otimes 1_N, \quad (3.3)$$

$$\mathcal{L}_{2'2}(w) = \sum_{k,l=1}^M 1_M \otimes E_{kl} \otimes 1_N \otimes \mathcal{L}^{kl}(w). \quad (3.4)$$

The Poisson brackets in the l.h.s. of (3.2) are given by (2.51)-(2.54).

Proof: The proof is direct. Let us demonstrate how to verify (3.2) for several components of $\overset{1'}{E}_{ij} \otimes \overset{2'}{E}_{kl}$, which are similar to those considered in (B.11)-(B.16) for the spin Calogero-Moser model.

the tensor component $\overset{1'}{E}_{ij} \otimes \overset{2'}{E}_{jk}$ ($i \neq j, j \neq k, i \neq k$):

l.h.s. of (3.2):

$$\text{tr}_{34} \left(R_{13}^z(q_{ij}) P_{13} R_{24}^w(q_{jk}) P_{24} \{ \mathcal{S}_3^{ij}, \mathcal{S}_4^{jk} \} \right) = -\text{tr}_3 \left(\mathcal{S}_3^{ik} R_{23}^w(q_{jk}) P_{23} R_{13}^z(q_{ij}) P_{13} \right). \quad (3.5)$$

r.h.s. of (3.2):

$$\mathrm{tr}_3 \left(\mathcal{S}_3^{ik} (R_{13}^z(q_{ik}) P_{13} R_{12}^{z-w}(q_{kj}) P_{12} - R_{12}^{z-w}(q_{ij}) P_{12} R_{23}^w(q_{ik}) P_{23}) \right). \quad (3.6)$$

Expressions (3.5) and (3.6) coincide due to (1.8) and (2.4).

the tensor component $E_{ii}^{1'} \otimes E_{ij}^{2'}$ ($i \neq j$):

l.h.s. of (3.2):

$$\begin{aligned} & \mathrm{tr}_4 \left(\{p_i, R_{24}^w(q_{ij})\} P_{24} \mathcal{S}_4^{ij} \right) + \mathrm{tr}_{34} \left(r_{13}(z) R_{24}^w(q_{ij}) P_{24} \{ \mathcal{S}_3^{ii}, \mathcal{S}_4^{ij} \} \right) = \\ & = \mathrm{tr}_3 \left(\mathcal{S}_3^{ij} \partial_{q_i} R_{23}^w(q_{ij}) P_{23} \right) - \mathrm{tr}_3 \left(\mathcal{S}_3^{ij} R_{23}^w(q_{ij}) P_{23} r_{13}(z) \right). \end{aligned} \quad (3.7)$$

r.h.s. of (3.2):

$$\mathrm{tr}_3 \left(\mathcal{S}_3^{ij} (R_{13}^w(q_{ij}) P_{13} R_{12}^{z-w}(q_{ji}) P_{12} - r_{12}(z-w) R_{23}^w(q_{ij}) P_{23}) \right) = \quad (3.8)$$

Expressions (3.7) and (3.8) coincide due to (2.32) rewritten through the Fourier symmetry (2.6) as

$$R_{ac}^{q_{ij}}(z) R_{bc}^{q_{ji}}(w) = -R_{ab}^{q_{ij}}(z-w) r_{ac}(z) + r_{bc}(w) R_{ab}^{q_{ij}}(z-w) - \partial_{q_i} R_{ab}^{q_{ij}}(z-w) \quad (3.9)$$

for distinct a, b, c .

the tensor component $E_{ij}^{1'} \otimes E_{ji}^{2'}$ ($i \neq j$):

l.h.s. of (3.2):

$$\mathrm{tr}_3 \left(\mathcal{S}_3^{jj} R_{13}^z(q_{ij}) P_{13} R_{23}^w(q_{ji}) P_{23} - \mathcal{S}_3^{ii} R_{23}^w(q_{ji}) P_{23} R_{13}^z(q_{ij}) P_{13} \right). \quad (3.10)$$

r.h.s. of (3.2):

$$\begin{aligned} & \mathrm{tr}_3 \left(\mathcal{S}_3^{ii} (r_{13}(z) R_{12}^{z-w}(q_{ij}) P_{12} - R_{12}^{z-w}(q_{ij}) P_{12} r_{23}(w)) \right) + \\ & + \mathrm{tr}_3 \left(\mathcal{S}_3^{jj} (-R_{12}^{z-w}(q_{ij}) P_{12} r_{13}(z) + r_{23}(w) R_{12}^{z-w}(q_{ij}) P_{12}) \right) - \\ & - \mathrm{tr}_3 \left(\left(\mathcal{S}_3^{ii} - \mathcal{S}_3^{jj} \right) \partial_{q_i} R_{12}^{z-w}(q_{ij}) P_{12} \right). \end{aligned} \quad (3.11)$$

The last term comes from the second line of (3.2). Again, expressions (3.10) and (3.11) coincide due to (3.9).

The rest of the components are verified similarly. ■

4 Examples

4.1 Elliptic models

Let us begin with the elliptic model [35, 25, 14]. The Lax pair is of the form:

$$\mathcal{L}(z) = \sum_{i,j=1}^M E_{ij} \otimes \mathcal{L}^{ij}(z), \quad \mathcal{L}^{ij}(z) \in \mathrm{Mat}_N \quad \mathcal{L}(z) \in \mathrm{Mat}_{NM}, \quad (4.1)$$

where

$$\begin{aligned} \mathcal{L}^{ij}(z) &= \delta_{ij} \left(p_i 1_N + \mathcal{S}_{(0,0)}^{ii} 1_N E_1(z) + \sum_{\alpha \neq 0} \mathcal{S}_{\alpha}^{ii} T_{\alpha} \varphi_{\alpha}(z, \omega_{\alpha}) \right) + \\ &+ (1 - \delta_{ij}) \sum_{\alpha} \mathcal{S}_{\alpha}^{ij} T_{\alpha} \varphi_{\alpha}(z, \omega_{\alpha} + \frac{q_{ij}}{N}), \end{aligned} \quad (4.2)$$

where the basis (C.8) in $\text{Mat}(N, \mathbb{C})$ is used. Similarly, the M -matrix is of the form

$$\begin{aligned} \mathcal{M}^{ij}(z) &= \delta_{ij} \mathcal{S}_{(0,0)}^{ii} \frac{E_1^2(z) - \wp(z)}{2N} 1_N + \frac{1}{N} \delta_{ij} \sum_{\alpha \neq 0} \mathcal{S}_{\alpha}^{ii} T_{\alpha} f_{\alpha}(z, \omega_{\alpha}) + \\ &+ \frac{1}{N} (1 - \delta_{ij}) \sum_{\alpha} \mathcal{S}_{\alpha}^{ij} T_{\alpha} f_{\alpha}(z, \omega_{\alpha} + \frac{q_{ij}}{N}). \end{aligned} \quad (4.3)$$

These formulae can be obtained from (2.15)-(2.17) and the R -matrix (C.15) together with (A.8).

The Lax equations hold on the constraints

$$\mathcal{S}_{(0,0)}^{ii} = \text{const}, \quad \forall i. \quad (4.4)$$

Instead of the standard basis (1.3) here we use the basis (C.8) for each $N \times N$ block. Then the Poisson structure (1.5) takes the form

$$\{\mathcal{S}_{\alpha}^{ij}, \mathcal{S}_{\beta}^{kl}\} = \delta_{il} \kappa_{\alpha, \beta} \mathcal{S}_{\alpha+\beta}^{kj} - \delta_{kj} \kappa_{\beta, \alpha} \mathcal{S}_{\alpha+\beta}^{il}, \quad (4.5)$$

where $\kappa_{\alpha, \beta}$ are the constants from (C.10).

The Hamiltonian easily follows from $\frac{1}{2N} \text{tr} \mathcal{L}^2(z) = \frac{1}{2N} E_2(z) \text{tr}(\mathcal{S}^2) + \mathcal{H}$ due to (C.11) and (A.8):

$$\mathcal{H} = \frac{1}{2} \sum_{i=1}^M p_i^2 - \frac{1}{2} \sum_{i=1}^M \sum_{\alpha \neq 0} \mathcal{S}_{\alpha}^{ii} \mathcal{S}_{-\alpha}^{ii} E_2(\omega_{\alpha}) - \frac{1}{2} \sum_{i \neq j}^M \sum_{\alpha} \mathcal{S}_{\alpha}^{ij} \mathcal{S}_{-\alpha}^{ji} E_2(\omega_{\alpha} + \frac{q_{ij}}{N}). \quad (4.6)$$

Let us show how this Hamiltonian is reproduced from the general formula (2.50). In order to get the second term in (4.6) one should substitute $m_{12}(0)$ into (2.50) from (C.17) and use relation (A.12). For evaluation of the last sum in (2.50) we need to calculate $F_{12}^0(q) P_{12}$. The answer for $F_{12}^0(q)$ is given in (C.18). Multiply it by $P_{12} = (1/N) \sum_b T_b \otimes T_{-b}$ from the left:

$$\begin{aligned} F_{12}^0(q) P_{12} &= -\frac{1}{N^2} E_2(q) \sum_b T_b \otimes T_{-b} \\ &+ \frac{1}{N^2} \sum_{a \neq (0,0); b} \varphi_a(q, \omega_a) (E_1(q + \omega_a) - E_1(q) + 2\pi i \partial_{\tau} \omega_a) \kappa_{a,b}^2 T_{a+b} \otimes T_{-a-b}. \end{aligned} \quad (4.7)$$

Let us redefine the summation index $b \rightarrow b - a$ in the last sum. Since $\kappa_{a,b} = \kappa_{a,b-a}$ we have

$$\begin{aligned} F_{12}^0(q) P_{12} &= \\ &= \frac{1}{N^2} \sum_b T_b \otimes T_{-b} \left(-E_2(q) + \sum_{a \neq 0; b} \varphi_a(q, \omega_a) (E_1(q + \omega_a) - E_1(q) + 2\pi i \partial_{\tau} \omega_a) \kappa_{a,b}^2 \right) \stackrel{(C.22)}{=} \\ &= -\frac{1}{N^2} \sum_b T_b \otimes T_{-b} E_2(\omega_b + \frac{q}{N}). \end{aligned} \quad (4.8)$$

Finally,

$$\mathrm{tr}_{12} \left(F_{21}^0(q_{ij}) P_{12} \mathcal{S}_1^{ij} \mathcal{S}_2^{ji} \right) = \sum_{\alpha} \mathcal{S}_{\alpha}^{ij} \mathcal{S}_{-\alpha}^{ji} E_2 \left(\omega_{\alpha} + \frac{q_{ij}}{N} \right). \quad (4.9)$$

In the rank 1 case the answer for the Hamiltonian is given by (2.60). Plugging (C.18) into (2.60) we get

$$\begin{aligned} \mathcal{H}^{\mathrm{tops}} &= \frac{1}{2} \sum_{i=1}^M p_i^2 - \frac{1}{2} \sum_{i=1}^M \sum_{\alpha \neq 0} \mathcal{S}_{\alpha}^{ii} \mathcal{S}_{-\alpha}^{ii} E_2(\omega_{\alpha}) - \\ &- \frac{N}{2} \sum_{i \neq j}^M \left(E_2(q_{ij}) \mathcal{S}_0^{ii} \mathcal{S}_0^{jj} - \sum_{a \neq 0} \varphi_a(q_{ij}, \omega_a) (E_1(q_{ij} + \omega_a) - E_1(q_{ij}) + 2\pi i \partial_{\tau} \omega_{\alpha}) \mathcal{S}_{-\alpha}^{ii} \mathcal{S}_{\alpha}^{jj} \right). \end{aligned} \quad (4.10)$$

Let us show how the latter expression appears from (4.6). In the rank one case using (C.11) (so that $\mathcal{S}_{\alpha}^{ij} = \mathrm{tr}(\mathcal{S}^{ij} T_{-\alpha})/N$) we get

$$\mathcal{S}_{\alpha}^{ij} \mathcal{S}_{-\alpha}^{ji} = \frac{\mathrm{tr}(\eta^j T_{-\alpha} \xi^i) \mathrm{tr}(\eta^i T_{\alpha} \xi^j)}{N^2} = \frac{\mathrm{tr}(\eta^j T_{-\alpha} \xi^i \eta^i T_{\alpha} \xi^j)}{N^2} = \frac{\mathrm{tr}(\mathcal{S}^{ii} T_{\alpha} \mathcal{S}^{jj} T_{-\alpha})}{N^2}. \quad (4.11)$$

In this way the Hamiltonian (4.6) acquires the form

$$\begin{aligned} \mathcal{H}^{\mathrm{tops}} &= \frac{1}{2} \sum_{i=1}^M p_i^2 - \frac{1}{2} \sum_{i=1}^M \sum_{\alpha \neq 0} \mathcal{S}_{\alpha}^{ii} \mathcal{S}_{-\alpha}^{ii} E_2(\omega_{\alpha}) - \\ &- \frac{1}{2} \sum_{i \neq j}^M \sum_{\alpha} \frac{\mathrm{tr}(\mathcal{S}^{ii} T_{\alpha} \mathcal{S}^{jj} T_{-\alpha})}{N^2} E_2 \left(\omega_{\alpha} + \frac{q_{ij}}{N} \right), \end{aligned} \quad (4.12)$$

which is the model of interacting tops of (1.26) type. The last terms in (4.12) can be simplified in the following way. Substitute $\mathcal{S}^{ii} = \sum_{\gamma} \mathcal{S}_{\gamma}^{ii} T_{\gamma}$ and $\mathcal{S}^{jj} = \sum_{\mu} \mathcal{S}_{\mu}^{jj} T_{\mu}$ into (4.12). It follows from (C.10)-(C.11) that

$$\mathrm{tr}(T_{\gamma} T_{\alpha} T_{\mu} T_{-\alpha}) = N \kappa_{\alpha, \mu}^2 \delta_{\mu + \gamma}. \quad (4.13)$$

Therefore,

$$\sum_{\alpha} \frac{\mathrm{tr}(\mathcal{S}^{ii} T_{\alpha} \mathcal{S}^{jj} T_{-\alpha})}{N^2} E_2 \left(\omega_{\alpha} + \frac{q_{ij}}{N} \right) = \frac{1}{N} \sum_{\alpha, \mu} \mathcal{S}_{-\mu}^{ii} \mathcal{S}_{\mu}^{jj} E_2 \left(\omega_{\alpha} + \frac{q_{ij}}{N} \right) \kappa_{\alpha, \mu}^2. \quad (4.14)$$

Using (C.20)-(C.21) and summing up over α we obtain the last term in (4.10).

4.2 Trigonometric models

The general classification of the unitary trigonometric R -matrices satisfying associative Yang-Baxter equation was given in [30]. It includes the 7-vertex deformation [8] of the 6-vertex R -matrix and its GL_N generalizations such as the non-standard R -matrix [2]. The integrable tops and related structures based on these R -matrices were described in [17].

Here we restrict ourselves to the case $N = 2$. The 7-vertex R -matrix is of the following form:

$$R_{12}^h(z) = \begin{pmatrix} \coth(z) + \coth(\hbar) & 0 & 0 & 0 \\ 0 & \sinh^{-1}(\hbar) & \sinh^{-1}(z) & 0 \\ 0 & \sinh^{-1}(z) & \sinh^{-1}(\hbar) & 0 \\ C \sinh(z + \hbar) & 0 & 0 & \coth(z) + \coth(\hbar) \end{pmatrix} \quad (4.15)$$

where C is a constant. In the limit $C \rightarrow 0$ the lower left-hand corner vanishes and we get the 6-vertex XXZ R -matrix. For the classical r -matrix and its derivative ($F_{12}^0(z) = \partial_z r_{12}(z)$) we have

$$r_{12}(z) = \begin{pmatrix} \coth(z) & 0 & 0 & 0 \\ 0 & 0 & \sinh^{-1}(z) & 0 \\ 0 & \sinh^{-1}(z) & 0 & 0 \\ C \sinh(z) & 0 & 0 & \coth(z) \end{pmatrix} \quad (4.16)$$

and

$$F_{12}^0(q) = \begin{pmatrix} -\frac{1}{\sinh^2(q)} & 0 & 0 & 0 \\ 0 & 0 & -\frac{\cosh(q)}{\sinh^2(q)} & 0 \\ 0 & -\frac{\cosh(q)}{\sinh^2(q)} & 0 & 0 \\ C \cosh(q) & 0 & 0 & -\frac{1}{\sinh^2(q)} \end{pmatrix} \quad (4.17)$$

respectively. The Fourier transformed F^0 matrix is of the form:

$$F_{12}^0(q)P_{12} = \begin{pmatrix} -\frac{1}{\sinh^2(q)} & 0 & 0 & 0 \\ 0 & -\frac{\cosh(q)}{\sinh^2(q)} & 0 & 0 \\ 0 & 0 & -\frac{\cosh(q)}{\sinh^2(q)} & 0 \\ C \cosh(q) & 0 & 0 & -\frac{1}{\sinh^2(q)} \end{pmatrix} \quad (4.18)$$

From the latter matrix using (1.32) we obtain

$$\begin{aligned} \mathcal{U}(\mathcal{S}^{ij}, \mathcal{S}^{ji}, q_i - q_j) &= \text{tr}_{12} \left(\partial_{q_i} r_{21}(q_{ij}) P_{12} \mathcal{S}_1^{ij} \mathcal{S}_2^{ji} \right) = \\ &= -\frac{1}{\sinh^2(q_{ij})} \left(\mathcal{S}_{11}^{ij} \mathcal{S}_{11}^{ji} + \mathcal{S}_{22}^{ij} \mathcal{S}_{22}^{ji} \right) - \frac{\cosh(q_{ij})}{\sinh^2(q_{ij})} \left(\mathcal{S}_{11}^{ij} \mathcal{S}_{22}^{ji} + \mathcal{S}_{22}^{ij} \mathcal{S}_{11}^{ji} \right) + C \cosh(q_{ij}) \mathcal{S}_{12}^{ij} \mathcal{S}_{12}^{ji}. \end{aligned} \quad (4.19)$$

Similarly, using (1.33) and (4.17) we get the potential for the model of interacting tops:

$$\begin{aligned} \mathcal{V}(\mathcal{S}^{ii}, \mathcal{S}^{jj}, q_i - q_j) &= \text{tr}_{12} \left(\partial_{q_i} r_{12}(q_{ij}) \mathcal{S}_1^{ii} \mathcal{S}_2^{jj} \right) = \\ &= -\frac{1}{\sinh^2(q_{ij})} \left(\mathcal{S}_{11}^{ii} \mathcal{S}_{11}^{jj} + \mathcal{S}_{22}^{ii} \mathcal{S}_{22}^{jj} \right) - \frac{\cosh(q_{ij})}{\sinh^2(q_{ij})} \left(\mathcal{S}_{12}^{ii} \mathcal{S}_{21}^{jj} + \mathcal{S}_{21}^{ii} \mathcal{S}_{12}^{jj} \right) + C \cosh(q_{ij}) \mathcal{S}_{12}^{ii} \mathcal{S}_{12}^{jj}. \end{aligned} \quad (4.20)$$

The top Hamiltonian $\mathcal{H}^{\text{top}}(\mathcal{S}^{ii})$ entering (1.1) or (1.26) is of the form:

$$\mathcal{H}^{\text{top}}(\mathcal{S}^{ii}) = \frac{1}{2} \left((\mathcal{S}_{11}^{ii})^2 + (\mathcal{S}_{22}^{ii})^2 \right) + C(\mathcal{S}_{12}^{ii})^2. \quad (4.21)$$

4.3 Rational models

The rational R -matrices satisfying the required properties are represented by the 11-vertex deformation [8] of the 6-vertex XXX (Yang's) R -matrix. Its higher rank analogues were derived in [34] and [20]. As in trigonometric case here we restrict ourselves to the case $N = 2$. The 11-vertex R -matrix is of the following form:

$$R_{12}^h(z) = \begin{pmatrix} \hbar^{-1} + z^{-1} & 0 & 0 & 0 \\ -\hbar - z & \hbar^{-1} & z^{-1} & 0 \\ -\hbar - z & z^{-1} & \hbar^{-1} & 0 \\ -\hbar^3 - 2z\hbar^2 - 2\hbar z^2 - z^3 & \hbar + z & \hbar + z & \hbar^{-1} + z^{-1} \end{pmatrix} \quad (4.22)$$

In order to get the XXX R -matrix one may take the limit $\lim_{\epsilon \rightarrow 0} \epsilon^{-1} R^{\epsilon h}(\epsilon z)$.

The classical r -matrix, the F_{12}^0 matrix and its Fourier dual are of the form:

$$r_{12}(z) = \begin{pmatrix} z^{-1} & 0 & 0 & 0 \\ -z & 0 & z^{-1} & 0 \\ -z & z^{-1} & 0 & 0 \\ -z^3 & z & z & z^{-1} \end{pmatrix} \quad (4.23)$$

$$F_{12}^0(q) = \begin{pmatrix} -q^{-2} & 0 & 0 & 0 \\ -1 & 0 & -q^{-2} & 0 \\ -1 & -q^{-2} & 0 & 0 \\ -3q^2 & 1 & 1 & -q^{-2} \end{pmatrix} \quad (4.24)$$

$$F_{12}^0(q)P_{12} = \begin{pmatrix} -q^{-2} & 0 & 0 & 0 \\ -1 & -q^{-2} & 0 & 0 \\ -1 & 0 & -q^{-2} & 0 \\ -3q^2 & 1 & 1 & -q^{-2} \end{pmatrix} \quad (4.25)$$

From (4.25) using (1.32) we obtain

$$\begin{aligned} \mathcal{U}(\mathcal{S}^{ij}, \mathcal{S}^{ji}, q_i - q_j) &= -\frac{1}{(q_i - q_j)^2} \left(\mathcal{S}_{11}^{ij} \mathcal{S}_{11}^{ji} + \mathcal{S}_{22}^{ij} \mathcal{S}_{22}^{ji} + \mathcal{S}_{11}^{ij} \mathcal{S}_{22}^{ji} + \mathcal{S}_{22}^{ij} \mathcal{S}_{11}^{ji} \right) + \\ &+ \mathcal{S}_{12}^{ij} \mathcal{S}_{22}^{ji} + \mathcal{S}_{22}^{ij} \mathcal{S}_{12}^{ji} - \mathcal{S}_{12}^{ij} \mathcal{S}_{11}^{ji} - \mathcal{S}_{11}^{ij} \mathcal{S}_{12}^{ji} - 3(q_i - q_j)^2 \mathcal{S}_{12}^{ij} \mathcal{S}_{12}^{ji}. \end{aligned} \quad (4.26)$$

Similarly, from (4.24) using (1.33) we obtain

$$\begin{aligned} \mathcal{V}(\mathcal{S}^{ii}, \mathcal{S}^{jj}, q_i - q_j) &= -\frac{1}{(q_i - q_j)^2} \left(\mathcal{S}_{11}^{ii} \mathcal{S}_{11}^{jj} + \mathcal{S}_{22}^{ii} \mathcal{S}_{22}^{jj} + \mathcal{S}_{12}^{ii} \mathcal{S}_{21}^{jj} + \mathcal{S}_{21}^{ii} \mathcal{S}_{12}^{jj} \right) + \\ &+ \mathcal{S}_{12}^{ii} \mathcal{S}_{22}^{jj} + \mathcal{S}_{22}^{ii} \mathcal{S}_{12}^{jj} - \mathcal{S}_{12}^{ii} \mathcal{S}_{11}^{jj} - \mathcal{S}_{11}^{ii} \mathcal{S}_{12}^{jj} - 3(q_i - q_j)^2 \mathcal{S}_{12}^{ii} \mathcal{S}_{12}^{jj}. \end{aligned} \quad (4.27)$$

The top Hamiltonian $\mathcal{H}^{\text{top}}(\mathcal{S}^{ii})$ entering (1.1) or (1.26) is of the form:

$$\mathcal{H}^{\text{top}}(\mathcal{S}^{ii}) = \mathcal{S}_{12}^{ii}(\mathcal{S}_{22}^{ii} - \mathcal{S}_{11}^{ii}). \quad (4.28)$$

5 Discussion

A possible application of the obtained family of integrable models is in constructing quantum integrable anisotropic long-rang spin chains. The basic idea is that such spin chains appear from the models of interacting tops by the so-called freezing trick likewise the Haldane-Shastry-Inozemtsev spin chains [15] come from the ordinary spin Calogero-Moser-Sutherland models. A direct quantization of the interacting tops is a separate problem, which will be discussed elsewhere. At the same time the quantum Hamiltonian of interacting tops appears in the so-called R -matrix-valued Lax pairs for the (classical) spinless Calogero-Moser model [21, 31, 14]. These are the Lax pairs in a large space $\text{Mat}(M, \mathbb{C}) \otimes \text{Mat}(N, \mathbb{C})^{\otimes M}$:

$$\mathcal{L}^{\text{CM}} = \sum_{a,b=1}^M E_{ab} \otimes \mathcal{L}_{ab}, \quad \mathcal{L}_{ab} = \delta_{ab} p_a 1_N^{\otimes M} + \nu(1 - \delta_{ab}) R_{ab}^z, \quad R_{ab}^z = R_{ab}^z(q_a - q_b). \quad (5.1)$$

and similarly for the accompany M -matrix

$$\mathcal{M}_{ab}^{\text{CM}} = \nu \delta_{ab} d_a + \nu(1 - \delta_{ab}) F_{ab}^z + \nu \delta_{ab} \mathcal{F}^0, \quad F_{ab}^z = \partial_{q_a} R_{ab}^z(q_a - q_b), \quad (5.2)$$

where

$$d_a = - \sum_{c: c \neq a}^M F_{ac}^0, \quad \mathcal{F}^0 = \sum_{b,c: b > c}^M F_{bc}^0 = \sum_{b,c: b > c}^M \partial_{q_b} r_{bc}(q_{bc}). \quad (5.3)$$

In the $N = 1$ case this Lax pair coincides with the widely known Krichever's result [18] for \mathfrak{gl}_M Calogero-Moser model. The last term \mathcal{F}^0 in (5.2) enters \mathcal{M} as a scalar (it is an identity matrix in $\text{Mat}(M, \mathbb{C})$ component) in the auxiliary space $\text{Mat}(M, \mathbb{C})$. Therefore, it can be moved to the l.h.s. of the Lax equation. This yields

$$\{H^{\text{CM}}, \mathcal{L}^{\text{CM}}\} + [\nu \mathcal{F}^0, \mathcal{L}^{\text{CM}}(z)] = [\mathcal{L}^{\text{CM}}(z), \bar{\mathcal{M}}^{\text{CM}}(z)], \quad (5.4)$$

where $\bar{\mathcal{M}}^{\text{CM}} = \mathcal{M}^{\text{CM}} - \nu 1_M \otimes \mathcal{F}^0$. On the one hand (5.4) is just a rewritten classical Lax equation for the spinless Calogero-Moser model. On the other hand we may treat it as half-quantum Lax equation in a sense that the dynamics is given by the interacting tops Hamiltonian (2.60), where the spin variables are already quantized, while the positions and momenta remain classical. Indeed, the quantization of \mathcal{S}_1^{ii} in fundamental representation of GL_N is given by the permutation operator P_{1j} . Plugging it into the potential of (2.60) we get the \mathcal{F}^0 term from (5.2) and (5.4).

Thus the R -matrix valued Lax pairs are multidimensional classical Lax pairs for the spinless Calogero-Moser models and at the same time they are quantum Lax pairs for the models of interacting tops with the spin variables being quantized in the fundamental representation of GL_N , i.e. the \mathcal{F}^0 term is the quantization of the potential $\mathcal{V}(\mathcal{S}^{ii}, \mathcal{S}^{jj}, q_i - q_j)$ (1.33).

Let us also mention that there is another class of integrable models with the Hamiltonian of type (1.26). These are the Gaudin type models [27]. The corresponding Lax matrix is of size

$M \times M$. It has simple poles at n points on elliptic curve (or its degenerations) with the classical spin variables matrices attached to each point. The number of points is not necessarily equal to M . It is an interesting task to find interrelations between the Gaudin models and the models of interacting tops through the spectral duality [26] based on the rank-size duality transformation.

The classical spinless Calogero-Moser model possesses an equilibrium position, where $p_i = 0$ and $q_i = x_i$ (for example, $x_i = i/M$ [9]). At this point the term $\{H^{\text{CM}}, \mathcal{L}^{\text{CM}}\}$ vanishes from the l.h.s. of (5.4), and we are left with the quantum Lax equation for some long-range (quantum) spin chain. It is an anisotropic generalization [31] of the Haldane-Shastry-Inozemtsev type chains. An open question is which \mathcal{F}^0 provide integrable spin chains? To confirm integrability we need to construct higher Hamiltonians, which commute with each other and with $\mathcal{F}^0(q_i = x_i)$. Taking into account all the above we guess that the model of interacting tops together with the freezing trick (the quantum version of the equilibrium position) can be used to calculate higher spin chain Hamiltonians and to prove their commutativity. For this purpose we need to construct a quantization for the model of interacting tops, which is the subject of our next paper.

Another one intriguing question is to construct relativistic generalization of the models discussed above. While the classical models of relativistic interacting tops are expected to be relatively simple (the block L^{ij} in (2.16) should be replaced by $\text{tr}_2(\mathcal{S}_2^{ij} R_{12}^z(q_{ij} + \eta)P_{12})$) its quantum versions and the related long-range spin chain were not studied yet as well as the corresponding R -matrix valued Lax pairs.

6 Appendix

6.1 A: Definitions and identities

The following set of functions is used in this paper [33]. The first one is the Kronecker function:

$$\phi(\eta, z) = \begin{cases} 1/\eta + 1/z & - \text{rational case,} \\ \coth(\eta) + \coth(z) & - \text{trigonometric case,} \\ \frac{\vartheta'(0)\vartheta(\eta+z)}{\vartheta(\eta)\vartheta(z)} & - \text{elliptic case.} \end{cases} \quad (\text{A.1})$$

Its elliptic version is given in terms of the odd theta-function

$$\vartheta(z) = \sum_{k \in \mathbb{Z}} \exp\left(\pi i \tau \left(k + \frac{1}{2}\right)^2 + 2\pi i \left(z + \frac{1}{2}\right) \left(k + \frac{1}{2}\right)\right) \quad (\text{A.2})$$

on elliptic curve with moduli τ ($\text{Im}(\tau) > 0$). Next are the first Eisenstein (odd) function and the Weierstrass (even) \wp -function:

$$E_1(z) = \begin{cases} 1/z, \\ \coth(z), \\ \vartheta'(z)/\vartheta(z), \end{cases} \quad \wp(z) = \begin{cases} 1/z^2, \\ 1/\sinh^2(z), \\ -\partial_z E_1(z) + \frac{1}{3} \frac{\vartheta'''(0)}{\vartheta'(0)}. \end{cases} \quad (\text{A.3})$$

We also need the derivatives

$$E_2(z) = -\partial_z E_1(z) \quad (\text{A.4})$$

and

$$f(z, q) \equiv \partial_q \phi(z, q) = \phi(z, q)(E_1(z+q) - E_1(q)). \quad (\text{A.5})$$

The one (A.4) is the second Eisenstein function.

The main relation is the Fay trisecant identity:

$$\phi(z, q)\phi(w, u) = \phi(z - w, q)\phi(w, q + u) + \phi(w - z, u)\phi(z, q + u). \quad (\text{A.6})$$

The following degenerations of (A.6) are necessary for the Lax equations and r -matrix structures:

$$\phi(z, x)f(z, y) - \phi(z, y)f(z, x) = \phi(z, x + y)(\wp(x) - \wp(y)), \quad (\text{A.7})$$

$$\phi(\eta, z)\phi(\eta, -z) = \wp(\eta) - \wp(z) = E_2(\eta) - E_2(z). \quad (\text{A.8})$$

Also

$$\begin{aligned} \phi(z, q)\phi(w, q) &= \phi(z + w, q)(E_1(z) + E_1(w) + E_1(q) - E_1(z + w + q)) = \\ &= \phi(z + w, q)(E_1(z) + E_1(w)) - f(z + w, q). \end{aligned} \quad (\text{A.9})$$

The local behavior of the Kronecker function and the first Eisenstein function near its simple pole at $z = 0$ is as follows:

$$\phi(z, u) = \frac{1}{z} + E_1(u) + \frac{z}{2}(E_1^2(u) - \wp(u)) + O(z^2), \quad (\text{A.10})$$

$$E_1(z) = \frac{1}{z} + \frac{z}{3} \frac{\wp'''(0)}{\wp'(0)} + O(z^3). \quad (\text{A.11})$$

From (A.10) and (A.5) it follows that

$$f(0, u) = -E_2(u). \quad (\text{A.12})$$

6.2 B: Spin gl_M Calogero-Moser model

The Lax equations

$$\dot{L}^{\text{spin}}(z) = [L^{\text{spin}}(z), M^{\text{spin}}(z)] \quad (\text{B.1})$$

with the Lax pair

$$L_{ij}^{\text{spin}}(z) = \delta_{ij}(p_i + S_{ii}E_1(z)) + (1 - \delta_{ij})S_{ij}\phi(z, q_{ij}), \quad (\text{B.2})$$

$$M^{\text{spin}}(z)_{ij} = (1 - \delta_{ij})S_{ij}f(z, q_i - q_j). \quad (\text{B.3})$$

provide (after restriction on the constraints (1.17)) equations of motion

$$\dot{q}_i = p_i, \quad \ddot{q}_i = \sum_{j \neq i}^M S_{ij}S_{ji}\wp'(q_i - q_j), \quad (\text{B.4})$$

$$\dot{S}_{ii} = 0, \quad \dot{S}_{ij} = \sum_{k \neq i, j}^M S_{ik}S_{kj}(\wp(q_i - q_k) - \wp(q_j - q_k)), \quad i \neq j. \quad (\text{B.5})$$

The l.h.s. of the Lax equations (B.1) is generated by the Hamiltonian (1.16)

$$\dot{L}^{\text{spin}}(z) = \{H^{\text{spin}}, L^{\text{spin}}(z)\} \quad (\text{B.6})$$

and the linear Poisson-Lie brackets on \mathfrak{gl}_M^* :

$$\{S_{ij}, S_{kl}\} = -S_{il}\delta_{kj} + S_{kj}\delta_{il} \quad \text{or} \quad \{S_1, S_2\} = [S_2, P_{12}]. \quad (\text{B.7})$$

Recall that the Poisson reduction with respect to Cartan action (1.18) is non-trivial. For instance, in the rank 1 case (1.22) such reduction leads to the spinless model (1.23). Explicit expression of the reduced Poisson structure depends on a choice of gauge fixation conditions. The equations of motion (B.4)-(B.5) are not the reduced. They are obtained by a simple restriction. To get the final equations one should perform the Dirac reduction and evaluate the Dirac terms.

The classical r -matrix structure is as follows:

$$\begin{aligned} \{L_1^{\text{spin}}(z), L_2^{\text{spin}}(w)\} &= [L_1^{\text{spin}}(z), r_{12}^{\text{spin}}(z, w)] - [L_2^{\text{spin}}(w), r_{21}^{\text{spin}}(w, z)] - \\ &\quad - \sum_{i \neq j} E_{ij} \otimes E_{ji} (S_{ii} - S_{jj}) f(z - w, q_{ij}) \end{aligned} \quad (\text{B.8})$$

with

$$r_{12}^{\text{spin}}(z, w) = E_1(z - w) \sum_{i=1}^M E_{ii} \otimes E_{ii} + \sum_{i \neq j}^M \phi(z - w, q_{ij}) E_{ij} \otimes E_{ji}. \quad (\text{B.9})$$

Here the linear Poisson brackets (B.7) are assumed as well. The Dirac reduction is not yet performed. However, we can see that the restriction on the constraints (1.17) kills the last term in (B.8), and we are left with the standard linear classical r -matrix structure. It is enough for Poisson commutativity

$$\{\text{tr}(L^k(z)), \text{tr}(L^n(w))\} = 0, \quad \forall k, n \in \mathbb{Z}_+, \quad z, w \in \mathbb{C} \quad (\text{B.10})$$

necessary for the Liouville integrability. The proof of (B.8) is direct. It is based on the identities (A.6)-(A.9). Let us write down a few examples of verification of (B.8):

the tensor component $E_{ij} \otimes E_{jk}$ ($i \neq j, j \neq k, k \neq i$):

l.h.s. of (B.8):

$$\{L_{ij}^{\text{spin}}(z), L_{jk}^{\text{spin}}(w)\} = \{S_{ij}, S_{jk}\} \phi(z, q_{ij}) \phi(w, q_{jk}) = -S_{ik} \phi(z, q_{ij}) \phi(w, q_{jk}). \quad (\text{B.11})$$

r.h.s. of (B.8):

$$S_{ik} \phi(z, q_{ik}) \phi(z - w, q_{kj}) + S_{ik} \phi(w, q_{ik}) \phi(w - z, q_{ji}). \quad (\text{B.12})$$

Expressions (B.11) and (B.12) coincide due to (A.6).

the tensor component $E_{ii} \otimes E_{ij}$ ($i \neq j$):

l.h.s. of (B.8):

$$\begin{aligned} \{L_{ii}^{\text{spin}}(z), L_{ij}^{\text{spin}}(w)\} &= \{p_i, \phi(w, q_{ij})\} S_{ij} + \{S_{ii}, S_{ij}\} E_1(z) \phi(w, q_{ij}) = \\ &= S_{ij} f(w, q_{ij}) - S_{ij} E_1(z) \phi(w, q_{ij}). \end{aligned} \quad (\text{B.13})$$

r.h.s. of (B.8):

$$S_{ij} \phi(z, q_{ij}) \phi(z - w, q_{ji}) + S_{ij} E_1(w - z) \phi(w, q_{ij}). \quad (\text{B.14})$$

Expressions (B.13) and (B.14) coincide due to (A.9).

the tensor component $E_{ij} \otimes E_{ji}$ ($i \neq j$):

l.h.s. of (B.8):

$$\begin{aligned} & \{L_{ij}^{\text{spin}}(z), L_{ji}^{\text{spin}}(w)\} = \\ & = \{S_{ij}, S_{ji}\} \phi(z, q_{ij}) \phi(w, q_{ji}) = (S_{ii} - S_{jj}) \phi(z, q_{ij}) \phi(-w, q_{ij}). \end{aligned} \quad (\text{B.15})$$

The last term from the r.h.s. of (B.8) contributes in this component. The r.h.s. of (B.8):

$$\begin{aligned} & (p_i + S_{ii} E_1(z) - p_j - S_{jj} E_1(z)) \phi(z - w, q_{ij}) - \\ & -(p_j + S_{jj} E_1(w) - p_i - S_{ii} E_1(w)) \phi(w - z, q_{ji}) - (S_{ii} - S_{jj}) f(z - w, q_{ij}) = \\ & = (S_{ii} - S_{jj}) \left((E_1(z) - E_1(w)) \phi(z - w, q_{ij}) - f(z - w, q_{ij}) \right). \end{aligned} \quad (\text{B.16})$$

Expressions (B.15) and (B.16) coincide due to (A.9).

6.3 C: Integrable \mathfrak{gl}_N tops

It was shown in [22] (see also [20]) that the Lax equations

$$\dot{L}(z, S) = [L(z, S), M(z, S)] \quad (\text{C.1})$$

are equivalent to equations

$$\dot{S} = [S, J(S)] \quad (\text{C.2})$$

for the Lax pair

$$L(z, S) = \text{tr}_2(r_{12}(z) S_2), \quad M(z, S) = \text{tr}_2(m_{12}(z) S_2), \quad S_2 = 1_N \otimes S \quad (\text{C.3})$$

and

$$J(S) = \text{tr}_2(m_{12}(0) S_2). \quad (\text{C.4})$$

constructed by means of the coefficients of the (classical limit) expansion (1.9) for an R -matrix satisfying the associative Yang-Baxter equation (1.8) and the properties (2.2)-(2.5). The answer (C.3) can be written more explicitly. For

$$r_{12}(z) = \sum_{i,j,k,l=1}^N r_{ijkl}(z) e_{ij} \otimes e_{kl} \quad (\text{C.5})$$

(C.3) means

$$L(z, S) = \sum_{i,j,k,l=1}^N r_{ijkl}(z) S_{lk} e_{ij} \quad (\text{C.6})$$

since $\text{tr}(e_{kl} S) = S_{lk}$.

Let us briefly describe how these formulae reproduce the elliptic top from [19]. In the elliptic case we need special matrix basis in $\text{Mat}(N, \mathbb{C})$. Consider the matrices

$$Q_{kl} = \delta_{kl} \exp\left(\frac{2\pi i}{N} k\right), \quad \Lambda_{kl} = \delta_{k-l+1=0 \bmod N}, \quad Q^N = \Lambda^N = 1_N. \quad (\text{C.7})$$

Then the basis in $\text{Mat}(N, \mathbb{C})$ is given by the following set:

$$T_a = T_{a_1 a_2} = \exp\left(\frac{\pi i}{N} a_1 a_2\right) Q^{a_1} \Lambda^{a_2}, \quad a = (a_1, a_2) \in \mathbb{Z}_N \times \mathbb{Z}_N. \quad (\text{C.8})$$

Since

$$\exp\left(\frac{2\pi i}{N} a_1 a_2\right) Q^{a_1} \Lambda^{a_2} = \Lambda^{a_2} Q^{a_1} \quad (\text{C.9})$$

we have

$$T_\alpha T_\beta = \kappa_{\alpha, \beta} T_{\alpha+\beta}, \quad \kappa_{\alpha, \beta} = \exp\left(\frac{\pi i}{N} (\beta_1 \alpha_2 - \beta_2 \alpha_1)\right), \quad (\text{C.10})$$

where $\alpha + \beta = (\alpha_1 + \beta_1, \alpha_2 + \beta_2)$. The non-degenerate pairing is given by the matrix trace:

$$\text{tr}(T_\alpha T_\beta) = N \delta_{\alpha+\beta}, \quad T_0 = 1_N. \quad (\text{C.11})$$

Define the set of functions numerated by $a = (a_1, a_2) \in \mathbb{Z}_N \times \mathbb{Z}_N$:

$$\varphi_a(z, \omega_a + u) = \exp(2\pi i \frac{a_2}{N} z) \phi(z, \omega_a + u), \quad \omega_a = \frac{a_1 + a_2 \tau}{N} \quad (\text{C.12})$$

and introduce notation

$$f_a(z, \omega_a + u) = \exp(2\pi i \frac{a_2}{N} z) f(z, \omega_a + u). \quad (\text{C.13})$$

The Baxter-Belavin R -matrix [5] satisfying all required properties including the Fourier symmetry (2.6) is of the form:

$$R_{12}^{BB}(\hbar, z) = \sum_{a \in \mathbb{Z}_N \times \mathbb{Z}_N} \varphi_a(z, \hbar + \omega_a) T_a \otimes T_{-a} \in \text{Mat}(N, \mathbb{C})^{\otimes 2}. \quad (\text{C.14})$$

This R -matrix satisfies required properties but with different normalizations. For example, the Fourier symmetry has form $R_{12}^{BB}(\hbar, z) P_{12} = R_{12}^{BB}(z/N, N\hbar)$ (see the Fourier transformation formulae in [37]). To fulfill all requirements including the normalization (2.5) we consider

$$R_{12}^h(z) = R_{12}^{BB}(\hbar/N, z) = \frac{1}{N} \sum_{a \in \mathbb{Z}_N \times \mathbb{Z}_N} \varphi_a(z, \frac{\hbar}{N} + \omega_a) T_a \otimes T_{-a} \in \text{Mat}(N, \mathbb{C})^{\otimes 2}. \quad (\text{C.15})$$

The corresponding classical r -matrix is as follows

$$r_{12}(z) = \frac{1}{N} E_1(z) 1_N \otimes 1_N + \frac{1}{N} \sum_{a \neq (0,0)} \varphi_a(z, \omega_a) T_a \otimes T_{-a} \in \text{Mat}(N, \mathbb{C})^{\otimes 2}, \quad (\text{C.16})$$

and

$$m_{12}(z) = \frac{E_1^2(z) - \wp(z)}{2N^2} 1_N \otimes 1_N + \frac{1}{N^2} \sum_{a \neq (0,0)} f_a(z, \omega_a) T_a \otimes T_{-a} \in \text{Mat}(N, \mathbb{C})^{\otimes 2}. \quad (\text{C.17})$$

Then the formulae for the Lax pair (C.3) reproduce the Lax pair of the elliptic top. It is contained in the Lax pair (4.1)-(4.3) as a diagonal $N \times N$ block.

The derivative of the classical r -matrix is obtained through (A.5):

$$F_{12}^0(z) = \partial_z r_{12}(z) = -\frac{1}{N} E_2(z) 1_N \otimes 1_N + \frac{1}{N} \sum_{a \neq (0,0)} \varphi_a(z, \omega_a) (E_1(z + \omega_a) - E_1(z) + 2\pi i \partial_\tau \omega_a) T_a \otimes T_{-a}. \quad (\text{C.18})$$

The Fourier symmetry $R_{12}^{\hbar}(z) = R_{12}^z(\hbar) P_{12}$ for the R -matrix (C.15) is based on the following set of identities for the functions (C.12):

$$\frac{1}{N} \sum_{\alpha} \kappa_{\alpha, \gamma}^2 \varphi_{\alpha}(N\hbar, \omega_{\alpha} + \frac{z}{N}) = \varphi_{\gamma}(z, \omega_{\gamma} + \hbar), \quad \forall \gamma \in \mathbb{Z}_N \times \mathbb{Z}_N. \quad (\text{C.19})$$

By degeneration of the latter identities one can deduce (see [37]):

$$\sum_{\alpha} E_2(\omega_{\alpha} + q) = N^2 E_2(Nq) \quad (\text{C.20})$$

and for $\gamma \neq 0$

$$\sum_{\alpha} \kappa_{\alpha, \gamma}^2 E_2(\omega_{\alpha} + q) = -N^2 \varphi_{\gamma}(Nq, \omega_{\gamma}) (E_1(Nq + \omega_{\gamma}) - E_1(Nq) + 2\pi i \partial_\tau \omega_{\gamma}). \quad (\text{C.21})$$

Conversely,

$$-E_2(q) + \sum_{\alpha} \kappa_{\alpha, \gamma}^2 \varphi_{\alpha}(q, \omega_{\alpha}) (E_1(q + \omega_{\alpha}) - E_1(q) + 2\pi i \partial_\tau \omega_{\alpha}) = -E_2(\omega_{\gamma} + \frac{q}{N}). \quad (\text{C.22})$$

Acknowledgments. The work was partially supported by RFBR grants 18-01-00926 (all authors) and 19-51-18006 Bolg_a (I. Sechin). The research of A. Zotov was also supported in part by the HSE University Basic Research Program, Russian Academic Excellence Project '5-100' and by the Young Russian Mathematics award. A. Zotov is also grateful to Sasha Alexandrov for hospitality at the 2-nd Workshop on integrable systems and applications (May, 2019) in IBS Center for Geometry and Physics (Pohang, Korea), where a part of the results was presented.

References

- [1] G. Aminov, S. Arthamonov, A. Smirnov, A. Zotov, J. Phys. A: Math. Theor. 47 (2014) 305207; arXiv:1402.3189. [hep-th].
- [2] A. Antonov, K. Hasegawa, A. Zabrodin, Nucl. Phys. B503 (1997) 747–770; hep-th/9704074.
- [3] V.I. Arnold, Annales de l'institut Fourier, 16:1 (1966) 319–361.
L.A. Dikii, Funct. Anal. Appl. 6:4 (1972) 326–327.
S.V. Manakov, Funct. Anal. Appl., 10:4 (1976) 328–329.
A.S. Mishenko, Funct. Anal. Appl. 4:3 (1970) 232–235.
A.S. Mishenko, A.T. Fomenko, Mathematics of the USSR-Izvestiya, 12:2 (1978) 371–389.

- [4] E. Billey, J. Avan, O. Babelon, *Physics Letters A*, 186 (1994) 114–118; hep-th/9312042.
 E. Billey, J. Avan, O. Babelon, *Physics Letters A*, 188 (1994) 263–271; hep-th/9401117.
 I. Krichever, O. Babelon, E. Billey, M. Talon, *Amer. Math. Soc. Transl. (2) Vol. 170* (1995) 83–120.
- [5] R.J. Baxter, *Ann. Phys.* 70 (1972) 193–228.
 A.A. Belavin, *Nucl. Phys. B*, 180 (1981) 189–200.
 M.P. Richey, C.A. Tracy, *J. Stat. Phys.* 42 (1986) 311–348.
- [6] M. Bertola, M. Cafasso, V. Rubtsov, *Commun. Math. Phys.* 363:2 (2018) 503–530;
 arXiv:1710.00736.
- [7] F. Calogero, *J. Math. Phys.* 10 (1969) 2191–2196.
 F. Calogero, *J. Math. Phys.* 12 (1971) 419–436.
 B. Sutherland, *Physical Review A*, 4:5 (1971) 2019–2021.
 B. Sutherland, *Physical Review A*, 5:3 (1972) 1372–1376.
 J. Moser, *Advances in mathematics* 16 (1975) 197–220.
- [8] I.V. Cherednik, *Theor. Math. Phys.*, 47:2 (1981) 422–425.
- [9] E. Corrigan, R. Sasaki, *J. Phys. A* 35 (2002) 7017–7062; arXiv:hep-th/0204039.
- [10] P. Etingof, O. Schiffmann, *Mathematical Research Letters*, 6 (1999) 593–612;
 arXiv:math/9908115 [math.QA].
 P. Etingof, O. Schiffmann, *London Math. Soc., Lecture Notes Series*, 290 (2001) 89–129;
 arXiv:math.QA/9908064.
- [11] L. Feher, B.G. Puztai, *Nuclear Physics B*, 621 (2002) 622–642; arXiv:math/0109132 [math.QA].
- [12] S. Fomin, A.N. Kirillov, *Advances in geometry; Progress in Mathematics book series, Vol. 172* (1999) 147–182.
 Anatol N. Kirillov, *SIGMA* 12 (2016), 002; arXiv:1502.00426 [math.RT].
- [13] J. Gibbons, T. Hermsen, *Physica D: Nonlinear Phenomena*, 11 (1984) 337–348;
 S. Wojciechowski, *Physics Letters A*, 111 (1985) 101–103.
- [14] A. Grekov, A. Zotov, *J. Phys. A: Math. Theor.* 51 (2018) 315202; arXiv: 1801.00245.
 A.V. Zotov, *Theoret. and Math. Phys.*, 197:3 (2018) 1755–1770.
- [15] F.D.M. Haldane, *Phys. Rev. Lett.* 60 (1988) 635–638.
 B.S. Shastri, *Phys. Rev. Lett.* 60 (1988) 639–642.
 V.I. Inozemtsev, *Journal of Statistical Physics*, 59 (1990) 1143–1155.
 Alexios P. Polychronakos, *Phys. Rev. Lett.* 70 (1993) 2329–2331; arXiv:hep-th/9210109.
- [16] K. Hikami, M. Wadati, *Physics Letters A*, 173 (1993) 263–266.
- [17] T. Krasnov, A. Zotov, *Annales Henri Poincaré*, 20:8 (2019) 2671–2697;
 arXiv:1812.04209 [math-ph].
- [18] I. Krichever, *Funct. Anal. Appl.*, 14:4 (1980) 282–290.
- [19] A. Levin, M. Olshanetsky, A. Zotov, *Commun. Math. Phys.* 236 (2003) 93–133;
 arXiv:nlin/0110045.
- [20] A. Levin, M. Olshanetsky, A. Zotov, *JHEP* 07 (2014) 012; arXiv:1405.7523 [hep-th].
 A. Levin, M. Olshanetsky, A. Zotov, *Nuclear Physics B* 887 (2014) 400–422, arXiv:1406.2995.

- [21] A. Levin, M. Olshanetsky, A. Zotov, JHEP 10 (2014) 109; arXiv:1408.6246 [hep-th].
A.M. Levin, M.A. Olshanetsky, A.V. Zotov, Theoret. and Math. Phys. 184:1 (2015) 924–939; arXiv:1501.07351 [math-ph].
- [22] A. Levin, M. Olshanetsky, A. Zotov, J. Phys. A: Math. Theor. 49:39 (2016) 395202; arXiv:1603.06101.
- [23] A. Levin, M. Olshanetsky, A. Smirnov, A. Zotov, Commun. Math. Phys., 316 (2012) 1–44; arXiv:1006.0702 [math-ph].
A. Levin, M. Olshanetsky, A. Smirnov, A. Zotov, J. Geom. Phys., 62:8 (2012) 1810–1850; arXiv:1007.4127 [math-ph].
- [24] A. Levin, M. Olshanetsky, A. Smirnov, A. Zotov, SIGMA 8 (2012) 095; arXiv:1207.4386 [math-ph].
- [25] A. Levin, M. Olshanetsky, A. Smirnov, A. Zotov, J. Phys. A: Math. Theor. 46:3 (2013) 035201; arXiv:1208.5750 [math-ph]
- [26] A. D. Mironov, A. Yu. Morozov, Y. Zenkevich, A. V. Zotov, JETP Letters, 97:1 (2013) 45-51; arXiv:1204.0913.
A. Mironov, A. Morozov, B. Runov, Y. Zenkevich, A. Zotov, Lett. Math. Phys., 103:3 (2013), 299-329; arXiv:1206.6349.
A. Mironov, A. Morozov, B. Runov, Y. Zenkevich, A. Zotov, JHEP 12 (2013) 034; arXiv:1307.1502.
- [27] N. Nekrasov, Commun. Math. Phys. 180 (1996) 587–604; arXiv:hep-th/9503157.
- [28] A. Polishchuk, Advances in Mathematics 168:1 (2002) 56-95.
- [29] A.P. Polychronakos, Phys. Rev. Lett. 89 (2002) 126403; hep-th/0112141.
A.P. Polychronakos, Nucl. Phys. B543 (1999) 485–498; hep-th/9810211.
A.P. Polychronakos, J. Phys. A: Math. Gen. 39 (2006) 12793; hep-th/0607033.
- [30] T. Schedler, Mathematical Research Letters, 10:3 (2003) 301–321; arXiv:math/0212258 [math.QA].
A. Polishchuk, Algebra, Arithmetic, and Geometry, Progress in Mathematics book series, Volume 270 (2010) 573–617; arXiv:math/0612761 [math.QA].
- [31] I. Sechin, A. Zotov, Phys. Lett. B, 781 (2018) 1-7 , arXiv: 1801.08908.
- [32] E.K. Sklyanin, Fuct. Anal. Appl. 16:4 (1982) 263–270.
A.G. Reyman, M.A. Semenov-Tian-Shansky, Zap. Nauchn. Semin. LOMI, 150 (1986) 104–118.
- [33] A. Weil, *Elliptic functions according to Eisenstein and Kronecker*, Springer-Verlag, (1976).
D. Mumford, *Tata Lectures on Theta I, II*, Birkhäuser, Boston, Mass. (1983, 1984).
- [34] A. Smirnov, Cent. Eur. J. Phys.,8:4 (2010) 542–554.
- [35] A.V. Zotov, A.M. Levin, Theoret. and Math. Phys., 146:1 (2006) 45-52.
- [36] A.V. Zotov, A.V. Smirnov, Theoret. and Math. Phys., 177:1 (2013), 1281-1338.
- [37] A. Zotov, Modern Phys. Lett. A, 32:32 (2017), 1750169 , 22 pp., arXiv: 1706.05601