

ON EIGENVALUES AND EIGENFUNCTIONS OF THE OPERATORS DEFINING MULTIDIMENSIONAL SCALING ON SOME SYMMETRIC SPACES

TIANYU MA AND EUGENE STEPANOV

ABSTRACT. We study asymptotics of the eigenvalues and eigenfunctions of the operators used for constructing multidimensional scaling (MDS) on compact connected Riemannian manifolds, in particular on closed connected symmetric spaces. They are the limits of eigenvalues and eigenvectors of squared distance matrices of an increasing sequence of finite subsets covering the space densely in the limit. We show that for products of spheres and real projective spaces, the numbers of positive and negative eigenvalues of these operators are both infinite. We also find a class of spaces (namely $\mathbb{R}\mathbb{P}^n$ with odd $n > 1$) whose MDS defining operators are not trace class, and original distances cannot be reconstructed from the eigenvalues and eigenfunctions of these operators.

1. INTRODUCTION

A problem frequently encountered in modern data science is that of reconstructing a metric space (X, d) and the Borel measure μ on it just from the information on the distances between points of a sufficiently large finite subset $\Sigma_k := \{x_1^k, \dots, x_k^k\} \subset X$. Here we require the subset Σ_k to cover X almost densely and with a density approximately μ . Of course, unless X is finite itself, no finite set of points will be sufficient to reconstruct the triple (X, d, μ) and one can only hope to do this in the limit as $k \rightarrow \infty$. To be more precise, we suppose to know distances between points of each set Σ_k of some chosen sequence of finite subsets of X , and would like to recover from it the information on (X, d, μ) . This is known as the *learning* problem (or *manifold learning*, when X is a priori supposed to be some smooth, say, Riemannian manifold, and d to be its geodesic distance).

One of the basic algorithms aimed to solve the learning problem and widely used in applications is *multidimensional scaling* (MDS) [21]. Although the latter has been originally proposed only for intrinsically Euclidean data (i.e., when X is a subset of a Euclidean space \mathbb{R}^n and the distance d is Euclidean), it has been extended to generic metric spaces. Moreover, in applicative “folklore”, it is often used not only when the distance d is non-Euclidean, but also when d is merely some symmetric function not necessarily satisfying the triangle inequality (the so-called *dissimilarity* function). Whether this application of MDS is justified, i.e., what will be reconstructed by MDS when d is a non-Euclidean distance has been recently posed and solved in [1] with quite an astonishing answer. Namely, take $X := \mathbb{S}^1$ as a unit circle endowed with its geodesic distance, and let the points of $\Sigma_k \subset X$ to be uniformly spaced so that in the limit as $k \rightarrow \infty$ they cover X uniformly. Then according to [1], MDS yields in the limit as $k \rightarrow \infty$ a closed curve in an

infinite-dimensional space, which is far from being a circle. An easy calculation shows that it is a fractal object, namely, a *snowflake* embedding [18] of a circle in an infinite-dimensional Hilbert space [14]. It becomes an isometric embedding if \mathbb{S}^1 is endowed with the geodesic distance raised to some power $\alpha = 1/2$. Although this may be unexpected in view of various commonly used applications of MDS, an explanation of this fact may be also traced back to the classical work [20] by Neumann and Schoenberg. In their paper, all the invariant metrics on the circle that embed isometrically into a Hilbert space are classified, including of course the 1/2-snowflake re-obtained via MDS, which is actually credited to the earlier work [22], see also the discussion in [9, section 7.3].

1.1. Asymptotics of MDS embeddings. In the study of asymptotical behaviour of the spectra of matrices of squared distances between points of finite samples $\Sigma_k \subset X$, as well as of the embedding maps $\mathcal{M}_k: \Sigma_k \rightarrow \mathbb{R}^k$ produced by MDS, the linear operator T over the space $L^2(X, \mu)$ defined by the following formulae plays an important role.

$$(1.1) \quad \begin{aligned} K(x, y) &:= -\frac{1}{2}d^2(x, y), \\ (\mathcal{K}u)(x) &:= \int_X K(x, y)u(y) d\mu(y), \\ T &:= PKP, \end{aligned}$$

where P is the projector operator to the orthogonal complement of constant functions in $L^2(X, \mu)$. Both \mathcal{K} and T are well-defined under just a mild assumption that μ have a finite 4-th order moment, i.e.

$$(1.2) \quad \int_X d^4(x_0, y) d\mu(y) < \infty$$

for some $x_0 \in X$ (which holds for instance when μ is finite and X is bounded). Moreover, in this case, they are self-adjoint Hilbert–Schmidt (and hence compact) operators. What is more important is the following: Under the same assumption (1.2), suppose the empirical measures μ_k of finite samples $\Sigma_k := \{x_1^k, \dots, x_k^k\}$ defined by

$$\mu_k := \frac{1}{k} \sum_{i=1}^k \delta_{x_i^k},$$

δ_y standing for the Dirac mass concentrated in $y \in X$, converge to μ as $k \rightarrow \infty$ in the Kantorovich¹ 4-distance W_4 , i.e. $\lim_k W_4(\mu_k, \mu) = 0$. Then according to [10, theorem 5.8], the maps \mathcal{M}_k viewed as functions from X to \mathbb{R}^∞ (with \mathbb{R}^k canonically identified with the subspace of \mathbb{R}^∞ having all zero coordinates except the first k coordinates) converge to some map $\mathcal{M}: X \rightarrow \mathbb{R}^\infty$, called further *infinite MDS map*, in measure μ with respect to the product topology on \mathbb{R}^∞ (see also [12]). The latter is given by the formula

$$(1.3) \quad \mathcal{M}(x) := \left(\sqrt{\lambda_1^+} \phi_1^+(x), \sqrt{\lambda_2^+} \phi_2^+(x), \dots, \sqrt{\lambda_j^+} \phi_j^+(x), \dots \right),$$

where $\lambda_1^+ \geq \lambda_2^+ \geq \dots > 0$ are positive eigenvalues of T (counting multiplicity), and $\{\phi_j^+\}_j \in L^2(X, \mu)$ is an orthonormal system in $L^2(X, \mu)$ made of the respective

¹Usually, though historically incorrectly, the distances W_p among probability measures are called Wassertsein distances.

eigenfunctions, i.e. $T\phi_j^+ = \lambda_j^+ \phi_j^+$. Note that the definition of M depends on the choice of ϕ_j^+ . Here we silently assume that if the set of positive eigenvalues of T contains $N < \infty$ elements, then $(M(x))_j := 0$ for $j > N$.

By calculating explicitly the eigenvalues and eigenfunctions of T and using (1.3), one shows in [10] that M gives a snowflake (Assouad-type) embedding of any m -dimensional sphere \mathbb{S}^m or any m -dimensional flat torus $(\mathbb{S}^1)^m$ into the Hilbert space ℓ^2 of square summable sequences (in the calculations one assumes μ to be the respective volume measure in all these cases).

Another important observation is the following: With the stronger assumption that T is a trace-class operator, i.e.,

$$\sum_i |\lambda_i| < +\infty,$$

where $\{\lambda_i\}$ stands for the sequence of *all* eigenvalues of T , let the metric measure space (X, d, μ) be, say, infinitesimally doubling see [5, theorem 3.4.3] (which includes any smooth Riemannian manifold equipped with geodesic distance and volume measure). Then by [10, theorem 5.8] (see also [12]), in a sense, distances between almost every pair of points can be recovered from the spectrum of T and the set of the respective eigenfunctions. Namely, in this case, we have

$$(1.4) \quad \sum_{i=1}^{\infty} \lambda_i (\phi_i(x) - \phi_i(y))^2 = d^2(x, y) \quad \text{for } \mu \otimes \mu\text{-a.e. } (x, y),$$

where $\{\phi_i\}$ stands for an orthonormal basis in $L^2(X, \mu)$ made of eigenfunctions of T with $T\phi_i = \lambda_i\phi_i$. The importance of the trace class condition on the operators T and \mathcal{K} for the asymptotics of the spectra of distance matrices has been also studied recently in [19].

1.2. Questions and results. The above-cited results raise a series of curious questions. Namely, one asks whether there are natural examples of spaces (X, d, μ) such that

- (Q1) no infinite MDS map (i.e., independently on the choice of eigenfunctions of T) gives a topological embedding of X into ℓ^2 ,
- (Q2) the operator T is not trace-class and/or the distance reconstruction formula (1.4) is not valid.

We find both examples among just compact Riemannian manifolds with volume measure, namely, both (Q1) and (Q2) are satisfied by projective spaces of sufficiently high dimension. We do so by studying the operator T , its eigenvalues and eigenfunctions for symmetric compact Riemannian manifolds with volume measure. Note that in general, there seems to be no easy way to find either the spectrum or eigenfunctions of T . However, in this particular case the situation greatly simplifies since we are able to show that T commutes with the Laplace-Beltrami operator, which allows us to search for its eigenfunctions among the eigenfunctions of the latter. We are able then to show that if X is a finite product of spheres of any dimensions, the infinite MDS map gives a snowflake embedding of X into ℓ^2 thus generalizing the results of [10], while if X is a projective space with sufficiently high dimension, then M does not send X to ℓ^2 at all, and in particular the distance reconstruction formula (1.4) is not valid. Curiously however, it happens that in all these cases. Moreover, if X is a finite product of spheres and projective spaces, then the spectrum of T contains infinitely many positive and negative eigenvalues.

This contrasts with the case that (X, d) is isometrically embeddable in a Hilbert space, in which all the eigenvalues of T are positive.

2. NOTATION AND PRELIMINARIES

For vectors x and y in the Euclidean space \mathbb{R}^n , we denote by $x \cdot y$ their Euclidean scalar product. The Euclidean norm is denoted by $|\cdot|$. Let ℓ^2 be the usual Banach space of square summable sequences equipped with its usual norm $\|\cdot\|_2$. The space \mathbb{R}^∞ stands for the linear space of all real-valued sequences (sometimes denoted by $\mathbb{R}^{\mathbb{N}}$ in the literature), equipped with its product topology. The norm $\|\cdot\|_2$ can be extended to a “metric” on \mathbb{R}^∞ taking values in $[0, +\infty]$.

If X is a smooth Riemannian manifold, we denote by $C^\infty(X)$ the set of infinitely smooth functions over X .

Throughout the paper, we sometimes use the big Theta notation by D. Knuth.

For a metric measure space (X, d, μ) , we will assume μ to be a Borel probability measure. By $\langle x, y \rangle$ we denote then the standard scalar product in the Hilbert space $L^2(X, \mu)$. For a $u \in L^2(X, \mu)$ we let u^\perp stand for its orthogonal complement in $L^2(X, \mu)$. The spectrum of a linear operator T counting multiplicity is denoted as $\text{Spec}(T)$, and its signature $\text{sgn}(T)$ consists of three values ordered in the numbers of zero, positive and negative eigenvalues.

Recall for a metric measure space (X, d, μ) , the MDS map $\mathcal{M} : X \rightarrow \mathbb{R}^\infty$ defined as in (1.3) are obtained from the positive eigenvalues and their corresponding eigenfunction the operator T . Such maps are not unique, a different choice of the orthonormal set $\{\phi_j^+\}$ yields a different map. However, they all have the common property given by the following statement.

Lemma 2.1. *If the operator T is Hilbert-Schmidt, then for any MDS maps $\mathcal{M}^1, \mathcal{M}^2$, we have*

$$\|\mathcal{M}^1(x) - \mathcal{M}^1(y)\|_2 = \|\mathcal{M}^2(x) - \mathcal{M}^2(y)\|_2, \quad \text{for all } x, y \in X.$$

Moreover, in this case the right-hand side of (1.4) is independent of the choice of eigenfunctions ϕ_j of \mathcal{K} .

Proof. Since T is compact, every non-zero eigenspace of T is finite-dimensional. A new choice of the orthonormal set of $\{\phi_j^+\}$ is obtained from $L^2(X, \mu)$ -orthogonal transformation of each eigenspace $E_{\lambda_\alpha^+}$, with the eigenvalue $\lambda_\alpha^+ > 0$. Let $\{\phi_j^\alpha\}$ and $\{\tilde{\phi}_j^\alpha\}$ be two orthonormal bases of $E_{\lambda_\alpha^+}$. The space $E_{\lambda_\alpha^+}$ with L^2 -norm can be identified as a Euclidean space with the standard norm, and orthogonal transformation on Euclidean spaces preserves Euclidean distances. Thus, we have

$$\sum_j (\phi_j^\alpha(x) - \phi_j^\alpha(y))^2 = \sum_j (\tilde{\phi}_j^\alpha(x) - \tilde{\phi}_j^\alpha(y))^2$$

Combining the equation above with the defining formula of MDS maps in (1.3), we obtain the desired equality. The independence of the right-hand side of (1.4) on the choice of eigenfunctions ϕ_j of \mathcal{K} is shown in the same way. \square

2.1. Review on Riemannian symmetric spaces. We give a brief review of the basic properties of Riemannian symmetric spaces in this section. We begin by recollecting the basic facts on Riemannian manifolds which will be used in this

paper. The metric g of a connected Riemannian manifold (M^n, g) defines a volume measure μ and a distance function d on M , where

$$d(x, y) = \inf\{\text{length}(\gamma) : \gamma \text{ is a curve from } x \text{ to } y\}.$$

This yields a metric measure space (M, d, μ) . By Hopf-Rinow theorem, (M, d) is a complete metric space if (M, g) is geodesically complete. If (M, g) is complete and connected, then for any $x, y \in M$, there exists a distance minimizing geodesic. For any $x \in M$, the tangential cut locus at x is the set of $v \in T_x M$ such that $\exp(tv)$ is a minimizing geodesic for $t \in [0, 1]$, but fails to be a minimizing geodesic for any $t > 1$. The cut locus \mathcal{C}_x at x is the image of this set under the exponential map at x . If $y \in \mathcal{C}_x$, we have either y is conjugate to x or there is more than 1 distance minimizing geodesic from x to y [15, lemma 8.2]. Therefore, we have $y \in \mathcal{C}_x$ if and only if $x \in \mathcal{C}_y$. Define the symmetric subset of $M \times M$ by

$$\mathcal{C} = \{(x, y) \in M \times M : x \in \mathcal{C}_y\}$$

It is a well-known fact that the function $d^2(x, \cdot)$ is smooth outside \mathcal{C}_x .

By a Laplacian on the Riemannian manifold, we always mean the Laplace-Beltrami operator.

Let (M, g) be a Riemannian manifold. Recall that a local geodesic symmetry at $p \in M$ is a local diffeomorphism r_p on a neighbourhood of p such that for all geodesic $\gamma(t)$ with $\gamma(0) = p$, we have $r_p(\gamma(t)) = \gamma(-t)$.

Definition 2.2. *A Riemannian manifold (M, g) is called a symmetric space if for every $p \in M$, the local geodesic symmetry r_p can be extended to a global isometry on M fixing p .*

It is obvious from the definition that all symmetric spaces are complete, and all connected symmetric spaces are homogeneous, i.e., the isometry group G acts transitively on (M, g) . In fact, every connected symmetric space is a reductive homogeneous space as follows. Denote G and K the isometry group and the isotropy of some $p \in M$, respectively. Let \mathfrak{g} and \mathfrak{k} be the corresponding Lie algebras of G and K . The geodesic symmetry $r_p \in G$ at p satisfies $r_p^2 = Id$. Hence $Ad(r_p)$ is an involutive Lie algebra automorphism of \mathfrak{g} . The Lie algebra \mathfrak{g} admits a decomposition $\mathfrak{g} = \mathfrak{k} + \mathfrak{m}$ such that \mathfrak{k} and \mathfrak{m} are the eigenspaces of $Ad(r_p)$ corresponding to the eigenvalues 1 and -1 , respectively.

We close this section with some geodesic properties of symmetric space which will be used later in the proof. Let $\gamma : \mathbb{R} \rightarrow M$ be a complete geodesic for the symmetric space (M, g) . Then the following family of composed map $\tau_s = r_{\gamma(s)} \circ r_{\gamma(0)}$ is a 1-parameter subgroup of the isometry group G . Easily we have

$$\tau_s(\gamma(t)) = \gamma(t + s).$$

Such maps are called *geodesic transvections* along γ . In terms of the cut locus, although determining the geometry of cut loci of a general Riemannian manifold can be difficult, the cut loci of symmetric spaces have been well studied, see [6, 16, 17]. In particular, for a compact symmetric space (M^n, g) , the cut locus at any $x \in M$ is a finite disjoint union of regular submanifolds with possible different dimensions [17, theorem 3.3]. Taking the union of the submanifolds of dimension $n - 1$ in this decomposition of \mathcal{C}_x , the Riemannian volume density μ together with the perpendicular unit vector fields define a measure $\underline{\mu}_x$ on \mathcal{C}_x . Hence, we may view \mathcal{C}_x as a ‘‘piece-wise smooth’’ manifold of dimension $n - 1$. Note that $\underline{\mu}_x$ can be zero

if the decomposition of \mathcal{C}_x has no components of dimension $n-1$, for example when M is the standard sphere.

3. THE MDS MAP FOR CLOSED CONNECTED SYMMETRIC SPACES

From now on, we focus on the MDS maps of closed connected symmetric spaces. Let (S, g) be a closed connected symmetric space. Denote d and μ the distance function and Borel measure induced by g as before (i.e., the geodesic distance and the Riemannian volume measure). Since S is compact, it is well known that there is an orthonormal basis of $L^2(S)$ contained in $C^\infty(S)$ consisting of Laplacian eigenfunctions, see e.g., [23, p. 2] and [11, theorem 2.2.17]. In addition, each eigenvalue of Δ has finite multiplicity, and eigenspaces of distinct eigenvalues are mutually orthogonal.

Several easy consequences follow from our assumption. Since S is compact, the integral kernel of \mathcal{K} is bounded and uniformly continuous on $S \times S$. Thus by Cauchy-Schwarz inequality, each eigenfunction of \mathcal{K} (also for T) is continuous. In addition, for each non-zero eigenvalue of \mathcal{K} , the corresponding eigenspace is finite-dimensional.

As both the Laplacian operator and the integral kernel $K(x, y)$ are closely related to the distance function, we would like to establish a relation between them. We begin with the following lemma leading to the symmetric property of the integral kernel of \mathcal{K} for symmetric spaces.

Lemma 3.1. *Let $\mu \otimes \mu$ be the product measure on $S \times S$. For a compact symmetric space S , the integral kernel of \mathcal{K} satisfies*

$$(3.1) \quad \Delta_x K(x, y) = \Delta_y K(x, y), \text{ for a.e. } (x, y) \in S \times S$$

Proof. First note that the integral kernel $K(x, y) = -\frac{1}{2}d^2(x, y)$ is always symmetric, i.e. $K(x, y) = K(y, x)$. For a fixed $x \in S$, the function $K(x, \cdot)$ is smooth on $S \setminus \mathcal{C}_x$, which is an open set of full measure in S . Since $x \in \mathcal{C}_y$ if and only if $y \in \mathcal{C}_x$, the functions $\Delta_x K(x, y)$ and $\Delta_y K(x, y)$ are well defined a.e. on $S \times S$ with respect to $\mu \otimes \mu$.

Suppose that $x \notin \mathcal{C}_y$, and let $\gamma : [-1, 1] \rightarrow S$ be the unique minimizing geodesic from x to y . The geodesic symmetry r_0 at the median $\gamma(0)$ is an isometry interchanging x and y . Denote K_y the function $K(\cdot, y)$. Since r_0 is an isometry, in a neighbourhood U of y we have

$$(\Delta K_y)(r_0(z)) = \Delta(K_y \circ r_0)(z), \quad \text{for all } z \in U.$$

Denote Δ_1, Δ_2 the Laplacian with respect to the first and second coordinates in $S \times S$, respectively. We also have

$$\Delta(K_y \circ r_0)(z) = \Delta_1 K(r_0(z), y) = \Delta_1 K(z, r_0(y)) = \Delta_1 K(z, x) = \Delta_2 K(x, z).$$

Taking $z := y$, we get the desired equality. \square

The following proposition shows that the operator T commutes with the extension of the Laplacian operator Δ^D .

Proposition 3.2. *The operator T commutes with Laplacian Δ on closed connected symmetric spaces in the following sense,*

$$(3.2) \quad \langle Tf, \Delta h \rangle = \langle T\Delta f, h \rangle, \quad \text{for all } f, h \in C^\infty(S).$$

Thus T preserves each Laplacian eigenspaces.

Proof. Since the only harmonic functions on closed manifolds are constants, the Laplacian operator Δ commutes with the projection P in (1.1) when acting on smooth functions. As $T = PKP$, it suffices to show \mathcal{K} commutes Δ by the following equality.

$$(3.3) \quad \langle \mathcal{K}f, \Delta h \rangle = \langle \mathcal{K}\Delta f, h \rangle, \quad \text{for all } f, h \in C^\infty(S).$$

Firstly, from Lemma 3.1, we have the following observation on symmetric spaces.

$$\Delta_x K(x, y) = \Delta_y K(x, y), \text{ for a.e. } (x, y) \in S \times S$$

For any $x \in S$, denote \mathcal{C}_x the cut locus of the function $K(x, \cdot)$ as before. We cut the manifold S from \mathcal{C}_x , and obtain a manifold M_x with boundary ∂M_x . Let $i_x : \partial M_x \rightarrow \mathcal{C}_x$ be the canonical projection on the boundary. Using the decomposition theorem of \mathcal{C}_x mentioned in Section 2.1 (see [17] for details), we obtain a decomposition of ∂M_x . Each component $\mathcal{C}_x(i)$ of dimension $n-1$ in \mathcal{C}_x is a regular submanifold, and its preimage under $(i_x)^{-1}$ is a double cover $\partial M_x^0(i) \sqcup \partial M_x^1(i)$. Thus, by restricting only to components of ∂M_x of dimension $n-1$, the unit outer normal N_x defines a measure $\underline{\mu}^x$ on ∂M_x . The space $(\partial M_x, \underline{\mu}^x)$ is isomorphic to $(\mathcal{C}_x, \underline{\mu}_x) \oplus (\mathcal{C}_x, \underline{\mu}_x)$ as measure spaces. For a compact symmetric space (S^n, g) , the conjugate locus at any $x \in S$ has dimension strictly less than $n-1$ [6, theorem 7.3.3]. Since \exp_x is non-singular outside the tangential conjugate locus, $\nabla_y K(x, y)$ is well-defined a.e. on ∂M_x .

For all $f, h \in C^\infty(S)$, we can apply the divergence theorem to obtain

$$\begin{aligned} \langle \Delta^D \mathcal{K}f, h \rangle &= \int_S f(y) \int_S \Delta_x h(x) K(x, y) d\mu(x) d\mu(y) \\ &= \int_S f(y) \int_{\partial M_y} K(x, y) \langle \nabla_x h, N^y(x) \rangle d\underline{\mu}^y(x) d\mu(y) \\ &\quad - \int_S f(y) \int_S \langle \nabla_x h, \nabla_x K(x, y) \rangle d\mu(x) d\mu(y) \end{aligned}$$

Note that for each $y \in S$, the integral

$$\int_{\partial M_y} K(x, y) \langle \nabla_x h, N^y(x) \rangle d\underline{\mu}^y(x)$$

vanishes because h is smooth on S . Therefore, we get

$$(3.4) \quad \langle \Delta^D \mathcal{K}f, h \rangle = - \int_S f(y) \int_S \langle \nabla_x h, \nabla_x K(x, y) \rangle d\mu(x) d\mu(y).$$

Analogously, we obtain

$$(3.5) \quad \langle \mathcal{K}\Delta f, h \rangle = - \int_S h(x) \int_S \langle \nabla_y f, \nabla_y K(x, y) \rangle d\mu(y) d\mu(x)$$

On the other hand, from (3.1) and the divergence theorem, we get

$$\begin{aligned} &\int_S h(x) \left(\int_{\partial M_x} f(y) \langle \nabla_y K, N_y^x \rangle d\underline{\mu}^x(y) - \int_S \langle \nabla_y f, \nabla_y K \rangle d\mu(y) \right) d\mu(x) \\ &= \int_S f(y) \left(\int_{\partial M_y} h(x) \langle \nabla_x K, N_x^y \rangle d\underline{\mu}^y(x) - \int_S \langle \nabla_x h, \nabla_x K \rangle d\mu(x) \right) d\mu(y) \end{aligned}$$

Comparing with the (3.4, 3.5) obtained earlier, we need to prove

$$(3.6) \quad \begin{aligned} & \int_S h(x) \int_{\partial M_x} f(y) \langle \nabla_y K, N_y^x \rangle d\bar{\mu}^x(y) d\mu(x) \\ &= \int_S f(y) \int_{\partial M_y} h(x) \langle \nabla_x K, N_x^y \rangle d\bar{\mu}^y(x) d\mu(y) \end{aligned}$$

For any $y \in \mathcal{C}_x(i)$ not conjugate to x , let $y_j \in M_x^j(i)$ be the preimage of y for $j = 0, 1$. Since distance minimizing geodesics segments starting from x cannot intersect \mathcal{C}_x except for the end-points, there exists distance minimizing unit speed geodesics γ_j in M_x from x to y_j for $j = 0, 1$. For simplicity, denote the geodesic in S corresponding to γ_j by the same notation. The geodesic transvection τ_j along γ_j sending x to y is an isometry. It maps \mathcal{C}_x to \mathcal{C}_y , and the curve γ_j into itself with $\tau_j \circ r_x(y) = x$. Thus, the component of \mathcal{C}_y containing x is a regular submanifold of dimension $n - 1$, and the preimage $i_y^{-1}(x)$ also contains exactly two points in ∂M_y . Let x_j be induced by the reversed curve of γ_j . Because y_j is not conjugate to x and x_j is not conjugate to y , we have $\nabla_y K(x, y_j)$ and $\nabla_x K(y, x_j)$ are well-defined. If we can show

$$(3.7) \quad \sum_{j=0}^1 \langle \nabla_y K(x, y_j), N_{y_j}^x \rangle = \sum_{j=0}^1 \langle \nabla_x K(y, x_j), N_{x_j}^y \rangle,$$

then Equality (3.6) is just an application of the Fubini theorem. Since γ_j is a distance minimizing geodesic segment from x to y_j , we have

$$\nabla_y K(x, y_j) = -d(x, y) \dot{\gamma}_j(d(x, y))$$

Combining the equality above with the fact $\tau_j \circ r_x$ is an isometry mapping y to x , we obtain

$$(\tau_j \circ r_x)_*(\nabla_y K(x, y_j)) = \nabla_x K(x_j, y), \quad (\tau_j \circ r_x)_*(N_{y_j}^x) = N_{x_j}^y, \quad \text{for all } j = 0, 1$$

Then the equation (3.7) holds. Hence, the Laplacian operator commutes with the integral operator T for closed symmetric spaces in the sense of (3.2).

Finally, note that for all $f_1, f_2 \in C^\infty(S)$ such that $\Delta f_i = \lambda_i f_i$ with $i = 1$ and 2 , we have

$$\langle T f_1, \lambda_2 f_2 \rangle = \langle T f, \Delta f_2 \rangle = \langle T \Delta f_1, f_2 \rangle = \langle \lambda_1 T f_1, f_2 \rangle.$$

Therefore, $T f_1$ is perpendicular to all eigenspaces of Δ corresponding to eigenvalues distinct from λ_1 . Since there is an orthonormal basis of $L^2(S)$ consisting of Δ -eigenfunctions, we see $T f_1$ is contained in the eigenspace of λ_1 for Δ . This completes the proof. \square

4. MDS FOR ELEMENTARY SYMMETRIC SPACES

4.1. Products of spheres. We can derive the spectrum of T for product spaces based on the components in the decomposition. This generalizes the discussion of the product of two spaces given in Section 6.2 of [10]. Let (X, d, μ) be the metric measure space induced by the product manifold

$$X = \prod_{i=1}^N X_i, \quad g = \sum_{i=1}^N g_i;$$

where each g_i is the Riemannian metric on a compact connected manifold X_i . Denote d_i and μ_i the Riemannian distance and the normalized Riemannian volume on (X_i, g_i) . It follows that for $x = (x_i) \in X$ and $y = (y_i) \in X$, we have

$$\mu = \prod_{i=1}^N \mu_i, \quad d^2(x, y) = \sum_{i=1}^N d_i^2(x_i, y_i).$$

Since the space (X_i, g_i) are all connected, the Hilbert spaces $L^2(X, \mu)$ and $L^2(X_i, \mu_i)$ are all separable. One would naturally expect the MDS maps and the associated operator to be in the form of Cartesian products of those for each component. This is exactly the case as in [10, section 6.2]. Suppose $\{\phi_i^j\}$ is an orthonormal basis diagonalizing the associated operator T_i for (X_i, d_i, μ_i) . Then $\{\phi_{i_1}^{j_1} \cdots \phi_{i_N}^{j_N}\}$ is an orthonormal basis for $L^2(X, \mu)$ by Fubini theorem. According to [10, proposition 6.2], the associated operator T is diagonalizable with respect to this basis. Since the constant functions are in the kernel of T_i for all $1 \leq i \leq N$, we can choose $\{\phi_i^j\}$ such that the non-constant elements are in 1^\perp . Then for $\Phi = \phi_{i_1}^{j_1} \cdots \phi_{i_N}^{j_N}$, we have $T(\Phi) = 0$ unless the components of Φ contain exactly 1 non-constant function. To see this, we first compute

$$-2K(\Phi)(x) = \sum_{i=1}^N \int_X d_i^2(x_i, y_i) \phi_{i_1}^{j_1}(y_1) \cdots \phi_{i_n}^{j_n}(y_n) d\mu(y)$$

If there is more than one non-constant component in Φ , apply Fubini theorem to the integrals above. We can re-order the integration such that inside each integral, the inner most term satisfies

$$\int_{X_i} \phi_i^{j_i} du_i = 0$$

Therefore, if $N > 1$, the kernel of T is infinite dimensional.

If all components of Φ are constant, clearly $T(\Phi) = PKP(\Phi) = 0$. Assume $\Phi = \phi_1^{j_1}$ is non-constant and $T_1(\phi_1^{j_1}) = \lambda_1^{j_1} \phi_1^{j_1}$. We obtain $T(\Phi) = \lambda_1^{j_1} \Phi$. As a result, for the non-zero spectra of T and T_i , we have

$$(4.1) \quad \text{Spec}(T) \setminus \{0\} = \sqcup_{i=1}^N (\text{Spec}(T_i) \setminus \{0\}),$$

where both sides of the relation above are eigenvalues counted with multiplicity. This implies T is a trace-class operator if and only if each T_i is trace-class.

Since the map T for any compact space has at least 1 positive eigenvalue, the product of N Riemannian manifolds shall have at least N positive eigenvalues counting multiplicity. Summing up, we have the following assertion.

Proposition 4.1. *Let $X = \prod_{i=1}^N X_i$ and $g = \sum_{i=1}^N g_i$ be the product space of compact connected Riemannian manifolds. Denote d and μ the distance and measure induced by g . Then the map T for (X, d, μ) has at least N positive eigenvalues counting multiplicity, and T is a trace-class operator if and only if each T_i is trace-class.*

From now on, let X be a product of spheres. We know from [10, section 6.1] the operator T_i for each component sphere is trace-class and has signature $(1, \infty, \infty)$. Combining (4.1) and the proposition above, we obtain the following statement.

Corollary 4.2. *Let X be a finite product of N spheres (including circles), and define the distance d and measure μ as above. The operator T for (X, d, μ) is trace-class and has signature (∞, ∞, ∞) for $N > 1$.*

As X is closed symmetric, the operator T preserves all eigenspaces of Δ . Let $E_{\lambda_i^\alpha}$ be an eigenspace for the Laplace operator of (X_i, g_i) and denote $\pi_i : X \rightarrow X_i$ the canonical projection, then T preserves the space $\pi_i^*(E_{\lambda_i^\alpha})$ and is self-adjoint on it with respect to the $L^2(X, \mu)$ -norm. In fact, the operator T is simply a constant scaling on $\pi_i^*(E_{\lambda_i^\alpha})$. To show this, denote G the isometry group of g , then for any $s \in G$ we have

$$\begin{aligned} \mathcal{K}(s \cdot f)(x) &= \int_X K(x, y) f(s \cdot y) d\mu(y) \\ &= \int_X K(s \cdot x, s \cdot y) f(s \cdot y) d\mu(y) \\ &= \int_X K(s \cdot x, z) f(z) d\mu(z) = \mathcal{K}(f)(s \cdot x). \end{aligned}$$

Therefore, the natural action of G commutes with the operator \mathcal{K} (hence T). Let G_i be the isometry group of (X_i, g_i) . Since each component (X_i, g_i) is a sphere, the isometry group G_i acts irreducibly on $E_{\lambda_i^\alpha}$ [7, theorem 3.1], and preserves the eigenspaces of T when viewed as a subgroup of G . Thus, T can have only 1-real eigenvalue on $\pi_i^*(E_{\lambda_i^\alpha})$. In particular, the map T has only 1 real eigenvalue on each Laplacian eigenspace of standard spheres.

4.2. Projective spaces.

4.2.1. *Signature of T for projective spaces.* Here we consider the case $X := \mathbb{RP}^n$, an n -dimensional real projective space equipped with its geodesic distance and Riemannian volume measure. Since $\mathbb{RP}^n = \mathbb{S}^n / \mathbb{Z}_2$, we expect the MDS for projective spaces behaves similarly compared to spheres. On the other hand, we will see in this section how the global topology makes a difference in MDS maps on projective spaces compared to the MDS on spheres. We start by determining the signature of T for projective spaces \mathbb{RP}^n .

The Laplacian eigenfunctions on \mathbb{RP}^n are well-defined projections of spherical harmonics. Hence, they are projections of spherical harmonics of even degree. Clearly as for the spheres, the group $SO(n)$ acts irreducibly on each Laplacian eigenspace of \mathbb{RP}^n . Thus, each eigenspace of T is a direct sum of eigenspace of Δ on \mathbb{RP}^n .

For $x, y \in \mathbb{S}^n$, the distance between the lines $[x]$ and $[y]$ on \mathbb{RP}^n is $\arccos(|\langle x, y \rangle|)$. We know that the operators T and \mathcal{K} share all eigenvalues and eigenfunctions except for those corresponding to constant functions. To compute the spectrum of \mathcal{K} for projective spaces, let $2k$ be even. Then by Funke-Hecke theorem [4, p. 98], the eigenvalue of \mathcal{K} for \mathbb{RP}^n corresponding to spherical harmonics of degree $2k$ is given by

$$(4.2) \quad \lambda_{2k}^n = \sigma_n \int_0^1 \arccos^2(t) P_{2k}^n(t) (1-t^2)^{(n-2)/2} dt,$$

where $P_{2k}^n(t)$ is the Legendre polynomials for \mathbb{S}^n of degree $2k$, and the numbers $\sigma_n = -\frac{\text{vol}(\mathbb{S}^{n-1})}{\text{vol}(\mathbb{S}^n)}$ are negative constants depending only on n . Inside the integral in (4.2), only the term

$$(4.3) \quad F_{2k}^n(t) = P_{2k}^n(t) (1-t^2)^{(n-2)/2} = R_{2k}^n \left(\frac{d}{dt} \right)^{2k} (1-t^2)^{2k+(n-2)/2}$$

is affected by the dimension n and even degree $2k$. For $2k$ even, the Rodrigues constants R_{2k}^n are given by [13, p. 22]

$$(4.4) \quad R_{2k}^n = \frac{1}{4^k} \frac{\Gamma(n/2)}{\Gamma(n/2 + 2k)}.$$

These constants are all positive. Note the equations (4.2,4.3) are still valid for the special case $n = 1$, where P_{2k}^1 are given by Chebyshev polynomials of the first kind.

For $n \geq 1$ and $2k \geq 2$, a substitution $t = \cos \theta$ using $\frac{d}{d\theta} = -\sin \theta \cdot \frac{d}{dt}$ yields

$$\begin{aligned} & F_{2k}^n(\cos \theta) \\ &= R_{2k}^n \left(\frac{-1}{\sin \theta} \frac{d}{d\theta} \right)^{2k} [(\sin \theta)^{4k+n-2}] \\ &= (2k+n-2) R_{2k}^n \left(\frac{-1}{\sin \theta} \frac{d}{d\theta} \right)^{2k-2} \left[(2(2k-1)+n-2)(\sin \theta)^{2(2k-2)+n-2} \right. \\ & \quad \left. - (2(2k-1)+s+1)(\sin \theta)^{2(2k-2)+n} \right] \\ &= (4k+n-2) \left[(2(2k-1)+n-2) \frac{R_{2k}^n}{R_{2k-2}^n} F_{2k-2}^n(\cos \theta) \right. \\ & \quad \left. - (2(2k-1)+n-1) \frac{R_{2k}^n}{R_{2k-2}^{n+2}} F_{2k-2}^{n+2}(\cos \theta) \right] \end{aligned}$$


By linearity of the integration, we combine the equation above and (4.2) to obtain

$$(4.5) \quad \frac{1}{\sigma_n} \lambda_{2k}^n = (4k+n-2) \left((4k+n-4) \frac{R_{2k}^n}{\sigma_n R_{2k-2}^n} \lambda_{2k-2}^n - (4k+n-3) \frac{R_{2k}^n}{\sigma_{n+2} R_{2k-2}^{n+2}} \lambda_{2k-2}^{n+2} \right)$$

Note for $n \geq 1$ and $2k \geq 2$, we always have $4k+n-4 \geq 1 > 0$. From the signs of the coefficients in the equation above, we see the following: If $\{\lambda_{2k}^n\}_{k=1}^\infty$ is a sequence of alternating signs indexed by even numbers $2k$, the signs of numbers in the sequence $\{\lambda_{2k}^{n+2}\}$ will also be alternating. For the basic case $n = 1$, the Riemannian manifold \mathbb{RP}^1 is isomorphic to \mathbb{S}^1 by doubling the angles between lines. For \mathbb{S}^1 , the signs of eigenvalues of \mathcal{K} corresponding to the eigenfunctions $\cos(m\theta)$ are alternating, depending on whether m is odd or even [2, section 2]. Thus, the signs of the sequence $\{\lambda_{2k}^1\}$ indexed by even $2k$ are also alternating. Combined with the argument above, we obtain the following result.

Proposition 4.3. *For odd number n , the MDS associated operator T on \mathbb{RP}^n has both infinitely many positive and negative eigenvalues.*

We then turn to the case of even-dimensional projective spaces, starting from the basic case of \mathbb{RP}^2 . We expect that the signature of T for even-dimensional projective spaces is similar to the odd dimension cases. A numeric computation for the first 6 eigenvalues (corresponding to even degree spherical harmonics up to $2k = 10$). From the pictures below, we can see that the first 6 eigenvalues have alternating signs.



Although computing the spectrum of T for projective spaces can be difficult, we can easily obtain some information on $\text{sgn}(T)$ for \mathbb{RP}^2 using Mercer's theorem.

Lemma 4.4. *The operator T on \mathbb{RP}^2 has infinitely many strictly positive and negative eigenvalues counting multiplicity.*

Proof. Since \mathbb{RP}^2 is homogeneous, we only need to prove this statement for \mathcal{K} .

Suppose \mathcal{K} has only finitely many strictly negative eigenvalues counting multiplicity. Let λ_{2k} be the eigenvalue of \mathcal{K} corresponding to the degree $2k$ spherical harmonics as before. Denote $\{\phi_{2k}^m\}$ with $-2k \leq m \leq 2k$ the standard L_2 -orthonormal basis with respect to μ of Laplacian eigenspace \mathcal{H}_{2k} of \mathbb{RP}^2 corresponding to degree $2k$ spherical harmonics. The function

$$(4.6) \quad N(x, y) = \sum_{\lambda_{2k} < 0} \lambda_{2k} \sum_{m=-2k}^{2k} \phi_{2k}^m(x) \phi_{2k}^m(y)$$

is smooth and bounded, since the summation above is finite. We can decompose the integral kernel as

$$K(x, y) = N(x, y) + H(x, y),$$

where $N(x, y)$ is a negative integral kernel and $H(x, y)$ is a positive integral kernel. Because the positive operator

$$H(f)(x) = \int_{\mathbb{RP}^2} f(y) d\mu(y)$$

has a continuous bounded kernel, Mercer's theorem [3, theorem 4.10] implies the following convergence with respect to summation of $\lambda_{2k} \phi_{2k}^m(x) \phi_{2k}^m(y)$ is absolute

and uniform:

$$(4.7) \quad H(x, y) = \sum_{\lambda_{2k} \geq 0} \sum_{m=-2k}^{2k} \lambda_{2k} \phi_{2k}^m(x) \phi_{2k}^m(y).$$

From the basics properties of spherical harmonics [4, theorem 3.3.3], we obtain

$$\begin{aligned} \sum_{m=-2k}^{2k} \phi_{2k}^m(x) \phi_{2k}^m(y) &= \dim(\mathcal{H}_k) P_{2k}(\cos(d(x, y))) \\ &= (4k + 1) P_{2k}(x \cdot y). \end{aligned}$$

Here we use x, y to denote both points in $\mathbb{R}\mathbb{P}^2$ and their lifts to \mathbb{S}^2 . Since P_{2k} is even, the value of $P_{2k}(x \cdot y)$ is always well-defined.

Therefore, we obtain the following absolute and uniform convergence

$$H(x, y) = \sum_{\lambda_{2k} \geq 0} \lambda_{2k} (4k + 1) P_{2k}(x \cdot y)$$

It follows that the following convergence of functions on $(\mathbb{R}\mathbb{P}^2)^2$

$$(4.8) \quad -\frac{1}{2} d^2(x, y) = -\frac{1}{2} \arccos^2(|x \cdot y|) = K(x, y) = \sum_{k=0}^{\infty} \lambda_{2k} (4k + 1) P_{2k}(x \cdot y)$$

is absolute and uniform. This implies for $t \in [0, 1]$, the convergence

$$(4.9) \quad \arccos^2(t) = -2 \sum_{k=0}^{\infty} \lambda_{2k} (4k + 1) P_{2k}(t)$$

is also absolute and uniform.

On the other hand, we know from (4.2), the eigenvalues of \mathcal{K} for $\mathbb{R}\mathbb{P}^1$ are given by

$$(4.10) \quad \lambda_{2j}^1 = -\frac{1}{\pi} \int_0^1 \arccos^2(t) C_{2j}(t) \frac{1}{\sqrt{1-t^2}} dt,$$

where C_{2j} are Chebyshev polynomials of the first kind of degree $2j$. Since the convergence in (4.9) is absolute and uniform, dominated convergence theorem implies

$$(4.11) \quad \lambda_{2j}^1 = \frac{2}{\pi} \sum_{k=0}^{\infty} \lambda_{2k} (4k + 1) \int_0^1 P_{2k}(t) C_{2j}(t) \frac{1}{\sqrt{1-t^2}} dt$$

According to [8, p. 96], the even degree Legendre polynomials can be expanded by Chebyshev polynomials as

$$P_{2k}(t) = \left(\frac{\Gamma(1/2 + k)}{\Gamma(1/2)\Gamma(k + 1)} \right)^2 C_0 + \frac{2}{\Gamma^2(1/2)} \sum_{i=1}^k \left(\frac{\Gamma(k - i + 1/2)\Gamma(k + i + 1/2)}{\Gamma(k - i + 1)\Gamma(k + i + 1)} \right) C_{2i}(t)$$

The functions $C_{2j}(t)$ are even, and we have

$$\int_0^1 C_{2i}(t) C_{2j}(t) \frac{1}{\sqrt{1-t^2}} dt = \frac{\pi}{4} (\delta_j^i + \delta_i^0 \delta_j^0)$$

Therefore, we obtain

$$(4.12) \quad \int_0^1 P_{2k} C_{2j} \frac{1}{\sqrt{1-t^2}} dt = 0, \quad 0 \leq k < j$$

$$(4.13) \quad \int_0^1 P_{2k} C_{2j} \frac{1}{\sqrt{1-t^2}} dt > 0, \quad k \geq j$$

These equalities and (4.11) imply λ_{2j}^1 is always non-negative for sufficiently large j . This leads to a contradiction. Hence, the operator \mathcal{K} (hence T) for $\mathbb{R}\mathbb{P}^2$ has to admit infinitely many negative eigenvalues counting multiplicity.

Similarly, we can prove that T on $\mathbb{R}\mathbb{P}^2$ also has infinitely many positive eigenvalues counting multiplicity. This completes the proof. \square

Starting from $\mathbb{R}\mathbb{P}^2$, an induction argument on the dimension of the projective space using (4.5) shows T for every even dimension projective space has to admit both infinitely many positive and negative eigenvalues counting multiplicity. Combined with the result of odd-dimensional projective spaces, we obtain the following.

Theorem 4.5. *For any projective space $\mathbb{R}\mathbb{P}^n$, the operator T always has both infinitely many positive and negative eigenvalues counting multiplicity.*

4.2.2. *Spectral asymptotics of T odd-dimensional projective spaces.* Now we estimate the norm of the eigenvalues of T for odd-dimensional projective spaces. This will show how the MDS maps of $\mathbb{R}\mathbb{P}^n$ differ from the MDS maps of \mathbb{S}^n for odd n . To estimate the asymptotics of the eigenvalues $\{\lambda_{2k}^n\}$, we need the following lemma.

Lemma 4.6. *For odd-dimensional projective space $\mathbb{R}\mathbb{P}^n$, we have*

$$\lambda_{2k}^n = \Theta(k^{-(n+3)/2}).$$

Proof. We prove this statement by induction on n . For the base case $n = 1$, a simple integration by parts shows

$$\lambda_{2k}^1 = \Theta(k^{-2}).$$

Suppose $\lambda_{2k}^n = \Theta(k^{-(n+3)/2})$ holds. All coefficients for the eigenvalues in (4.5) are non-zero. Solving this equation for λ_{2k-2}^n and we get

$$\begin{aligned} |\lambda_{2k-2}^{n+2}| &\leq C_1(n, k) \left(\frac{R_{2k-2}^{n+2}}{R_{2k}^n} \right) |\lambda_{2k}^n| + C_2(n, k) \left(\frac{R_{2k-2}^{n+2}}{R_{2k-2}^n} \right) |\lambda_{2k-2}^n| \\ |\lambda_{2k-2}^{n+2}| C_2(n, k) &\geq C_2(n, k) \left(\frac{R_{2k-2}^{n+2}}{R_{2k-2}^n} \right) |\lambda_{2k-2}^n|, \end{aligned}$$

where the second inequality comes from the fact that the sequence $\{\lambda_{2k}^n\}$ indexed by $2k$ has alternating signs. Here $C_1(n, k)$ and $C_2(n, k)$ are positive functions such that $C_1(n, k) = \Theta(k^{-2})$ and $C_2(n, k) = \Theta(1)$ when n is fixed. Moreover, from the formula of Rodrigues constants, we get

$$(4.14) \quad 0 < \frac{R_{2k-2}^{n+2}}{R_{2k}^n} = \Theta(k) \text{ and } 0 < \frac{R_{2k-2}^{n+2}}{R_{2k-2}^n} = \Theta(k^{-1}).$$

Then these inequalities together imply $\lambda_{2k}^{n+2} = \Theta(k^{-1-(n+3)/2})$. \square

Recall that by Lemma 2.1, for any $x, y \in X$, the value of $\|\mathcal{M}(x) - \mathcal{M}(y)\|_2$ is independent of the choice of the eigenfunctions of \mathcal{K} defining \mathcal{M} . We are able to state the following result regarding the MDS maps for odd-dimensional projective spaces.

Theorem 4.7. *For every odd $n > 1$, the operator T for \mathbb{RP}^n is not trace-class. Furthermore, given any $x \in \mathbb{RP}^n$, there exists a positive measure set U_x such that*

$$(4.15) \quad \|\mathcal{M}(x) - \mathcal{M}(y)\|_2 = +\infty, \quad \text{for all } y \in U_x,$$

and the distance reconstruction formula (1.4) does not hold $\mu \otimes \mu$ almost everywhere.

Proof. According to [4, theorem 3.1.4], the dimension of the space of degree $2k$ spherical harmonics on \mathbb{RP}^n is given by

$$(4.16) \quad \dim(\mathcal{H}_{2k}^n) = \frac{4k + n - 1}{2k + n - 1} \binom{2k + n - 1}{n - 1}.$$

Using the relation of binomial coefficients, we easily deduce that $\dim(\mathcal{H}_{2k}^n)$ is a polynomial in k of degree $n - 1$. Then for all odd n , we have

$$(4.17) \quad |\lambda_{2k}^n| \dim(\mathcal{H}_{2k}^n) = \Theta(k^{(n-5)/2})$$

Therefore, for all odd n with $n > 1$, the operator T on \mathbb{RP}^n is not a trace-class operator.

First we show $\|\mathcal{M}(x) - \mathcal{M}(y)\|_2 = +\infty$ when $x \cdot y = 0$. Let $\{\phi_{2k}^i\}$ with $i \in I_{2k}^n$ be an orthonormal basis with respect to μ of \mathcal{H}_{2k}^n . For $x, y \in \mathbb{RP}^n$, we have

$$(4.18) \quad \begin{aligned} \|\mathcal{M}(x) - \mathcal{M}(y)\|_2^2 &= \sum_{\lambda_{2k} > 0} \lambda_{2k} \sum_{i \in I_{2k}^n} (\phi_{2k}^i(x) - \phi_{2k}^i(y))^2 \\ &= \sum_{k=0}^{\infty} \lambda_{4k+2} \sum_{i \in I_{4k+2}^n} (\phi_{4k+2}^i(x) - \phi_{4k+2}^i(y))^2 \\ &= \sum_{k=0}^{\infty} \lambda_{4k+2} \cdot \dim(\mathcal{H}_{4k+2}^n) \cdot 2(1 - P_{4k+2}^n(x \cdot y)) \end{aligned}$$

The Legendre polynomials have the following recurrence relation (see [4, proposition 3.3.11])

$$(4.19) \quad (k + n - 1)P_{k+1}^n(t) - (2k + n - 1)tP_k^n(t) + kP_{k-1}^n(t) = 0.$$

Therefore, for fixed n we can see $\{|P_{4k+2}^n(0)|\}$ for $k \geq 0$ is a decreasing sequence uniformly bounded away from 1. Since $x \cdot y = 0$, for any $n > 1$, there exists some $\delta_n > 0$ such that

$$\|\mathcal{M}(x) - \mathcal{M}(y)\|_2^2 \geq 2\delta_n \sum_{k=0}^{\infty} \lambda_{4k+2} \cdot \dim(\mathcal{H}_{4k+2}^n).$$

Using (4.17), we can see $\|\mathcal{M}(x) - \mathcal{M}(y)\|_2^2 = +\infty$ for odd $n > 1$.

From (4.18), we see that $\|\mathcal{M}(x) - \mathcal{M}(y)\|_2$ depends only on $|x \cdot y|$. Moreover, if $x_0 \cdot x_1 = 0$, the triangle inequality implies there is a set of positive measure U_{x_i} such that $\|\mathcal{M}(y) - \mathcal{M}(x_i)\| = +\infty$ for all $y \in U_{x_i}$ and for some $i = 0$ or 1 . The operator T commutes with the action of the isometry group G , so the distance $\|\mathcal{M}(x) - \mathcal{M}(y)\|_2$ is invariant under the action by G . Since G acts transitively on \mathbb{RP}^n , it follows that for every $x \in \mathbb{RP}^n$ with odd $n > 1$, there exists a positive measure set U_x such that (4.15) holds.

Finally, by Fubini theorem one has then that the series on the left-hand side of the distance reconstruction formula (1.4) does not converge on a set of couples (x, y) of positive $\mu \otimes \mu$ measure, and hence (1.4) does not hold $\mu \otimes \mu$ almost everywhere, which concludes the proof. \square

REFERENCES

- [1] H. Adams, M. Blumstein, and L. Kassab. Multidimensional scaling on metric measure spaces. *Rocky Mountain Journal of Mathematics*, 50(2):397–413, 2020.
- [2] D. Azevedo and V.A. Menegatto. Eigenvalues of dot-product kernels on the sphere. *Proceeding Series of the Brazilian Society of Computational and Applied Mathematics*, 3(1), 2015.
- [3] Felipe Cucker and Ding Xuan Zhou. *Learning theory: an approximation theory viewpoint*, volume 24. Cambridge University Press, 2007.
- [4] Helmut Groemer. *Geometric applications of Fourier series and spherical harmonics*, volume 61. Cambridge University Press, 1996.
- [5] J. Heinonen, P. Koskela, N. Shanmugalingam, and J.T. Tyson. *Sobolev spaces on metric measure spaces*. Number 27. cambridge university press, 2015.
- [6] Sigurdur Helgason. *Differential geometry and symmetric spaces*, volume 341. American Mathematical Soc., 2001.
- [7] Sigurdur Helgason. *Groups and geometric analysis: integral geometry, invariant differential operators, and spherical functions*, volume 83. American Mathematical Society, 2022.
- [8] Mourad Ismail. *Classical and quantum orthogonal polynomials in one variable*, volume 13. Cambridge university press, 2005.
- [9] Lara Kassab. Multidimensional scaling: Infinite metric measure spaces. *arXiv preprint arXiv:1904.07763*, 2019. Masters thesis, Colorado State University.
- [10] Alexey Kroshnin, Eugene Stepanov, and Dario Trevisan. Infinite multidimensional scaling for metric measure spaces. *ESAIM: Control, Optimisation and Calculus of Variations*, 28:58, 2022.
- [11] Michael Levitin, Dan Mangoubi, and Iosif Polterovich. *Topics in spectral geometry*, volume 237. American Mathematical Society, 2023.
- [12] Sunhyuk Lim and Facundo Memoli. Classical multidimensional scaling on metric measure spaces. *arXiv preprint arXiv:2201.09385*, 2022.
- [13] Claus Müller. *Analysis of spherical symmetries in Euclidean spaces*, volume 129. Springer Science & Business Media, 2012.
- [14] Nikita P., Vladimir S., Eugene S., and Dario T. Reconstruction of manifold embeddings into euclidean spaces via intrinsic distances, 2020. arxiv 2012.13770.
- [15] Peter Petersen. *Riemannian geometry (1st ed.)*, volume 171. Springer-Verlag, 1998.
- [16] Takashi Sakai. On the structure of cut loci in compact riemannian symmetric spaces. *Mathematische Annalen*, 235:129–148, 1978.
- [17] Masaru Takeuchi. On conjugate loci and cut loci of compact symmetric spaces i. *Tsukuba journal of mathematics*, 2:35–68, 1978.
- [18] J.T. Tyson and J.-M. Wu. Characterizations of snowflake metric spaces. *Ann. Acad. Sci. Fenn. Math*, 30(2):313–336, 2005.
- [19] A Vershik and F Petrov. Limit spectral measures of matrix distributions of metric triples. *Functional Analysis and Its Applications*, 57:169–172, 2023.
- [20] J. Von Neumann and I.J. Schoenberg. Fourier integrals and metric geometry. *Transactions of the American Mathematical Society*, 50(2):226–251, 1941.
- [21] J. Wang. *Geometric structure of high-dimensional data and dimensionality reduction*. Springer, 2012.
- [22] W.A. Wilson. On certain types of continuous transformations of metric spaces. *American Journal of Mathematics*, 57(1):62–68, 1935.
- [23] Steve Zelditch. *Eigenfunctions of the Laplacian on a Riemannian manifold*, volume 125. American Mathematical Soc., 2017.

EIGENVALUES AND EIGENFUNCTIONS OF MDS DEFINING OPERATORS ON SYMMETRIC SPACES

HSE UNIVERSITY, MOSCOW, RUSSIAN FEDERATION

Email address: tma@hse.ru

ST.PETERSBURG BRANCH OF THE STEKLOV MATHEMATICAL INSTITUTE OF THE RUSSIAN ACADEMY OF SCIENCES, ST.PETERSBURG, RUSSIAN FEDERATION AND DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DI PISA, LARGO BRUNO PONTECORVO 5, 56127 PISA, ITALY AND HSE UNIVERSITY, MOSCOW, RUSSIAN FEDERATION

Email address: stepanov.eugene@gmail.com