

DIVIDED DIFFERENCE OPERATORS ON POLYTOPES

VALENTINA KIRITCHENKO

ABSTRACT. We define convex-geometric counterparts of divided difference (or Demazure) operators from the Schubert calculus and representation theory. These operators are used to construct inductively polytopes that capture Demazure characters of representations of reductive groups. In particular, Gelfand–Zetlin polytopes and twisted cubes of Grossberg–Karshon are obtained in a uniform way.

1. INTRODUCTION

Polytopes play a prominent role in representation theory and algebraic geometry. In algebraic geometry, there are Okounkov convex bodies introduced by Kaveh–Khovanskii and Lazarsfeld–Mustata (see [KKh] for the references). These convex bodies turn out to be polytopes in many important cases (e.g. for spherical varieties). In representation theory, there are string polytopes introduced by Berenstein–Zelevinsky and Littelmann [BZ, L]. String polytopes are associated with the irreducible representations of a reductive group G , namely, the integer points inside and at the boundary of a string polytope parameterize a canonical basis in the corresponding representation. A classical example of a string polytope for $G = GL_n$ is a Gelfand–Zetlin polytope.

There is a close relationship between string polytopes and Okounkov bodies. String polytopes were identified with Okounkov polytopes of flag varieties for a geometric valuation [K] and were also used in [KKh] to give a more explicit description of Okounkov bodies associated with actions of G on algebraic varieties. Natural generalizations of string polytopes are Okounkov polytopes of Bott–Samelson resolutions of Schubert varieties for various geometric valuations (an example of such a polytope is computed in [Anderson]).

In this paper, we introduce an elementary convex-geometric construction that yields polytopes with the same properties as string polytopes and Okounkov polytopes of Bott–Samelson resolutions. Namely, exponential sums over the integer points inside these polytopes coincide with Demazure characters. We start from a single point and apply a sequence of simple convex-geometric operators that mimic

Key words and phrases. Gelfand–Zetlin polytope, divided difference operator, Demazure character.

The author was supported by Dynasty foundation, AG Laboratory NRU HSE, MESRF grants ag. 11.G34.31.0023, MK-983.2013.1 and by RFBR grants 12-01-31429-mol-a, 12-01-33101-mol-a-ved. This study was carried out within “The National Research University Higher School of Economics’ Academic Fund Program in 2013-2014, research grant No. 12-01-0194”.

the well-known *divided difference* or *Demazure operators* from the Schubert calculus and representation theory. Convex-geometric Demazure operators act on convex polytopes and take a polytope to a polytope of dimension one greater. In particular, classical Gelfand–Zetlin polytopes can be obtained in this way (see Section 3.2). More generally, these operators act on *convex chains*. The latter were defined and studied in [PKh] and used in [PKh2] to prove a convex-geometric variant of the Riemann–Roch theorem.

When $G = GL_n$, convex-geometric Demazure operators were implicitly used in [KST] to calculate Demazure characters of Schubert varieties in terms of the exponential sums over unions of faces of Gelfand–Zetlin polytopes and to represent Schubert cycles by unions of faces. A motivation for the present paper is to create a general framework for extending results of [KST] on Schubert calculus from type A to arbitrary reductive groups. In particular, convex-geometric divided difference operators allow one to use in all types a geometric version of mitosis (mitosis on parallelepipeds) developed in [KST, Section 6]. This might help to find an analog of mitosis of [KnM] in other types.

Another motivation is to give a tool for describing inductively Okounkov polytopes of Bott–Samelson resolutions. We describe polytopes that conjecturally coincide with Okounkov polytopes of Bott–Samelson resolutions for a natural choice of a geometric valuation (see Conjecture 4.1). Another application is an inductive description of Newton–Okounkov polytopes for line bundles on Bott towers (in particular, on toric degenerations of Bott–Samelson resolutions) that were first described by Grossberg and Karshon [GK] (see Section 4.1 and Remark 4.6).

This paper is organized as follows. In Section 2, we give background on convex chains and define convex-geometric divided difference operators. In Section 3, we relate these operators with Demazure characters and their generalizations. In Section 4, we outline possible applications to Okounkov polytopes of Bott towers and Bott–Samelson varieties.

I am grateful to Dave Anderson, Joel Kamnitzer, Kiumars Kaveh and Askold Khovanskii for useful discussions. I would also like to thank the referee and Megumi Harada for the careful reading and valuable comments.

2. MAIN CONSTRUCTION

2.1. String spaces and parapolytopes.

Definition 1. A *string space* of rank r is a real vector space \mathbb{R}^d together with a direct sum decomposition

$$\mathbb{R}^d = \mathbb{R}^{d_1} \oplus \dots \oplus \mathbb{R}^{d_r}$$

and a collection of linear functions $l_1, \dots, l_r \in (\mathbb{R}^d)^*$ such that l_i vanishes on \mathbb{R}^{d_i} .

We choose coordinates in \mathbb{R}^d such that they are compatible with the direct sum decomposition. The coordinates will be denoted by $(x_1^1, \dots, x_{d_1}^1; \dots; x_1^r, \dots, x_{d_r}^r)$ so

that the summand \mathbb{R}^{d_i} is given by vanishing of all coordinates except for $x_1^i, \dots, x_{d_i}^i$. In what follows, we regard \mathbb{R}^d as an affine space.

Let $\mu = (\mu_1, \dots, \mu_{d_i})$ and $\nu = (\nu_1, \dots, \nu_{d_i})$ be two collections of real numbers such that $\mu_j \leq \nu_j$ for all $j = 1, \dots, d_i$. By the *coordinate parallelepiped* $\Pi(\mu, \nu) \subset \mathbb{R}^{d_i}$ we mean the parallelepiped

$$\Pi(\mu, \nu) = \{(x_1^i, \dots, x_{d_i}^i) \in \mathbb{R}^{d_i} \mid \mu_j \leq x_j^i \leq \nu_j, j = 1, \dots, d_i\}.$$

Definition 2. A convex polytope $P \subset \mathbb{R}^d$ is called a *parapolytope* if for $i = 1, \dots, r$, and any vector $c \in \mathbb{R}^d$ the intersection of P with the parallel translate $c + \mathbb{R}^{d_i}$ of \mathbb{R}^{d_i} is either empty or the parallel translate of a coordinate parallelepiped, i.e.,

$$P \cap (c + \mathbb{R}^{d_i}) = c + \Pi(\mu_c, \nu_c)$$

for μ_c and ν_c that depend on c .

For instance, if $d = r$ (i.e., $d_1 = \dots = d_r = 1$) then every polytope is a parapolytope. Below is a less trivial example of a parapolytope in a string space.

Example 2.1. Consider the string space

$$\mathbb{R}^d = \mathbb{R}^{n-1} \oplus \mathbb{R}^{n-2} \oplus \dots \oplus \mathbb{R}^1$$

of rank $r = (n - 1)$ and dimension $d = \frac{n(n-1)}{2}$.

Let $\lambda = (\lambda_1, \dots, \lambda_n)$ be a non-increasing collection of integers. For each λ , define the *Gelfand–Zetlin polytope* Q_λ by the inequalities

$$\begin{array}{cccccccc} \lambda_1 & & \lambda_2 & & \lambda_3 & & \dots & & \lambda_n \\ & x_1^1 & & x_2^1 & & \dots & & & x_{n-1}^1 \\ & & x_1^2 & & \dots & & & x_{n-2}^2 & \\ & & & \ddots & & \ddots & & & \\ & & & & x_1^{n-2} & & x_2^{n-2} & & \\ & & & & & x_1^{n-1} & & & \end{array}$$

where the notation

$$\begin{array}{cc} a & b \\ & c \end{array}$$

means $a \geq c \geq b$. It is easy to check that Q_λ is a parapolytope. Indeed, consider the parallel translate of \mathbb{R}^{n-i} by a vector $c = (c_1^1, \dots, c_{n-1}^1; \dots; c_1^{n-1})$. Put $c_i^0 = \lambda_i$ for $i = 1, \dots, n$. The intersection of Q_λ with $c + \mathbb{R}^{n-i}$ is given by the the following inequalities:

$$\begin{array}{cccccccc} c_1^{i-1} & & c_2^{i-1} & & c_3^{i-1} & & \dots & & c_{n-i+1}^{i-1} \\ & x_1^i & & x_2^i & & \dots & & & x_{n-i}^i \\ & & c_1^{i+1} & & \dots & & c_{n-i-1}^{i+1} & & \end{array}.$$

Therefore, the intersection can be identified with the coordinate parallelepiped $c + \Pi(\mu, \nu) \subset c + \mathbb{R}^{n-i}$, where $\mu_j = \max(c_j^{i-1}, c_{j-1}^{i+1})$ and $\nu_j = \min(c_{j+1}^{i-1}, c_j^{i+1})$ (put $c_0^{i+1} = -\infty$ and $c_{n-i}^{i+1} = +\infty$).

2.2. Polytopes and convex chains. Consider the set of all convex polytopes in \mathbb{R}^d . This set can be endowed with the structure of a commutative semigroup using *Minkowski sum*

$$P_1 + P_2 = \{x_1 + x_2 \in \mathbb{R}^d \mid x_1 \in P_1, x_2 \in P_2\}$$

It is not hard to check that this semigroup has cancelation property. We can also multiply polytopes by positive real numbers using dilation:

$$\lambda P = \{\lambda x \mid x \in P\}, \quad \lambda \geq 0.$$

Hence, we can embed the semigroup S of convex polytopes into its Grothendieck group V , which is a real (infinite-dimensional) vector space. The elements of V are called *virtual polytopes*.

It is easy to check that the set of parapolytopes in \mathbb{R}^d is closed under Minkowski sum and under dilations. Hence, we can define the subspace $V_{\square} \subset V$ of *virtual parapolytopes* in the string space \mathbb{R}^d .

Example 2.2. If \mathbb{R}^d is a string space of rank 1, i.e. $d_1 = d$, then parapolytopes are coordinate parallelepipeds $\Pi(\mu, \nu)$. Clearly,

$$\Pi(\mu, \nu) + \Pi(\mu', \nu') = \Pi(\mu + \mu', \nu + \nu').$$

Hence, virtual parapolytopes can be identified with the pairs of vectors $\mu, \nu \in \mathbb{R}^d$. This yields an isomorphism $V_{\square} \simeq \mathbb{R}^d \oplus \mathbb{R}^d$. Under this isomorphism, the semigroup of (true) coordinate parallelepipeds gets mapped to the convex cone in $\mathbb{R}^d \oplus \mathbb{R}^d$ given by the inequalities $\mu_i \leq \nu_i$ for $i = 1, \dots, d$.

We now define the space \tilde{V} of *convex chains* following [PKh]. A *convex chain* is a function on \mathbb{R}^d that can be represented as a finite linear combination

$$\sum_P c_P \mathbb{I}_P,$$

where $c_P \in \mathbb{R}$, and \mathbb{I}_P is the characteristic function of a convex polytope $P \subset \mathbb{R}^d$, that is,

$$\mathbb{I}_P(x) = \begin{cases} 1, & x \in P \\ 0, & x \notin P \end{cases}.$$

The semigroup S of convex polytopes can be naturally embedded into \tilde{V} :

$$\iota : S \hookrightarrow \tilde{V}; \quad \iota : P \mapsto \mathbb{I}_P$$

In what follows, we will work in the space of convex chains and freely identify a polytope P with the corresponding convex chain \mathbb{I}_P . However, note that the embedding ι is not a homomorphism, that is, $\mathbb{I}_{P+Q} \neq \mathbb{I}_P + \mathbb{I}_Q$ (the sum of convex chains is defined as the usual sum of functions).

Remark 2.3. The embedding $\iota : S \hookrightarrow \tilde{V}$ can be extended to the space V of all virtual polytopes. Namely, there exists a commutative operation $*$ on \tilde{V} (called *product of convex chains*) such that

$$\mathbb{I}_{P+Q} = \mathbb{I}_P * \mathbb{I}_Q \quad (M)$$

for any two convex polytopes P and Q (see [PKh, Section 2, Proposition-Definition 3]). Virtual polytopes can be identified with the convex chains that are invertible with respect to $*$.

Similarly to the space of convex chains, define the subspace $\tilde{V}_\square \subset \tilde{V}$ of *convex parachains* using only parapolytopes instead of all polytopes. We will use repeatedly the following example of a parachain.

Example 2.4. Consider the simplest case $d = 1$. Let $[\mu, \nu] \subset \mathbb{R}$ be a segment (i.e., $\mu < \nu$), and $[\nu, \mu]$ — a virtual segment. Using the existence of the operation $*$ satisfying (M), it is easy to check that

$$\iota([\nu, \mu]) = -\mathbb{I}_{[-\nu, -\mu]} + \mathbb{I}_{\{-\nu\}} + \mathbb{I}_{\{-\mu\}}$$

(note that the right hand side is the characteristic function of the open interval $(-\nu, -\mu)$). Indeed,

$$\begin{aligned} \mathbb{I}_{[\mu, \nu]} * (-\mathbb{I}_{[-\nu, -\mu]} + \mathbb{I}_{\{-\nu\}} + \mathbb{I}_{\{-\mu\}}) &= -\mathbb{I}_{[\mu, \nu]} * \mathbb{I}_{[-\nu, -\mu]} + \mathbb{I}_{[\mu, \nu]} * \mathbb{I}_{\{-\nu\}} + \mathbb{I}_{[\mu, \nu]} * \mathbb{I}_{\{-\mu\}} \\ &= -\mathbb{I}_{[\mu-\nu, \nu-\mu]} + \mathbb{I}_{[\mu-\nu, 0]} + \mathbb{I}_{[0, \nu-\mu]} = \mathbb{I}_{\{0\}}. \end{aligned}$$

More generally, if $P \subset \mathbb{R}^d$ is a convex polytope then

$$(-1)^{\dim P} \mathbb{I}_P * \mathbb{I}_{\text{int}(P^\vee)} = \mathbb{I}_{\{0\}},$$

where $P^\vee = \{-x \mid x \in P\}$, and $\text{int}(P^\vee)$ denotes the interior of P^\vee (see [PKh, Section 2, Theorem 2]).

2.3. Divided difference operators on parachains. For each $i = 1, \dots, r$, we now define a *divided difference (or Demazure) operator* D_i on the space of convex parachains \tilde{V}_\square . Let P be a parapolytope. Choose the smallest $j = 1, \dots, d_i$ such that P lies in the hyperplane $\{x_j^i = \text{const}\}$. If no such j exists, then $D_i(\mathbb{I}_P)$ is not defined. Otherwise, we expand P in the direction of x_j^i as follows.

First, suppose that a parapolytope P lies in $(c + \mathbb{R}^{d_i})$ for some $c \in \mathbb{R}^d$, i.e., $P = c + \Pi(\mu, \nu)$ is a coordinate parallelepiped. We always fix the choice of c by requiring that c lies in the direct complement to \mathbb{R}^{d_i} with respect to the decomposition $\mathbb{R}^d = \mathbb{R}^{d_1} \oplus \dots \oplus \mathbb{R}^{d_i} \oplus \dots \oplus \mathbb{R}^{d_r}$. Consider $\nu' = (\nu'_1, \dots, \nu'_{d_i})$, where $\nu'_k = \nu_k$ for all $k \neq j$, and ν'_j is defined by the equality

$$\sum_{k=1}^{d_i} (\mu_k + \nu'_k) = l_i(c).$$

If $\nu'_j \geq \nu_j$, then $D_i^+(P) := c + \Pi(\mu, \nu')$ is a true coordinate parallelepiped. Note that P is a facet of $D_i^+(P)$ unless $\nu' = \nu$.

If $\nu'_j < \nu_j$, define $\mu' = (\mu'_1, \dots, \mu'_{d_i})$ by setting $\mu'_k = \mu_k$ for all $k \neq j$, and $\mu'_j = \nu'_j$. Then $D_i^-(P) := c + \Pi(\mu', \nu)$ is a true coordinate parallelepiped, and P is a facet of $D_i^-(P)$. Let P' be the facet of $D_i^-(P)$ parallel to P .

We now define $D_i(\mathbb{I}_P)$ as follows:

$$D_i(\mathbb{I}_P) = \begin{cases} \mathbb{I}_{D_i^+(P)} & \text{if } \nu_j \leq \nu'_j, \\ -\mathbb{I}_{D_i^-(P)} + \mathbb{I}_P + \mathbb{I}_{P'} & \text{if } \nu_j > \nu'_j. \end{cases}$$

Remark 2.5. This definition is motivated by the following observation. Let μ and ν be integers such that $\mu < \nu$. Define the function $f(\mu, \nu, t)$ of a complex variable t by the formula

$$f(\mu, \nu, t) = t^\mu + t^{\mu+1} + \dots + t^\nu,$$

that is, f is the exponential sum over all integer points in the segment $[\mu, \nu] \subset \mathbb{R}$. Computing the sum of the geometric progression, we get that

$$f(\mu, \nu, t) = \frac{t^\mu - t^{\nu+1}}{1 - t}.$$

This formula gives a meromorphic continuation of $f(\mu, \nu, t)$ to all real μ and ν . In particular, for integer μ and ν such that $\mu > \nu$ we obtain

$$f(\mu, \nu, t) = \frac{t^\mu - t^{\nu+1}}{1 - t} = -(t^{\nu+1} + \dots + t^{\mu-1}),$$

that is, f is minus the exponential sum over all integer points in the open interval $(\nu, \mu) \subset \mathbb{R}$ (cf. Example 2.4).

Definition 3. Let $P \subset \mathbb{R}^d$ be a parapolotope such that P lies in the hyperplane $\{x_j^i = \text{const}\}$ for some j but does not lie in any hyperplane $\{x_k^i = \text{const}\}$ for $k < j$. Define $D_i(\mathbb{I}_P)$ by setting


$$D_i(\mathbb{I}_P) |_{c+\mathbb{R}^{d_i}} = D_i^{(j)}(\mathbb{I}_{P \cap (c+\mathbb{R}^{d_i})})$$

for all c in the complement to \mathbb{R}^{d_i} . The superscript (j) on the right hand side means that we always expand $P \cap (c + \mathbb{R}^{d_i})$ in the direction of x_j^i as explained above (even when $P \cap (c + \mathbb{R}^{d_i})$ for some c lies in a hyperplane $\{x_k^i = \text{const}\}$ for $k < j$).

It is not hard to check that this definition yields a convex chain. In many cases (see examples in Section 3), $D_i(\mathbb{I}_P)$ is the characteristic function of a polytope (and P is a facet of this polytope unless $D_i(\mathbb{I}_P) = \mathbb{I}_P$). This polytope will be denoted by $D_i(P)$. The definition of D_i can be extended by linearity to the other parachains, however, $D_i(\delta)$ for a convex chain δ in general depends on a representation $\delta = \sum_P c_P \mathbb{I}_P$.

The definition immediately implies that similarly to the classical Demazure operators the convex-geometric ones satisfy the identity $D_i^2 = D_i$. It would be interesting to find an analog of braid relations for these operators.

2.4. Examples.


 FIGURE 1. Trapezoids $D_2(P)$ for different segments $P = AB$.

Dimension 2. The simplest meaningful example is $\mathbb{R}^2 = \mathbb{R} \oplus \mathbb{R}$. Label coordinates in \mathbb{R}^2 by $x := x_1^1$ and $y := x_1^2$. Assume that $l_1 = y$ and $l_2 = x$. If $P = \{(\mu_1, \mu_2)\}$ is a point, and $\mu_2 \geq 2\mu_1$, then $D_1(P)$ is a segment:

$$D_1(P) = [(\mu_1, \mu_2), (\mu_2 - \mu_1, \mu_2)].$$

If $\mu_2 < 2\mu_1$, then $D_1(\mathbb{I}_P)$ is a virtual segment, that is,

$$D_1(\mathbb{I}_P) = -\mathbb{I}_{[(\mu_2 - \mu_1, \mu_2), (\mu_1, \mu_2)]} + \mathbb{I}_P + \mathbb{I}_{(\mu_2 - \mu_1, \mu_2)}.$$

If $P = AB$ is a horizontal segment, where $A = (\mu_1, \mu_2)$ and $B = (\nu_1, \mu_2)$, then $D_2(P)$ is the trapezoid $ABCD$ given by the inequalities

$$\mu_1 \leq x \leq \nu_1, \quad \mu_2 \leq y \leq x - \mu_2.$$

See Figure 1 for $D_2(P)$ in the case $\mu_1 = -1$, $\nu_1 = 2$, $\mu_2 = -1$ (left) and $\mu_1 = -1$, $\nu_1 = 2$, $\mu_2 = 0$ (right). In the latter case, the convex chain $D_2(\mathbb{I}_P)$ is equal to

$$\mathbb{I}_{OBC} - \mathbb{I}_{ADO} + \mathbb{I}_{OA} + \mathbb{I}_{DO} - \mathbb{I}_O.$$

Dimension 3. A more interesting example is $\mathbb{R}^3 = \mathbb{R}^2 \oplus \mathbb{R}$. Label coordinates in \mathbb{R}^3 by $x := x_1^1$, $y := x_1^2$ and $z := x_1^3$. Assume that $l_1 = z$ and $l_2 = x + y$. If $P = (\mu_1, \mu_2, \mu_3)$ is a point, then $D_1(P)$ is a segment:

$$D_1(P) = [(\mu_1, \mu_2, \mu_3), (\mu_3 - \mu_1 - 2\mu_2, \mu_2, \mu_3)].$$

Similarly, if $P = [(\mu_1, \mu_2, \mu_3), (\nu_1, \mu_2, \mu_3)]$ is a segment in \mathbb{R}^2 , then $D_1(P)$ is the rectangle given by the equation $z = \mu_3$ and the inequalities

$$\mu_1 \leq x \leq \nu_1, \quad \mu_2 \leq y \leq \mu_3 - \mu_1 - \nu_1 - \mu_2.$$




FIGURE 2. Trapezoid $D_2D_1(P)$ and polytope $D_1D_2D_1(P)$ for a point $P = (0, -3, -3)$

Using the previous calculations, it is easy to show that if $P = (\lambda_2, \lambda_3, \lambda_3)$ is a point and $\lambda_3 < \lambda_2 < -\lambda_2 - \lambda_3$, then $D_1D_2D_1(P)$ is the 3-dimensional Gelfand–Zetlin polytope Q_λ (as defined in Example 2.1) for $\lambda = (\lambda_1, \lambda_2, \lambda_3)$, where $\lambda_1 = -\lambda_2 - \lambda_3$. Indeed, $D_2D_1(P)$ is the trapezoid (see Figure 2) given by the equation $y = \lambda_3$ and the inequalities

$$\lambda_2 \leq x \leq \lambda_1, \quad \lambda_3 \leq z \leq x.$$

Then $D_1D_2D_1(P)$ is the union of all rectangles $D_2(I_a)$ for $a \in [\lambda_3, \lambda_1]$, where I_a is the segment $D_2D_1(P) \cap \{z = a\}$, that is, $I_a = [(\max\{z, \lambda_2\}, \lambda_3, a), (\lambda_1, \lambda_2, a)]$. Hence,

$$\lambda_3 \leq y \leq \min\{\lambda_2, z\}.$$

Similarly to the last example, we construct Gelfand–Zetlin polytopes for arbitrary n using the string space from Example 2.1 (see Theorem 3.4).

3. POLYTOPES AND DEMAZURE CHARACTERS

3.1. Characters of polytopes. For a string space $\mathbb{R}^d = \mathbb{R}^{d_1} \oplus \dots \oplus \mathbb{R}^{d_r}$, denote by $\sigma_i(x)$ the sum of the coordinates of $x \in \mathbb{R}^d$ that correspond to the subspace \mathbb{R}^{d_i} , i.e., $\sigma_i(x) = \sum_{k=1}^{d_i} x_k^i$. With each integer point $x \in \mathbb{R}^d$ in the string space, we associate the *weight* $p(x) \in \mathbb{R}^r$ defined as $(\sigma_1(x), \dots, \sigma_r(x))$. For the rest of the paper, we will always assume that $l_i(x)$ depends only on $p(x)$, that is, l_i comes from a linear

function on \mathbb{R}^r (the latter will also be denoted by l_i). In addition, we assume that l_i is integral, i.e., $l_i(x) \in \mathbb{Z}$ for all $x \in \mathbb{Z}^d$.

Denote the basis vectors in \mathbb{R}^r by $\alpha_1, \dots, \alpha_r$, and denote the coordinates with respect to this basis by (y_1, \dots, y_r) . For each $i = 1, \dots, r$, define the affine reflection $s_i : \mathbb{R}^r \rightarrow \mathbb{R}^r$ by the formula

$$s_i(y_1, \dots, y_i, \dots, y_r) = (y_1, \dots, l_i(y) - y_i, \dots, y_r).$$

Example 3.1. For the string space $\mathbb{R}^d = \mathbb{R}^{n-1} \oplus \mathbb{R}^{n-2} \oplus \dots \oplus \mathbb{R}^1$ from Example 2.1, define the functions l_i by the formula

$$l_i(x) = \sigma_{i-1}(x) + \sigma_{i+1}(x),$$

where we put $\sigma_0 = \sigma_n = 0$. Identify \mathbb{R}^{n-1} with the weight lattice of SL_n so that α_i is identified with the i -th simple root. In this case, the reflection s_i coincides with the simple reflection in the hyperplane perpendicular to the root α_i .

We now consider the ring R of Laurent polynomials in the formal exponentials $t_1 := e^{\alpha_1}, \dots, t_r := e^{\alpha_r}$ (that is, R is the group algebra of the lattice $\mathbb{Z}^r \subset \mathbb{R}^r$). Let $P \subset \mathbb{R}^d$ be a *lattice* polytope in the string space, i.e., the vertices of P belong to \mathbb{Z}^r . Define the *character of P* as the sum of formal exponentials $e^{p(x)}$ over all integer points x inside and at the boundary of P :

$$\chi(P) := \sum_{x \in P \cap \mathbb{Z}^d} e^{p(x)}.$$

In particular, if $d = r$, then $\chi(P)$ is exactly the integer point transform of P . The R -valued function χ can be extended by linearity to all *lattice* convex chains, that is, to the chains $\sum_P c_P \mathbb{I}_P$ such that P is a lattice polytope and $c_P \in \mathbb{Z}$.

Define the *Demazure operator* T_i on R as follows:

$$T_i f = \frac{f - t_i \cdot s_i f}{1 - t_i},$$

where the action of s_i on R is defined by $s_i e^\lambda := e^{s_i(\lambda)}$ for $\lambda \in \mathbb{Z}^r$. For the string space of Example 3.1, these operators reduce to the classical Demazure operators on the group algebra of the weight lattice of SL_n .

The following result motivates the definition of divided difference operators D_i on convex chains (Definition 3).

Theorem 3.2. *Let $P \subset \mathbb{R}^d$ be a lattice parapolytope. Then*

$$\chi(D_i(\mathbb{I}_P)) = T_i \chi(P).$$

Proof. By definition of $D_i(\mathbb{I}_P)$, it suffices to prove this identity when $P = c + \Gamma$, where c lies in the complement to \mathbb{R}^{d_i} and $\Gamma := \Pi(\mu, \nu) \subset \mathbb{R}^{d_i}$ is a coordinate parallelepiped. Then

$$\chi(P) = e^{p(c)} \sum_{z \in \Gamma \cap \mathbb{Z}^{d_i}} t_i^{\sigma(z)}.$$

Hence,

$$T_i(\chi(P)) = e^{p(c)} T_i \left(\sum_{z \in \Gamma \cap \mathbb{Z}^{d_i}} t_i^{\sigma(z)} \right).$$

Recall that by definition of ν' we have

$$\sum_{k=1}^{d_i} (\mu_k + \nu'_k) = l_i(c).$$

Assume that $\nu'_j \geq \nu_j$. Let Π denote $\Pi(\mu, \nu')$. Then Γ , Π and T_i satisfy the hypothesis of [KST, Proposition 6.3]. Applying this proposition we get that

$$T_i \left(\sum_{z \in \Gamma \cap \mathbb{Z}^{d_i}} t_i^{\sigma(z)} \right) = \sum_{z \in \Pi \cap \mathbb{Z}^{d_i}} t_i^{\sigma(z)}.$$

Hence, $T_i(\chi(P)) = \chi(D_i(P))$.

The case $\nu'_j < \nu_j$ is completely analogous. \square

Note that Theorem 3.2 for $d_i = 1$ follows directly from the definitions of T_i and D_i (see Remark 2.5).

Example 3.3. Figure 3 illustrates Theorem 3.2 when $d_i = 2$ and $P = c + \Gamma$ where $\Gamma \subset \mathbb{R}^{d_i}$ is the segment $[(-1, -1), (2, -1)]$. Namely, $T_i(t_i^{x_1^i + x_2^i})$ is equal to the character of the segment $[(x_1^i, x_2^i), (x_1^i, l_i(c) - 2x_1^i - x_2^i)]$ for every $(x_1^i, x_2^i) \in \Gamma \cap \mathbb{Z}^2$ by definition of T_i . Hence,

$$\sum_{(x_1^i, x_2^i) \in \Gamma \cap \mathbb{Z}^2} T_i(t_i^{x_1^i + x_2^i})$$

for $l_i(c) = 3$ coincides with the character of the trapezoid shown on Figure 3 (left). It is easy to construct a bijective correspondence between the integer points in the trapezoid and those in the rectangle $D_i(P)$ in such a way that the sum of coordinates is preserved. The former are marked by black dots, and the latter by empty circles.

Theorem 3.2 allows one to construct various polytopes (possibly virtual) and convex chains whose characters yield the Demazure characters (in particular, the Weyl character) of irreducible representations of reductive groups (see Section 3.3). The same character can be captured using string spaces for different partitions $d = d_1 + d_2 + \dots + d_n$ (see Section 4.3). The case $d_1 = \dots = d_n = 1$ produces polytopes with very simple combinatorics, namely, multidimensional versions of trapezoids that are combinatorially equivalent to cubes (they are called *twisted cubes* in [GK]). However, twisted cubes that represent the Weyl characters are virtual. Considering string spaces with $d_i > 1$ allows one to represent the Weyl character by a true though more intricate polytope (see Example 3.4 for SL_3 and Section 4.3). The reason is illustrated by Figure 3 (right) that depicts a virtual trapezoid and a (true) rectangle




FIGURE 3. Rectangle and trapezoid yield the same character

with the same character. Note that the point $(2, 1)$ (marked by -1) contributes a negative summand to the character of the trapezoid.

3.2. Gelfand–Zetlin polytopes for SL_n . Let $\mathbb{R}^d = \mathbb{R}^{n-1} \oplus \mathbb{R}^{n-2} \oplus \dots \oplus \mathbb{R}^1$ be the string space of rank $(n - 1)$ from Example 3.1. The theorem below shows how to construct the classical Gelfand–Zetlin polytopes (see Example 2.1) via the convex-geometric Demazure operators D_1, \dots, D_{n-1} .

Theorem 3.4. *For every strictly dominant weight $\lambda = (\lambda_1, \dots, \lambda_n)$ (that is, $\lambda_1 > \dots > \lambda_n$) of GL_n such that $\lambda_1 + \dots + \lambda_n = 0$, the Gelfand–Zetlin polytope Q_λ coincides with the polytope*

$$[(D_1)(D_2 D_1)(D_3 D_2 D_1) \dots (D_{n-1} \dots D_1)](a_\lambda),$$

where $a_\lambda \in \mathbb{R}^d$ is the point $(\lambda_2, \dots, \lambda_n; \lambda_3, \dots, \lambda_n; \dots; \lambda_n)$.

Proof. Let us define the polytope

$$P_\lambda(i, j) := \left[(\hat{D}_{n-j} \dots \hat{D}_i D_{i-1} \dots D_1) \dots (D_{n-1} \dots D_1) \right](a_\lambda)$$

for every pair (i, j) such that $1 \leq i \leq (n - j) \leq (n - 1)$. Put $x_l^0 = \lambda_l$ for $l = 1, \dots, n$. We will show by induction on dimension that $P_\lambda(i, j)$ is the face of the Gelfand–Zetlin polytope Q_λ given by the equations $x_l^k = x_{l+1}^{k-1}$ for all pairs (k, l) such that either $l > j$, or $l = j$ and $k \geq i$. The induction base is $P_\lambda(1, 1) = a_\lambda$, which is clearly a vertex of Q_λ by our assumption. The induction step follows from Lemma

(recall that $\sigma_i(x) = \sum_{j=1}^{d_i} x_j^i$). Here (α_k, α_i) is determined by the simple reflection s_i as follows:

$$s_i(\alpha_k) = \alpha_k + (\alpha_k, \alpha_i)\alpha_i,$$

(that is, the function (\cdot, α_i) is minus the coroot corresponding to α_i). In particular, if $G = SL_n$ and $w_0 = (s_1)(s_2s_1)(s_3s_2s_1) \dots (s_{n-1} \dots s_1)$, then we get the string space from Example 3.1.

Define the projection p of the string space to the real span \mathbb{R}^r of the weight lattice of G by the formula $p(x) = \sigma_1(x)\alpha_1 + \dots + \sigma_r(x)\alpha_r$. Note that after identifying the root lattice of G with $\mathbb{Z}^r \subset \mathbb{R}^r$ the map p is the same as the map p defined earlier in Section 3.1.

Theorem 3.6. *For every dominant weight λ in the root lattice of G , and every point $a_\lambda \in \mathbb{Z}^d$ such that $p(a_\lambda) = w_0\lambda$ the convex chain*

$$P_\lambda := D_{i_1}D_{i_2} \dots D_{i_d}(a_\lambda)$$

yields the Weyl character $\chi(V_\lambda)$ of the irreducible G -module V_λ , that is,

$$\chi(V_\lambda) = \chi(P_\lambda).$$

Proof. By the Demazure character formula [Andersen] we have

$$\chi(V_\lambda) = T_{i_1} \dots T_{i_d} e^{w_0\lambda}.$$

This formula together with Theorem 3.2 implies by induction the desired statement. \square

As a corollary, we get that $p_*(P_\lambda)$ is the weight polytope of V_λ in \mathbb{R}^r . Here p_* denotes the push-forward of convex chains (see [PKh, Proposition-Definition 2]).

Remark 3.7. A slight modification of Theorem 3.2 makes it applicable to all dominant weights (not only those inside the root lattice). Namely, instead of the lattices $\mathbb{Z}^d \subset \mathbb{R}^d$ and $\mathbb{Z}^r \subset \mathbb{R}^r$ one should consider the shifted lattices $a_\lambda + \mathbb{Z}^d \subset \mathbb{R}^d$ and $\lambda + \mathbb{Z}^r \subset \mathbb{R}^r$, and define characters of polytopes with respect to these new lattices. The convex chain P_λ will be lattice with respect to the lattice $a_\lambda + \mathbb{Z}^d$.

In the same way, we can construct convex chains that capture the characters of Demazure submodules of V_λ for any element w in the Weyl group and a reduced decomposition $w = s_{j_1} \dots s_{j_\ell}$ (see Corollary 4.5). In particular, if $s_{j_1} \dots s_{j_\ell}$ is a terminal subword of $s_{i_1} s_{i_2} \dots s_{i_d}$ (that is, $j_\ell = i_d, j_{\ell-1} = i_{d-1}$, etc.) then the corresponding convex chain is a face of P_λ . It is interesting to check whether this convex chain is always a true polytope. One way to do this would be to identify it with an Okounkov polytope of the Bott–Samelson resolution corresponding to the word $s_{j_1} \dots s_{j_\ell}$ (see Conjecture 4.1).

3.4. Examples.

Sp(4). Take $G = Sp(4)$ and $w_0 = s_2 s_1 s_2 s_1$ (here α_1 denotes the shorter root and α_2 denotes the longer one). The corresponding string space of rank 2 is $\mathbb{R}^4 = \mathbb{R}^2 \oplus \mathbb{R}^2$ together with $l_1 = 2(x_1^2 + x_2^2)$ and $l_2 = x_1^1 + x_2^1$. Let $\lambda = -p_1 \alpha_1 - p_2 \alpha_2$ be a dominant weight, that is, $\lambda_1 := (p_2 - p_1) \geq 0$ and $\lambda_2 := (p_1 - 2p_2) \geq 0$. Choose a point $a_\lambda = (a, b, c, d)$ such that $(a + b) = p_1$ and $(c + d) = p_2$ (that is, $p(a_\lambda) = w_0 \lambda = -\lambda$). Label coordinates in \mathbb{R}^4 by $x := x_1^1$, $y := x_2^1$, $z := x_1^2$ and $t := x_2^2$. Then the polytope $D_2 D_1 D_2 D_1(a_\lambda)$ is given by inequalities

$$0 \leq x - a \leq 2\lambda_1, \quad z - c \leq x - a + \lambda_2, \quad y - b \leq 2(z - c),$$

$$y - b \leq z - c + \lambda_2, \quad 0 \leq t - d \leq \lambda_2, \quad t - d \leq \frac{y - b}{2}.$$

It is not hard to show that the polytopes $D_1 D_2 D_1 D_2(a_\lambda)$ and $D_2 D_1 D_2 D_1(a_\lambda)$ are the same up to a linear transformation of \mathbb{R}^4 . Each polytope has 11 vertices, hence, they are not combinatorially equivalent to string polytopes for $s_1 s_2 s_1 s_2$ or $s_2 s_1 s_2 s_1$ defined in [L].

SL(3). Take $G = SL(3)$ and $w_0 = s_1 s_2 s_1$. The corresponding string space of rank 2 coincides with the one from Section 2.4, namely, $\mathbb{R}^3 = \mathbb{R}^2 \oplus \mathbb{R}$, and $l_1 = x_1^2$, $l_2 = x_1^1 + x_2^1$. If $a_\lambda = (b, c, c)$ where $-b - c \geq b \geq c$, then the polytope $D_1 D_2 D_1(a_\lambda)$ is the Gelfand–Zetlin polytope Q_λ for $\lambda = (-b - c, b, c)$. Label coordinates in \mathbb{R}^3 by $x := x_1^1$, $y := x_2^1$ and $z := x_1^2$. We now introduce a different structure of a string space in \mathbb{R}^3 by splitting \mathbb{R}^2 , namely, $\mathbb{R}^3 = \mathbb{R}^1 \oplus \mathbb{R}^1 \oplus \mathbb{R}^1$ with coordinates $\tilde{x}_1^1, \tilde{x}_2^1, \tilde{x}_3^1$ such that $\tilde{x}_1^1 = x$, $\tilde{x}_2^1 = z$, $\tilde{x}_3^1 = y$. Put $\tilde{l}_1 = l_1 - 2y$, $\tilde{l}_2 = l_2$ and $\tilde{l}_3 = l_1 - 2x$. It is easy to check that $\tilde{D}_2 \tilde{D}_1(a_\lambda) = D_2 D_1(a_\lambda)$ and deduce by arguments of Example 3.3 that the virtual polytope $\tilde{D}_3 \tilde{D}_2 \tilde{D}_1(a_\lambda)$ (see Figure 4) has the same character as the polytope $D_1 D_2 D_1(a_\lambda)$. In particular, the image of $\tilde{D}_3 \tilde{D}_2 \tilde{D}_1(a_\lambda)$ under the projection $(x, y, z) \mapsto (x + y, z)$ coincides with the weight polytope of the irreducible representation of SL_3 with the highest weight $-c\alpha_1 - (a + b)\alpha_2$ (provided that the latter is dominant, that is, $a + b - 2c \geq 0$ and $c - 2(a + b) \geq 0$). The virtual polytope $\tilde{D}_3 \tilde{D}_2 \tilde{D}_1(a_\lambda)$ is a twisted cube of Grossberg–Karshon (cf. [GK, Figure 2]) given by the inequalities

$$a \leq x \leq c - 2b - a, \quad c \leq z \leq x + b - c, \quad b \leq y \leq -2x + z - b. \quad (GK)$$

Note that the last pair of inequalities is inconsistent when $b > -2x + z - b$, and should be interpreted in the sense of convex chains. More precisely, $\tilde{D}_3 \tilde{D}_2 \tilde{D}_1(a_\lambda) = \mathbb{I}_P - \mathbb{I}_Q$, where P is the convex polytope given by inequalities (GK) and Q is the set given by the inequalities

$$a \leq x \leq c - 2b - a, \quad c \leq z \leq x + b - c, \quad b > y > -2x + z - b.$$

(cf. [GK, Formula (2.21)]). A generalization of this example will be given in Section 4.3.




FIGURE 4. Virtual polytope $\tilde{D}_3\tilde{D}_2\tilde{D}_1(a_\lambda)$ for $a_\lambda = (0, -3, -3)$

4. BOTT TOWERS AND BOTT–SAMELSON RESOLUTIONS

In this section, we outline possible algebro-geometric applications of the convex-geometric Demazure operators.

4.1. Bott towers. Let us recall the definition of a *Bott tower* (see [GK] for more details). It is a toric variety obtained from a point by iterating the following step. Let X be a toric variety, and L a line bundle on X . Define a new toric variety $Y := \mathbb{P}(L \oplus \mathcal{O}_X)$ as the projectivization of the split rank two vector bundle $L \oplus \mathcal{O}_X$ on X . Consider a sequence of toric varieties

$$Y_0 \leftarrow Y_1 \leftarrow \dots \leftarrow Y_d,$$

where Y_0 is a point, and $Y_i = \mathbb{P}(L_{i-1} \oplus \mathcal{O}_{Y_{i-1}})$ for a line bundle L_{i-1} on Y_{i-1} . In particular, $Y_1 = \mathbb{P}^1$ and $Y_2 = \mathbb{P}(\mathcal{O}_{\mathbb{P}^1} \oplus \mathcal{O}_{\mathbb{P}^1}(k))$ is a Hirzebruch surface. We call Y_d the *Bott tower* corresponding to the collection of line bundles (L_1, \dots, L_{d-1}) . Note that the collection (L_1, \dots, L_{d-1}) depends on $\frac{d(d-1)}{2}$ integer parameters since $\text{Pic}(Y_i) = \mathbb{Z}^i$. Recall that the Picard group of a toric variety of dimension d can be identified with a group of virtual lattice polytopes in \mathbb{R}^d in such a way that very ample line bundles get identified with their Newton polytopes. One can describe the (possibly virtual) polytope $P(\mathcal{L})$ of a given line bundle \mathcal{L} on Y_d using a suitable string space.

Consider a string space with $d = r$, that is, $d_1 = \dots = d_r = 1$. We have the decomposition

$$\mathbb{R}^d = \underbrace{\mathbb{R} \oplus \dots \oplus \mathbb{R}}_d.$$

Label coordinates in \mathbb{R}^d as follows: $x_1^i := y_i$ for $i = 1, \dots, d$. Since we will be interested in the polytope $P := D_1 \dots D_d(a)$, we can assume that the linear function l_i for $i < d$ does not depend on y_1, \dots, y_i , and $l_d = y_1$. Hence, the collection (l_1, \dots, l_{d-1}) of linear functions also depends on $\frac{d(d-1)}{2}$ parameters.

The projective bundle formula gives a natural basis (η_1, \dots, η_d) in the Picard group of Y_d . Namely, for $d = 1$, the basis in $\text{Pic}(\mathbb{P}^1)$ consists of the class of a point in \mathbb{P}^1 . We now proceed by induction. Let $(\eta_1, \dots, \eta_{i-1})$ be the basis in $\text{Pic}(Y_{i-1})$ (we identify $\text{Pic}(Y_{i-1})$ with its pull-back to $\text{Pic}(Y_d)$). Put $\eta_i = c_1(\mathcal{O}_{Y_i}(1))$ where $c_1(\mathcal{O}_{Y_i}(1))$ denotes the first Chern class of the tautological quotient line bundle $\mathcal{O}_{Y_i}(1)$ on Y_i . Decompose L_1, \dots, L_{d-1} in the basis (η_1, \dots, η_d) :

$$L_1 = a_{1,1}\eta_1, \quad \dots, \quad L_{d-1} = a_{d-1,1}\eta_1 + \dots + a_{d-1,d-1}\eta_{d-1}. \quad (*)$$

Similarly, decompose l_1, \dots, l_{d-1} in the basis of coordinate functions (y_1, \dots, y_d) :

$$l_1 = b_{1,1}y_2 + \dots + b_{1,d-1}y_d, \quad \dots, \quad l_{d-1} = b_{d-1,d-1}y_d. \quad (**)$$

Let Y_d be the Bott tower corresponding to the collection $(*)$ of line bundles, and \mathbb{R}^d the string space corresponding to the collection $(**)$ of linear functions. One can show (cf. [GK, Theorem 3]) that if $a_{i,j} = b_{j,i}$, then there exists $a_{\mathcal{L}} \in \mathbb{R}^d$ such that

$$P(\mathcal{L}) = D_1 D_2 \dots D_d(a_{\mathcal{L}}).$$

In particular, when \mathcal{L} is very ample the polytope $D_1 D_2 \dots D_d(a_{\mathcal{L}})$ coincides with the Newton polytope of the pair (Y_d, \mathcal{L}) . Note that the intermediate polytopes $\{a_{\mathcal{L}}\} \subset D_d(a_{\mathcal{L}}) \subset D_{d-1} D_d(a_{\mathcal{L}}) \subset \dots$ correspond to the flag of toric subvarieties $Z_0 = \{pt\} \subset Z_1 \subset \dots \subset Z_d = Y_d$, where $Z_i = p_{d-i}^{-1}(Z_0)$ and p_i is the projection $p_i : Y_d \rightarrow Y_i$.

4.2. Bott–Samelson varieties. Similarly to Bott towers, Bott–Samelson varieties can be obtained by successive projectivizations of rank two vector bundles. In general, these bundles are no longer split, so the resulting varieties are not toric. In [GK], Bott–Samelson varieties were degenerated to Bott towers by changing complex structure (in particular, Bott–Samelson varieties are diffeomorphic to Bott towers when regarded as real manifolds). Below we define these varieties using notation of Section 3.3.

Fix a Borel subgroup $B \subset G$. With every collection of simple roots $(\alpha_{i_1}, \dots, \alpha_{i_\ell})$, one can associate a *Bott–Samelson variety* $R_{(i_1, \dots, i_\ell)}$ and a map $R_{(i_1, \dots, i_\ell)} \rightarrow G/B$ by the following inductive procedure. Put $R_\emptyset = pt$. For every ℓ -tuple $I = (i_1, \dots, i_\ell)$ denote by I^j the $(\ell - 1)$ -tuple $(i_1, \dots, \hat{i}_j, \dots, i_\ell)$. Define R_I as the fiber product $R_{I^\ell} \times_{G/P_\ell} G/B$, where P_{i_ℓ} is the minimal parabolic subgroup corresponding to the




FIGURE 5. Polytope $D_1E_uD_2D_1(a)$ for $a = (0, -1, -1)$ and $u = (0, -1/2, 0)$

root α_{i_ℓ} . The map $r_I : R_I \rightarrow G/B$ is defined as the projection to the second factor. There is a natural embedding

$$R_{I^\ell} \hookrightarrow R_I; \quad x \mapsto (x, r_I(x)).$$

In particular, any subsequence $J \subset I$ yields the embedding $R_J \hookrightarrow R_I$.

It follows from the projective bundle formula that the Picard group of R_I is freely generated by the divisors $R_{I^1}, \dots, R_{I^\ell}$. Denote by v the geometric valuation on $\mathbb{C}(R_I)$ defined by the flag $R_\emptyset \subset R_{(i_\ell)} \subset R_{(i_{\ell-1}, i_\ell)} \subset \dots \subset R_{I^1} \subset R_I$. Let \mathcal{L} be a line bundle on R_I , and $P_v(\mathcal{L})$ its Okounkov body with respect to the valuation v . Conjecturally, $P_v(\mathcal{L})$ can be described using string spaces as follows.

Replace a reduced decomposition of w_0 in the definition of the string space from Section 3.3 by a sequence $(\alpha_{i_1}, \dots, \alpha_{i_\ell})$ that defines R_I (we no longer require that $s_{i_1} \dots s_{i_\ell}$ be reduced). More precisely, let d_i the number of α_{i_j} in this sequence such that $i_j = i$. We get the following string space \mathcal{S}_I of rank $\leq r$ and dimension ℓ :

$$\mathbb{R}^\ell = \mathbb{R}^{d_1} \oplus \dots \oplus \mathbb{R}^{d_r},$$

where the functions l_i are given by the formula:

$$l_i(x) = \sum_{k \neq i} (\alpha_k, \alpha_i) \sigma_k(x). \quad (BS)$$

In particular, if $\ell = d$ and $s_{i_1} \dots s_{i_\ell}$ is reduced then R_I is a Bott–Samelson resolution of the flag variety G/B , and \mathbb{R}^ℓ is exactly the string space from Section 3.3. Denote by E_u the parallel translation in the string space by a vector $u \in \mathbb{R}^\ell$.

Conjecture 4.1. *For every line bundle \mathcal{L} on R_I , there exists a point $\mu \in \mathbb{R}^r$ and vectors $u_1, \dots, u_\ell \in \mathbb{R}^\ell$ such that we have*

$$P_v(\mathcal{L}) = E_{u_1} D_{i_1} E_{u_2} D_{i_2} \dots E_{u_\ell} D_{i_\ell}(a_\mu)$$

for any point $a_\mu \in \mathbb{R}^\ell$ that satisfies $p(a) = \mu$.

In particular, if $\mathcal{L} = r_I^* \mathcal{L}(\lambda)$, where $\mathcal{L}(\lambda)$ is the line bundle on G/B corresponding to the dominant weight λ , then one can take $u_2 = \dots = u_\ell = 0$ and $\mu = \lambda$. This conjecture agrees with the example computed in [Anderson, Section 6.4] for SL_3 and the Bott–Samelson resolution $R_{(1,2,1)}$ (cf. Figure 5 and Figure 3(b) in loc.cit.). Figure 5 shows the polytope $D_1 E_u D_2 D_1(a)$ for the string space (BS) when $G = SL_3$ and $I = (1, 2, 1)$.

4.3. Degenerations of string spaces. While twisted cubes of Grossberg–Karshon for GL_n and Gelfand–Zetlin polytopes have different combinatorics they produce the same Demazure characters. We now reproduce this phenomenon for general string spaces. In particular, we transform the string space (BS) from Section 4.2 into a string space from Section 4.1.

Let \mathcal{S} be a string space $\mathbb{R}^d = \mathbb{R}^{d_1} \oplus \dots \oplus \mathbb{R}^{d_r}$ with functions l_1, \dots, l_r . Suppose that $d_i > 1$.

Definition 4. The i -th degeneration of the string space \mathcal{S} is the string space

$$\mathbb{R}^d = \mathbb{R}^{d_1} \oplus \dots \oplus \underbrace{\mathbb{R}^{d_i-1} \oplus \mathbb{R}^1}_{\mathbb{R}^{d_i}} \oplus \dots \oplus \mathbb{R}^{d_r}$$

of rank $(r + 1)$ with functions $l_1, \dots, l'_i, l''_i, \dots, l_r$, where

$$l'_i(x) = l_i(x) - 2x_{d_i}^i; \quad l''_i(x) = l_i(x) - 2 \sum_{k=1}^{d_i-1} x_k^i.$$

Example 4.2. The string space $\mathbb{R} \oplus \mathbb{R} \oplus \mathbb{R}$ from Example 3.4 is the 1-st degeneration of the space $\mathbb{R}^2 \oplus \mathbb{R}$ with $l_1 = x_1^1 + x_2^1$, $l_2 = x_1^2$.

Define the projection $p_i : \mathbb{R}^{r+1} \rightarrow \mathbb{R}^r$ by sending $(y_1, \dots, y'_i, y''_i, \dots, y_r)$ to $(y_1, \dots, y'_i + y''_i, \dots, y_r)$. This projection induces a homomorphism of group algebras of the lattices \mathbb{Z}^{r+1} and \mathbb{Z}^r , which we will also denote by p_i . It is easy to check that

$$T_i \circ p_i = p_i \circ T'_i = p_i \circ T''_i.$$

Combining this observation with Theorem 3.2, we get the following proposition.

Proposition 4.3. *For a lattice polytope $P \subset \mathbb{R}^d$, we have*

$$\chi(D_i(P)) = p_i(\chi(D'_i(P))) = p_i(\chi(D''_i(P))).$$

We now degenerate successively the string spaces from Section 4.2. Let $I = (\alpha_{i_1}, \dots, \alpha_{i_\ell})$ be a sequence of simple roots, and

$$\mathbb{R}^\ell = \mathbb{R}^{d_1} \oplus \dots \oplus \mathbb{R}^{d_r}$$

is the corresponding string space \mathcal{S}_I with the functions l_1, \dots, l_r given by (BS) .

Let $\tilde{\mathcal{S}}_I$ be the string space of rank ℓ obtained from \mathcal{S}_I by $(d_1 - 1)$ first degenerations, $(d_2 - 1)$ second degenerations etc., that is,

$$\tilde{\mathcal{S}}_I = \underbrace{\mathbb{R}_1^{(1)} \oplus \dots \oplus \mathbb{R}_{d_1}^{(1)}}_{\mathbb{R}^{d_1}} \oplus \dots \oplus \underbrace{\mathbb{R}_1^{(\ell)} \oplus \dots \oplus \mathbb{R}_{d_\ell}^{(\ell)}}_{\mathbb{R}^{d_\ell}},$$

where the functions $l_j^{(i)}$ are given by the formula:

$$l_j^{(i)}(x) = l_i(x) - 2 \sum_{k \neq j} x_k^i.$$

Denote the Demazure operators associated with $\tilde{\mathcal{S}}_I$ by $\tilde{D}_j^{(i)}$.

For a point $a \in \mathbb{R}^\ell$, consider the convex chain $P_I = D_{i_1} \dots D_{i_\ell}(a)$. For every $k = 1, \dots, r$, we now replace the rightmost D_k in the expression $D_{i_1} \dots D_{i_\ell}$ by $D_1^{(k)}$, the next one by $D_2^{(k)}, \dots$, the leftmost by $D_{i_k}^{(k)}$. Denote the resulting convex chain by \tilde{P}_I .

Example 4.4. If $r = 3$, $\ell = 6$ and $I = (\alpha_1, \alpha_2, \alpha_1, \alpha_3, \alpha_2, \alpha_1)$, then

$$P_I = D_1 D_2 D_1 D_3 D_2 D_1, \quad \tilde{P}_I = \tilde{D}_3^{(1)} \tilde{D}_2^{(2)} \tilde{D}_2^{(1)} \tilde{D}_1^{(3)} \tilde{D}_1^{(2)} \tilde{D}_1^{(1)}(a).$$

Proposition 4.3 implies that P_I and \tilde{P}_I have the same character (with respect to the map $p : x \mapsto \sigma_1(x)\alpha_1 + \dots + \sigma_r(x)\alpha_r$).

Corollary 4.5. *If $p(a)$ is dominant and $s_{i_1} \dots s_{i_\ell}$ is reduced, then the corresponding Demazure character coincides with $\chi(P_I) = \chi(\tilde{P}_I)$.*

The proof is completely analogous to the proof of Theorem 3.6.

Remark 4.6. Note that the convex chain \tilde{P}_I coincides with the twisted cube constructed in [GK] for the corresponding Bott–Samelson resolution. Indeed, $(\alpha_i, \alpha_i) = -2$ according to our definition of the function (\cdot, α_i) (see Section 3.3), hence, we can write

$$l_j^{(i)}(x) = \sum_{(p,q) \neq (i,j)} (\alpha_p, \alpha_i) x_q^p.$$

It is now easy to check that the defining inequalities for \tilde{P}_I coincide with the inequalities given by [GK, Formula (2.21)] together with computations of [GK, Section 3.7].

The string space \mathcal{S}_I and its complete degeneration $\tilde{\mathcal{S}}_I$ are two extreme cases that yield convex chains for given Demazure characters. By taking partial degenerations of \mathcal{S}_I one can construct intermediate convex chains with the same character. However, only \mathcal{S}_I might produce true polytopes (such as Gelfand–Zetlin polytopes for $G = SL_n$ or polytope of Example 3.4 for $G = Sp_4$) that represent the Weyl characters. Indeed, such a polytope must have a face $D_i^2(a)$ (in the case of \mathcal{S}_I) or

a face $D'_i D''_i(a)$ (in the case of the i -th degeneration of \mathcal{S}_I). The former is a true segment since $D_i^2 = D_i$, while the latter is necessarily a virtual trapezoid with the same character due to cancelations (cf. Figure 3).

REFERENCES

- [Andersen] H. H. ANDERSEN, *Schubert varieties and Demazure's character formula*, Invent. Math., **79** (1985), no. 3, 611–618
- [Anderson] D. ANDERSON, *Okounkov bodies and toric degenerations*, preprint arXiv:1001.4566v3 [math.AG], to appear in Math. Ann.
- [BZ] A. BERENSTEIN, A. ZELEVINSKY, *Tensor product multiplicities, canonical bases and totally positive varieties*, Invent. Math. **143** (2001), no.1, 77–128
- [GK] M. GROSSBERG AND Y. KARSHON, *Bott towers, complete integrability, and the extended character of representations*, Duke Math. J., **76** (1994), no. 1, 23–58.
- [K] K. KAVEH, *Crystal basis and Newton–Okounkov bodies*, preprint arXiv:1101.1687 [math.AG]
- [KKh] K. KAVEH, A. KHOVANSKII, *Convex bodies associated to actions of reductive groups*, Moscow Math. J., **12** (2012), 369–396
- [PKh] A. G. KHOVANSKII, A. V. PUKHLIKOV, *Finitely additive measures of virtual polytopes*, St. Petersburg Mathematical Journal, **4** (1993), no.2, 337–356
- [PKh2] A. G. KHOVANSKII AND A. V. PUKHLIKOV, *The Riemann–Roch theorem for integrals and sums of quasipolynomials on virtual polytopes*, Algebra i Analiz, **4** (1992), no.4, 188–216
- [KST] V. KIRITCHENKO, E. SMIRNOV, V. TIMORIN, *Schubert calculus and Gelfand–Zetlin polytopes*, Russian Math. Surveys, **67** (2012), no.4, 685–719
- [L] P. LITTELMANN, *Cones, crystals and patterns*, Transformation Groups, **3** (1998), pp. 145–179
- [KnM] A. KNUTSON AND E. MILLER, *Gröbner geometry of Schubert polynomials*, Ann. of Math. (2), **161** (2005), 1245–1318
- E-mail address:* vkiritch@hse.ru

LABORATORY OF ALGEBRAIC GEOMETRY AND FACULTY OF MATHEMATICS, HIGHER SCHOOL OF ECONOMICS, VAVILOVA ST. 7, 112312 MOSCOW, RUSSIA

INSTITUTE FOR INFORMATION TRANSMISSION PROBLEMS, MOSCOW, RUSSIA