

DOUBLE PANTS DECOMPOSITIONS OF 2-SURFACES

ANNA FELIKSON AND SERGEY NATANZON

Dedicated to Sabir Gusein-Zade on the occasion of his 60th birthday

ABSTRACT. We consider the union of two pants decompositions of the same orientable 2-dimensional surface of any genus g . Each pants decomposition corresponds to a handlebody bounded by this surface, so two pants decompositions correspond to a Heegaard splitting of a 3-manifold. We introduce a groupoid acting on double pants decompositions. This groupoid is generated by two simple transformations (called flips and handle twists), each transformation changing only one curve of the double pants decomposition. We prove that the groupoid acts transitively on all double pants decompositions corresponding to Heegaard splittings of a 3-dimensional sphere. As a corollary, we prove that the mapping class group of the surface is contained in the groupoid.

2000 MATH. SUBJ. CLASS. 57M50.

KEY WORDS AND PHRASES. Pants decomposition, Heegaard splitting, Mapping class group.

INTRODUCTION

Consider a 2-dimensional orientable surface S of genus g with n holes. A pants decomposition of S is a decomposition into 3-holed spheres (called “pairs of pants”). For each pants decomposition P of S one may construct a handlebody S_+ such that S is the boundary of S_+ and all curves of P are contractible inside S_+ . A union of two pants decompositions of the same surface define two different handlebodies bounded by S , attaching these handlebodies along S one obtains a Heegaard splitting of some 3-manifold.

Below we define double pants decompositions of surfaces as a union of two pants decompositions (with an additional assumption that the homology classes of the curves contained in the double pants decomposition generate the entire homology lattice $H_1(S, \mathbb{Z})$). We introduce a simple groupoid acting on double pants decompositions and prove that this groupoid acts transitively on all double pants decompositions (of the same surface) resulting in Heegaard splitting of a 3-sphere.

More precisely, there are two simple invertible operations acting on the set of double pants decompositions. One of these operations is a flip (or Whitehead move)

Received June 18, 2010; in revised form December 07, 2010.

Research of the second named author is supported by grants RFBR 10-01-00678 and NSH 709.2008.1.

performed in any of the decompositions. Another operation is called a “handle twist”, it is performed only if a handle is cut out by a curve contained in both pants decompositions. In this case the handle contains exactly one curve from each of the two pants decompositions, and one applies a Dehn twist along one of them to the other.

Main Theorem. *Let S be a topological 2-surface of genus g with n holes, where $2g + n > 2$. Then the groupoid generated by flips and handle twists acts transitively on the set of all double pants decompositions of S corresponding to Heegaard splittings of 3-sphere.*

It is possible to show that the mapping class group acts effectively on double pants decompositions of some special combinatorial class. Together with the Main Theorem this implies the following result:

Corollary. *The category of double pants decompositions of a topological surface with morphisms generated by flips and handle twists contains a subcategory isomorphic to a category of topological surfaces with the mapping class group as the set of morphisms.*

The idea to consider two pants decompositions as a Heegaard splitting of a 3-manifold already appeared in the literature ([3], [8], [11]). The main tool of study there is the curve complex defined by Harvey [4], which is a simplicial complex whose vertices are homotopy classes of closed curves on the surface and simplices are spanned by mutually disjoint curves. In particular, Hempel [8] defines a distance between two pants decompositions in terms of the distance in the curve complex and uses this distance to describe topological properties of the resulting 3-manifolds. Our construction is also naturally related to the pants complex described by Hatcher and Thurston ([7], [5]) and to investigation of the mapping class group through various complexes ([1], [2], [10], [9], [12] and many others). The notion of a groupoid generated by flips and twists on the Heegaard splittings seems to be new.

The paper is organized as follows. In Section 1 we discuss properties of ordinary pants decompositions. In Section 2 we introduce double pants decompositions, admissible double pants decompositions and the groupoid generated by flips and twists. We prefer to work with surfaces containing neither holes nor marked points and postpone all details concerning the holes (marked points) till Section 5. In Section 3, we prove transitivity theorem for the case of surfaces of genus $g = 2$ without marked points (Theorem 3). In Section 4, we prove the Main Theorem for the case of surfaces of any genus (Theorem 4.20). In Section 5, we extend basic definitions to the case of surfaces with holes or marked points and complete the proof of Main Theorem. Finally, in Section 6, we prove the Corollary (Theorem 6.3).

Acknowledgments. We are grateful to Allen Hatcher, Saul Schleimer, Anton Zorich, Karen Vogtmann and Ursula Hamenstädt for many helpful conversations. The work was initiated during the authors’ stay at Max Planck Institute for Mathematics and completed during the stay of the first author at the same Institute. We are grateful to the Institute for hospitality, support and a nice working atmosphere. Finally, we are grateful to the anonymous referee for a bunch of valuable suggestions.

1. PANTS DECOMPOSITIONS

1.1. Zipped pants decompositions. Let $S = S_{g,n}$ be an oriented surface of genus g with n or marked points, where $2g + n > 2$. Throughout Sections 1–4 we assume in addition $n = 0$ (this assumption will be removed in Section 5).

A *curve* c on S is an embedded closed non-contractible curve considered up to a homotopy of S .

Given a set of curves we always assume that there are no “unnecessary intersections”, so that if two curves of this set intersect each other in k points then there are no homotopy equivalent pair of curves intersecting in less than k points.

For a pair of curves c_1 and c_2 we denote by $|c_1 \cap c_2|$ the geometric number of intersections (i.e., counting without multiplicities) of c_1 with c_2 .

Definition 1.1 (Pants decomposition). A *pants decomposition* P of S is a system of nonoriented mutually disjoint curves $P = P_S = \{c_1, \dots, c_k\}$ decomposing S into pairs of pants (i.e., into spheres with 3 holes).

It is easy to see that any pants decomposition of a surface of genus g consists of $3g - 3$ curves. Note that we do allow self-folded pants, two of whose boundary components are identified in S as shown in Fig. 1.1.

A surface which consists of one self-folded pair of pants will be called *handle*.

We say that a curve $c \in P$ is *non-regular* if c is contained in a handle \mathfrak{h} such that the boundary c_1 of \mathfrak{h} is also contained in P (see Fig. 1.1). Otherwise, we say that c is *regular*.




FIGURE 1.1. (a) A pair of pants; (b) a pair of self-folded pants composing a handle (the handle contains a non-regular curve c)

Remark 1.2. A set of curves forming a pants decomposition is maximal in sense that any larger set of mutually disjoint non-oriented curves contains homotopy equivalent curves.

Definition 1.3 (Zipper system). A union $Z = \{z_1, \dots, z_{g+1}\}$ of mutually disjoint curves is a *zipper system* if Z decomposes S into two spheres with $g + 1$ holes.

Definition 1.4 (Z compatible with P). A zipper system Z is *compatible with a pants decomposition* $P = \{c_1, \dots, c_{3g-3}\}$ if $|c_i \cap (\bigcup_{j=1}^{g+1} z_j)| = 2$ for each $i = 1, \dots, 3g - 3$.

Fig. 1.2 contains an example of a zipper system Z compatible with a pants decomposition P .




FIGURE 1.2. A zipper system Z compatible with a pants decomposition P

Lemma 1.5. *If Z is a zipper system compatible with a pants decomposition P then $\bigcup_{j=1}^{g+1} z_j$ decomposes each pair of pants in P into two hexagons.*

Proof. Suppose that a curve z_j intersects a curve c_i contained in the boundary of a pair of pants p_1 . The curves of the pants decomposition cut z_j into segments. Let l be a segment of z_j (or one of such segments) contained in p_1 . Since curves do not have unnecessary intersections, l looks as shown in Fig. 1.3(a) or (b). If l looks as in Fig. 1.3(a) then for some of the three boundary curves of p_1 the condition $|c_i \cap (\bigcup_{j=1}^{g+1} z_j)| = 2$ is broken. This implies that it is as one shown in Fig. 1.3(b). Therefore, p_1 looks like in Fig. 1.3(c), i.e., p_1 is decomposed into two hexagons. \square

Remark 1.6. If Z is a zipper system compatible with a pants decomposition P then there exists an orientation reversing involution σ of S such that σ preserves Z pointwise and such that $\sigma(c_i) = c_i$ for each $c_i \in P$. To build this involution one needs only to switch the pairs of hexagons described in Lemma 1.5.




FIGURE 1.3. Zipper system Z decomposes each pair of pants into two hexagons

Definition 1.7 (Zipped flip). Given a pants decomposition $P = \{c_1, \dots, c_{3g-3}\}$ and a zipper system Z compatible with P we define a *zipped flip* of pants decomposition as it is shown in Fig. 1.4. Formally speaking, a *zipped flip* f_i of a pants decomposition $P = \{c_1, \dots, c_{3g-3}\}$ in the curve c_i is a replacing a regular curve $c_i \subset P$ by a unique curve c'_i satisfying the following properties:

- c'_i does not coincide with any of c_1, \dots, c_{3g-3} ;
- c'_i intersects Z exactly in two points;
- $c'_i \cap c_j = \emptyset$ for all $j \neq i$.

Clearly, $f_i(P)$ is a new pants decomposition of S , and Z is a zipper system compatible with $f_i(P)$. The uniqueness of the curve c'_i satisfying the properties in Definition 1.7 verifies trivially. In particular, it is easy to see that $f_i \circ f_i(c_i) = c_i$.




FIGURE 1.4. A zipped flip of a pants decomposition

1.2. Unzipped pants decompositions

Definition 1.8 (Lagrangian plane of pants decomposition). Let $P = \{c_1, \dots, c_{3g-3}\}$ be a pants decomposition. A Lagrangian plane $\mathcal{L}(P) \subset H_1(S, \mathbb{Z})$ is a sublattice spanned by the homology classes $h(c_i)$, $i = 1, \dots, 3g-3$ (here c_i is taken with any orientation).

Notation. By $\langle h(c_1), \dots, h(c_k) \rangle$ we denote the sublattice of $H_1(S, \mathbb{Z})$ spanned by $h(c_1), \dots, h(c_k)$, so that $\mathcal{L}(P) = \langle h(c_1), \dots, h(c_{3g-3}) \rangle$.

Definition 1.9 (Unzipped flip). Let $P = \{c_1, \dots, c_n\}$ be a pants decomposition. Define an *unzipped flip of P in the curve c_i* (or just a *flip*) as a replacing a regular curve $c_i \subset P$ by any curve c'_i satisfying the following properties:

- c'_i does not coincide with any of c_1, \dots, c_n ;
- $|c'_i \cap c_i| = 2$;
- $c'_i \cap c_j = \emptyset$ for all $j \neq i$.

Proposition 1.10. $\mathcal{L}(P) = \mathcal{L}(f(P))$ for any unzipped flip f of P .

Proof. Let $p_1 \cup p_2$ be two pairs of pants glued along a curve e changed under the flip f . Let a, b, c, d be four curves cutting $p_1 \cup p_2$ out of S . Suppose that $f(e)$ separates a and b from c and d in $p_1 \cup p_2$. Denote by $h(c)$ the homology class of c . Then

$$h(f(e)) = h(a) + h(b) = h(c) + h(d) \in \mathcal{L}(P).$$

The similar relation holds when $f(e)$ separates a and c from b and d or a and d from b and c . \square

Lemma 1.11. Let P be a pants decomposition and $c \in P$ be a curve. A flip f of P in the curve c is defined uniquely by a homology class of $f(c)$ up to a Dehn twist along c .

Proof. Consider two pairs of pants p_1 and p_2 adjacent to c , let $M = p_1 \cup p_2$. Since $|(f(c) \cap c)| = 2$ and $f(c) \cap \partial M = \emptyset$, the segment $f(c) \cap p_i$, $i = 1, 2$ looks as shown in Fig. 1.3(a). The homology class of $f(c)$ defines which of the ends of $f(c) \cap p_1$ are glued to which of the ends of $f(c) \cap p_2$. So, the only freedom in gluing of p_1 to p_2 is generated by a Dehn twist along c . \square

The result of Lemma 1.11 may be restated as follows.

Lemma 1.12. *Let P be a pants decomposition compatible with a zipper system Z . Let f_c be a flip of P in a curve $c \in P$. Then f_c is a zipped flip for some $Z' = T_c^m(Z)$, where $m \in \mathbb{Z}$ and T_c is a Dehn twist along c .*

Remark 1.13. In particular, Lemma 1.12 implies that if P_0 is a pants decomposition compatible with a zipper system Z then after any sequence of flips f_1, \dots, f_k we obtain a decomposition P_k compatible with some zipper system Z_k . Moreover, one can choose zipper systems $Z_1, \dots, Z_k, Z_0 = Z$ so that f_i is a zipped flip with respect to Z_i and $Z_{i+1} = T_{c_i}^{m_i}(Z_i)$ for the curve $c_i \in P_i$ changed by f_i .

A Dehn twist along a curve $c \in P$ is a composition of two flips (see Fig. 1.5).




FIGURE 1.5. Dehn twist as a composition of two flips

Definition 1.14 (Category of unzipped pants decompositions). A category $\mathfrak{P}_g(\mathcal{L})$ for the given Lagrangian plane \mathcal{L} is the following:

Objects: pants decompositions P of a genus g surface S satisfying $\mathcal{L}(P) \in \mathcal{L}$;

Elementary morphisms: unzipped flips.

Other *morphisms* are compositions of elementary ones.

Example 1.15 (An orbit of a pants decomposition of S_2). In this example we describe an orbit of an arbitrary pants decomposition of a surface of genus 2. The description is in terms of a graph Γ where a vertex $v_P \in \Gamma$ corresponds to a pants decomposition P and an edge $e \in \Gamma$ connecting v_P to $v_{P'}$ corresponds to a flip f such that $P' = f(P)$.

A pants decomposition P of S_2 may be of one of two types:

- (a) “non-self-folded”: P consists of two non-self-folded pairs of pants (the pants decompositions of this type are denoted by squares in Fig. 1.6);
- (b) “self-folded”: P consists of two handles glued along the holes (the pants decompositions of this type are denoted by circles in Fig. 1.6).

For each of the two types of vertices of Γ we need to understand how many edges are incident to the vertex. In fact, the number of such edges is always infinite: if $c \in P$ is a regular curve, T_c is a Dehn twist along c and $f(c)$ is a flip of the curve c then $T_c(f(c))$ is also a flip of c (see Fig. 1.5). On the other hand, Lemma 1.11 states that modulo Dehn twist T_c there are exactly two possibilities for the flip $f(c)$. Therefore, instead of Γ we will draw the simplified graph $\bar{\Gamma}$ obtained from Γ after factorizing by Dehn twists T_c for each flipped curve c . Then for each regular curve $c \in P$ there are exactly 2 edges emanating from vertex $v_P \in \bar{\Gamma}$. If P is of non-self-folded type, there are 3 regular curves in P , so there are 6 edges incident to $v_P \in \bar{\Gamma}$. If P is of self-folded type, there is a unique regular curve in P , so there are only two edges incident to $v_P \in \bar{\Gamma}$. The graph $\bar{\Gamma}$ is shown in Fig. 1.6.

We will say that a path $\gamma \in \bar{\Gamma}$ is *alternating* if any edge of γ connects two vertices of different types. It is easy to see that for each path $\gamma \in \bar{\Gamma}$ there exists an alternating path $\gamma' \in \bar{\Gamma}$ with the same endpoints. Indeed, each edge of $\bar{\Gamma}$ connecting two vertices of the same type (those are always “square” vertices) may be substituted by an alternating path of two edges. Since a Dehn twist along a curve of pants decompositions is a composition of two flips (each changing the type of a pants decomposition in case of S_2), we obtain the same property for an arbitrary path in Γ :

For each path $\gamma \in \Gamma$ there exists an alternating path $\gamma' \in \Gamma$ with the same endpoints.




FIGURE 1.6. An orbit of a pants decomposition of S_2 (modulo Dehn twists). Vertices marked by squares and circle correspond to non-self-folded and self-folded pants decompositions respectively. Edges correspond to flips. The labels 1, 2, 3 on the edges show which of the three curves of the decomposition is flipped along this edge

Example 1.15 shows that the pants decompositions containing the curves separating handles play special role among other pants decompositions of a closed genus 2 surface.

Definition 1.16 (Standard pants decomposition). A pants decomposition P is *standard* if P contains g curves c_1, \dots, c_g such that c_i cuts out of S a handle \mathfrak{h}_i .

1.3. Pants decompositions and handlebodies. A 3-dimensional *handlebody* is a 3-dimensional disk with several handles attached (where a handle is a solid torus, and it is attached to a 3-disk along a 2-disk).

Given a pants decomposition P one may construct a handlebody in the following way: for each pair of pants P_i one considers a 3-disk D_i with three marked 2-disks on its boundary, then for each pair of adjacent (in P) pairs of pants P_i, P_j one attaches D_i to D_j along the marked disks. One obtains a handlebody $H(P)$ whose boundary coincides with S and carries the structure of the initial pants decomposition P .

The following proposition is evident:

Proposition 1.17. *Let f be a flip of P . Then $H(f(P)) = H(P)$.*

Proposition 1.18 (A. Hatcher [6]). *Morphisms of $\mathfrak{P}_g(\mathcal{L})$ do not act transitively on the objects of the same category.*

Proof. Suppose that the surface S is embedded in \mathbb{R}^3 and a pants decomposition P is such that each curve of P is contractible inside the inner handlebody defined by $S \subset \mathbb{R}^3$. Then each flip preserves this property of P . On the other hand there exists a pants decomposition $P' \in \mathcal{L}(P)$ with non-contractible (inside the given handlebody) curves (see Fig. 1.7 for a non-contractible curve c such that $h(c) \in \mathcal{L}(P)$). \square




FIGURE 1.7. Pants decomposition $P = \{c_1, c_2, c_3\}$ and a curve c such that $h(c) = h(c_2)$ and c is not contractible inside the inner handlebody (the curve c is linked non-trivially with c_3)

Proposition 1.19 (F. Luo [11]). *Flips do act transitively on the pants decompositions whose curves are contractible in a given handlebody of a surface $S \subset \mathbb{R}^3$.*

Corollary 1.20. *Flip-equivalence classes of pants decompositions of a given surface are in one-to-one correspondence with handlebodies defined by pants decompositions.*

2. DOUBLE PANTS DECOMPOSITIONS

2.1. Double pants decompositions and flip-twist groupoid

Definition 2.1 (Lagrangian planes in general position). Two *Lagrangian planes* \mathcal{L}_1 and \mathcal{L}_2 are in *general position* if $H_1(S, \mathbb{Z}) = \langle \mathcal{L}_1, \mathcal{L}_2 \rangle$.

See Fig. 2.1 for an example of two pants decompositions spanning a pair of Lagrangian planes in general position.




FIGURE 2.1. Pair of pants decompositions (P_a, P_b)

Definition 2.2 (Double pants decomposition). A *double pants decomposition* $DP = (P_a, P_b)$ is a pair of pants decompositions P_a and P_b of the same surface such that the Lagrangian planes $\mathcal{L}_a = \mathcal{L}(P_a)$ and $\mathcal{L}_b = \mathcal{L}(P_b)$ spanned by these pants decompositions are in general position.

Definition 2.3 (Handle twists). A *Handle twist* is a transformation of a double pants decomposition $DP = (P_a, P_b)$ which may be performed if P_a and P_b contain the same curve $a_i = b_i$ separating the same handle \mathfrak{h} , see Fig. 2.2. Let $a \in \mathfrak{h}$ and $b \in \mathfrak{h}$ be the only curves of P_a and P_b contained in \mathfrak{h} . Then a *handle twist in \mathfrak{h}* is a Dehn twist along a or along b in any of two directions (see Fig. 2.2(b)).




FIGURE 2.2. Handle twists: (a) handle (double self-folded pair of pants); (b) the same handle after one of the four possible handle twists

Definition 2.4 (Category of double pants decompositions). A category $\mathfrak{DP}_{g,0}$ of *double pants decompositions* of a genus g surface $S = S_{g,0}$ is the following:

Objects: double pants decompositions $DP = (P_a, P_b)$ of S .

Elementary morphisms:

- unzipped flips of P_i ($i \in \{a, b\}$);
- handle twists.

Other *morphisms* are compositions of elementary ones.

Remark 2.5. The index “ $g, 0$ ” in the notation $\mathfrak{DP}_{g,0}$ is to underline that the objects of this category are double pants decompositions of surfaces of genus g *without* marked points.

Definition 2.6 (\mathfrak{DP} -equivalence). Two double pants decompositions are called \mathfrak{DP} -equivalent if there exists a morphism of $\mathfrak{DP}_{g,0}$ taking one of them to the other.

Definition 2.7 (Flip-twist groupoid). All morphisms of $\mathfrak{DP}_{g,0}$ are reversible, so the morphisms form a groupoid acting on the objects of $\mathfrak{DP}_{g,0}$. We will call it a *flip-twist groupoid* and denote FT .

2.2. Admissible double pants decompositions

Definition 2.8 (Standard double pants decomposition, principle curves). A double pants decomposition (P_a, P_b) is *standard* if there exist g curves c_1, \dots, c_g such that the following two conditions hold:

- $c_i \in P_a \cap P_b$;
- c_i cuts out of S a handle \mathfrak{h}_i .

The set of curves $\{c_1, \dots, c_g\}$ in this case is the *set of principle curves* of (P_a, P_b) .

See Fig. 2.3 for an example of a standard double pants decomposition. Notice that a standard double pants decomposition is not unique: many elements of the mapping class group of S act non-trivially on the decomposition shown in Fig. 2.3, moreover applying a flip to the curve in a middle of P_a one may obtain a decomposition which is combinatorially different from one shown in the figure (in the former there are 5 curve contained in $P_a \cap P_b$, while in the latter there are 4 ones).




FIGURE 2.3. Standard double pants decomposition (P_a, P_b) . The principle curves are shown by bold lines

Definition 2.9 (Admissible decomposition). A double pants decomposition (P_a, P_b) is *admissible* if there exists a standard double pants decomposition (P'_a, P'_b) such that P'_a is flip-equivalent to P_a and P'_b is flip-equivalent to P_b .

Example 2.10. It is easy to check that a double pants decomposition (P_a, P_b) shown in Fig. 2.1 is admissible.

Lemma 2.11. Let DP be an admissible double pants decomposition and t be a handle twist of DP . Then $t(DP)$ is also admissible.

Proof. Denote by \bar{t} the Dehn twist of S that acts on DP as the handle twist t . Notice that applying \bar{t} to a double pants decomposition we obtain the decomposition with the same combinatorial properties, in particular, \bar{t} takes flippable curves to flippable. Moreover, for each double pants decomposition DP_0 if f is a flip of a curve $c \in DP_0$ then $f' = \bar{t}f\bar{t}^{-1}$ is a flip of the curve $\bar{t}(c) \in \bar{t}(DP_0)$ and $\bar{t}f\bar{t}^{-1}(\bar{t}(DP_0)) = \bar{t}(f(DP_0))$.

Suppose now that f_1, \dots, f_k is a sequence of flips taking DP to a standard decomposition DP' . Then the sequence $f'_1, \dots, f'_k, f'_i = \bar{t}f_i\bar{t}^{-1}$ of flips takes $t(DP)$ to a standard decomposition $\bar{t}(DP')$. \square

The set of admissible double pants decompositions is closed under the action of flips and handle twists, so we may define a subcategory of $\mathfrak{DP}_{g,0}$:

Definition 2.12 (Category of admissible double pants decompositions). A category $\mathfrak{ADP}_{g,0}$ of *admissible double pants decompositions* of a genus g surface is the following:

Objects: admissible double pants decompositions $DP = (P_a, P_b)$ of a genus g surface.

Elementary morphisms:

- unzipped flips of P_i ($i \in \{a, b\}$);
- handle twists.

Other *morphisms* are compositions of elementary ones.

2.3. Admissible decompositions as Heegaard splittings of \mathbb{S}^3 . Given a pants decomposition P_a on S , one can define a handlebody S_+ such that all curves of P_a are contractible in S_+ . A choice of two pants decompositions P_a and P_b gives two handlebodies which could be attached along S , so that we obtain a Heegaard splitting of a 3-manifold denoted by $M(P_a, P_b)$.

Lemma 2.13. *Suppose that (P_a, P_b) is a standard double pants decomposition. Then $M(P_a, P_b) = \mathbb{S}^3$.*

Proof. We consider the surface $S = S_{g,0}$ with a standard pants decomposition as a union of g handles and a g -holed sphere. Notice that on each of these surfaces the homology classes of the curves $c \in P_a \cup P_b$ generate entire homology lattice.

First, we consider a handle (a one holed torus) \mathfrak{h}_i with double pants decomposition $DP = (P_a, P_b) \cap \mathfrak{h}_i$. In this case DP consists only of two curves a_i and b_i . Since $\langle h(a_i), h(b_i) \rangle = H_1(\mathfrak{h}, \mathbb{Z})$, a_i intersects b_i at a unique point. It is easy to see that \mathfrak{h} may be embedded in \mathbb{S}^3 so that a_i is contractible inside the inner solid torus (bounded by \mathfrak{h} and a disk attached to $\partial\mathfrak{h}$) and b_i is contractible outside one: to see that first we embed a_i correctly and then apply several twists along a_i to “unwrap” b_i .

Now we embed each handle $\mathfrak{h}_1, \dots, \mathfrak{h}_g$ in the same \mathbb{S}^3 so that all images are disjoint. Then we attach the sphere $S_{0,g}$. Since all curves on $S_{0,g}$ are homologically trivial, they are contractible both inside and outside. So, we constructed an embedding of S to \mathbb{S}^3 such that all curves of P_a and P_b are contractible in inner and outer handlebodies respectively, which is equivalent to $M(P_a, P_b) = \mathbb{S}^3$. \square

Lemma 2.14. *If $M(P_a, P_b) = \mathbb{S}^3$ then (P_a, P_b) is admissible.*

Proof. If $M(P_a, P_b) = \mathbb{S}^3$ then there exists an embedding $S \hookrightarrow \mathbb{S}^3$ such that the curves of P_a are contractible in inner handlebody for this embedding, while the curves of P_b are contractible in the outer one. It is easy to find a standard double pants decomposition (P'_a, P'_b) such that the curves of P'_a and the curves of P'_b are contractible in the inner and outer handlebodies respectively. In view of Corollary 1.20 flips act transitively on pants decompositions contractible inside a given handlebody, so the decomposition (P_a, P_b) is flip-equivalent to (P'_a, P'_b) , which is standard. So, (P_a, P_b) is admissible. \square

Combining Lemma 2.13 with Lemma 2.14 we arrive at the following characterization of admissible decompositions (it also may be taken as an alternative definition):

Theorem 2.15. *A double pants decomposition (P_a, P_b) is admissible if and only if $M(P_a, P_b) = \mathbb{S}^3$.*

Our next aim is to prove that morphisms of $\mathcal{ADP}_{g,0}$ act transitively on the objects of $\mathcal{ADP}_{g,0}$. In other words, we will prove that flips and twists act transitively

on double pants decompositions corresponding to Heegaard splittings of S^3 . This is done in Section 3 for the case of $g = 2$ and in Section 4 for a general case.

Remark 2.16. An admissible double pants decomposition can be also defined as any decomposition obtained from a standard decomposition via a sequence of flips and handle twists. In principle, this class of decompositions may be wider than one in Definition 2.9 (where only flips are allowed), so the transitivity theorem for this class looks stronger than one we are going to prove. In fact, Lemma 2.11 shows that these two classes coincide.

3. TRANSITIVITY OF MORPHISMS IN CASE $g = 2$

In this section we prove the Main Theorem for the case of surface of genus $g = 2$ containing no marked points. The proof is based on the following result 3.2 of Hatcher and Thurston [7].

Definition 3.1 (\mathcal{S} -moves). Let P be a pants decomposition of S and $a, c \in P$ be two curves such that c cuts out of S a handle \mathfrak{h} and let $a \subset \mathfrak{h}$. An \mathcal{S} -move of a pants decomposition P in a curve a is a substitution of a by a curve a' , where $a' \in \mathfrak{h}$ is an arbitrary curve such that $|a \cap a'| = 1$.

Theorem 3.2 (A. Hatcher, W. Thurston [7], [5]). *Let $S_{g,n}$ be a surface of genus g with n holes. Any pants decomposition of $S_{g,n}$ can be transformed to any other pants decomposition of $S_{g,n}$ via flips and \mathcal{S} -moves.*

Remark 3.3. In the initial paper of Hatcher and Thurston [7] the surface $S_{g,n}$ is supposed to be closed surface containing no marked points. This assumption is removed in [5].

Remark 3.4. Theorem 3.2 does not imply immediately transitivity of morphisms in $\mathfrak{AP}_{g,0}$ (Theorem 4.20) since the set of handle twists in $\mathfrak{AP}_{g,0}$ is much smaller than the set of \mathcal{S} -moves in Theorem 3.2 (the former depends on the relative position of two decompositions).


Now we will prove several lemmas: Lemmas 3.6 and 3.8 will be used for the proof of transitivity both in case of $g = 2$ and in general case. Lemma 3.9 is specific for genus 2, its generalization requires more work for general genus.

Definition 3.5 (Double \mathcal{S} -move). Under the conditions of definition of a handle twist (Definition 2.3), a *double \mathcal{S} -move* in \mathfrak{h} is the move switching the curves a and b .

Lemma 3.6. *Double \mathcal{S} -move is a morphism of $\mathfrak{AP}_{g,0}$.*

Proof. Any double \mathcal{S} -move is a composition of 3 handle twists, see Fig. 3.1. □

Lemma 3.7. *Let \mathfrak{h} be a handle and $c \in \mathfrak{h}$ be a closed non-self-intersecting curve with non-trivial homology class $h(c)$. Then the homotopy class of c is completely determined by $h(c)$.*

FIGURE 3.1. Double \mathcal{S} -move as a composition of three handle twists

Proof. Attach a disk D to the boundary $\partial\mathfrak{h}$ so that we obtain a closed torus $\mathfrak{t} = \mathfrak{h} \cup D$. The statement of the lemma holds for a closed torus, so the homotopy class of c is determined by $h(c)$ together with the relative position of the curve c and the hole $\partial\mathfrak{h}$ in the closed torus \mathfrak{t} . However, $\mathfrak{t} \setminus c$ is connected so the relative position of c and $\partial\mathfrak{h}$ is unique up to isotopy of \mathfrak{t} , which implies that the homotopy class of c is determined by $h(c)$. \square

Lemma 3.8. *Let (P_a, P_b) and (P'_a, P'_b) be two standard double pants decompositions containing the same handle \mathfrak{h} . Then $(P_a, P_b)|_{\mathfrak{h}}$ may be transformed to $(P'_a, P'_b)|_{\mathfrak{h}}$ by a sequence of handle twists in \mathfrak{h} (where $(P_1, P_2)|_{\mathfrak{h}}$ is a restriction of the double pants decomposition to the handle \mathfrak{h}).*

Proof. Let a, b, a', b' be the curves of P_a, P_b, P'_a, P'_b contained in \mathfrak{h} . We need to find a composition $\bar{\psi}$ of handle twists in \mathfrak{h} such that $\bar{\psi}(a) = a', \bar{\psi}(b) = b'$.

First, suppose that $a = a'$. Since $\langle h(a), h(b) \rangle = \langle h(a), h(b') \rangle$, one has

$$h(b') = h(b) + l_a h(a)$$

for $l_a \in \mathbb{Z}$. Applying l_a -tuple twist along a to the curve b we obtain a curve c such that $h(c) = h(b')$. In view of Lemma 3.7 this implies that $b' = c$. So, in the case $a = a'$, $\bar{\psi}$ is a composition of flips along a .

Next, suppose that $a' = b$. Then a double \mathcal{S} -move interchanging a with b reduces the question to the previous case.

Now, suppose that $a' \neq a, b$. Then we have

$$h(a') = l_a h(a) + l_b h(b),$$

where $l_a, l_b \in \mathbb{Z}$ are coprime. In view of Lemma 3.7, we only need to find a sequence $\bar{\psi}$ of morphisms of $\mathfrak{DP}_{g,0}$ taking a to any curve $x \in S'$ such that $h(x) = l_a h(a) + l_b h(b)$. Since l_a and l_b are coprime and a handle twist transforms $(h(a), h(b))$ into either $(h(a) \pm h(b), h(b))$ or $(h(a), h(b) \pm h(a))$, this sequence of handle twists do exists. \square

Lemma 3.9. *Let (P_a, P_b) be a standard double pants decomposition of $S_{2,0}$. Let $\varphi = \varphi_k \circ \dots \circ \varphi_1$ be a sequence of flips of P_a such that $\varphi(P_a)$ is a standard decomposition. Then there exists a morphism η of $\mathfrak{ADP}_{2,0}$ and a pants decomposition P'_b of S such that $\eta((P_a, P_b))$ is a standard double pants decomposition and $\eta((P_a, P_b)) = (\varphi(P_a), P'_b)$.*

Proof. Recall from Example 1.15 that for each two pants decompositions P_1 and P_2 connected by a sequence of flips there exists a sequence of flips connecting these pants decompositions and such that each flip in this sequence changes the type of

pants decomposition from “self-folded” into “non-self-folded” or back (in the other word takes a standard pants decomposition to a non-standard one and back). This implies that it is sufficient to show the lemma for compositions $\varphi = \varphi_2 \circ \varphi_1$ of two flips.

If φ_2 changes the same curve as φ_1 does, then φ is a twist and the required composition η is shown in Fig. 3.2.




FIGURE 3.2. (a) Twist $\varphi(P_a)$: $\varphi = \varphi_2 \circ \varphi_1$; (b) Composition $\eta(P_a, P_b)$ for the twist φ

If the flips φ_1 and φ_2 change different curves then (modulo twists) φ looks like in Fig. 3.3(a) and the required composition η is shown in Fig. 3.3(b).

This completes the proof of Lemma 3.9. \square

Theorem 3.10. *Morphisms of $\mathfrak{ADP}_{2,0}$ act transitively on the objects of $\mathfrak{ADP}_{2,0}$.*

Proof. By Definition 2.4, the objects of $\mathfrak{ADP}_{2,0}$ are admissible double pants decompositions, so, it is sufficient to prove the transitivity on the set of standard pants decompositions.




FIGURE 3.3. (a) Composition $\varphi(P_a)$ of two flips $\varphi = \varphi_2 \circ \varphi_1$;
 (b) Composition $\eta = \eta((P_a, P_b))$ for φ : $\eta(P_a) = \varphi(P_a)$, $\eta(P_b)$ is shown

Let (P_a, P_b) and (P'_a, P'_b) be two standard double pants decompositions. If the principle curve of (P_a, P_b) coincides with one of (P'_a, P'_b) then Lemma 3.8 implies that (P_a, P_b) is \mathfrak{DP} -equivalent to (P'_a, P'_b) .

Suppose that the principle curves of (P_a, P_b) and (P'_a, P'_b) are different. It is left to show that there exists a sequence $\eta = \eta_n \circ \cdots \circ \eta_1$ of morphisms of $\mathfrak{ADP}_{2,0}$ such that $\eta((P_a, P_b)) = (P'_a, P'_b)$ and P'_b is arbitrary pants decomposition turning the pair (P'_a, P'_b) into a standard pants decomposition.

By Theorem 3.2, there exists a sequence $\psi = \psi_k \circ \cdots \circ \psi_1$ of flips and \mathcal{S} -moves taking P_a to P'_a . By definition, in case of $g = 2$ an \mathcal{S} -move is applicable only to standard double pants decompositions. This implies that the sequence ψ is a composition of several subsequences of two types:

- subsequences of flips, each subsequence takes a standard pants decomposition to another standard one;
- \mathcal{S} -moves.

By Lemma 3.9, a subsequence of the first type may be extended to a morphism of $\mathfrak{ADP}_{2,0}$ taking a standard pants decomposition to another a standard one. By Lemma 3.8 any \mathcal{S} -move of the component P_a may be realized as a sequence of morphisms of $\mathfrak{ADP}_{2,0}$ taking a standard double pants decomposition to a standard one. This implies that ψ may be extended to a morphism of $\mathfrak{ADP}_{2,0}$ and the theorem is proved. \square

4. TRANSITIVITY OF MORPHISMS IN CASE OF SURFACES WITHOUT MARKED POINTS

In this section we adjust the proof of Theorem 3.10 to the case of higher genus.

4.1. Preparatory lemmas

Lemma 4.1. *Let P be a pants decomposition and \mathfrak{h} be a handle cut out by some $c \in P$. Let φ be an \mathcal{S} -move in \mathfrak{h} . Then $\varphi = (\bar{\varphi}_1)^{-1} \circ \varphi_2 \circ \bar{\varphi}_1$, where $\bar{\varphi}_1$ is a sequence of flips preserving \mathfrak{h} and φ_2 is an \mathcal{S} -move of a standard pants decomposition.*

Proof. Consider the surface $S \setminus \mathfrak{h}$. By Theorem 3.2 it is possible to transform any pants decomposition of $S \setminus \mathfrak{h}$ to a standard one (clearly it may be done using flips only: one may use zipped flips with respect to any zipper system compatible with P). This defines the sequence $\bar{\varphi}_1$. Then we apply \mathcal{S} -move in the handle \mathfrak{h} and apply φ_1^{-1} to bring the pants decomposition of $S \setminus \mathfrak{h}$ into initial position. \square

In the proof of Theorem 3.10 we used the fact that in case of $g = 2$ \mathcal{S} -moves are defined for standard decompositions only. This does not hold for $g > 2$. However, in view of Lemma 4.1 the following holds.

Lemma 4.2. *Let P_a and P'_a be standard pants decompositions and there exists a sequence $\varphi = \varphi_k \circ \dots \circ \varphi_1$ of flips and \mathcal{S} -moves such that $\varphi(P_a) = P'_a$. Then it is possible to choose φ in such a way that all \mathcal{S} -moves in φ are applied to standard decompositions.*

Lemma 4.3. *Let $S_{0,g}$ be a sphere with g holes. Then any pants decomposition of $S_{0,g}$ may be transformed to any other pants decomposition of $S_{0,g}$ by a sequence of flips.*

Proof. The statement follows immediately from Theorem 3.2 since \mathcal{S} -moves could not be applied to $S_{0,g}$. \square

4.2. Transitivity in case of the same zipper system

Definition 4.4 ((P_a, P_b) compatible with Z). One says that a double pants decomposition (P_a, P_b) is *compatible* with a zipper system Z if P_a is compatible with Z and P_b is compatible with Z .

In this section we will prove that if (P_a, P_b) and (P'_a, P'_b) are two admissible double pants decompositions compatible with the same zipper system then (P_a, P_b) is \mathfrak{DP} -equivalent to (P'_a, P'_b) .

Definition 4.5 (Strictly standard decomposition). An admissible double pants decomposition (P_a, P_b) is *strictly standard* if it is standard and $c \in \{P_a \cup P_b\} \setminus \{P_a \cap P_b\}$ if and only if c is contained inside some handle cut out by a principle curve of the decomposition.

Example 4.6. The double pants decomposition in Fig. 2.3 is standard but not strictly standard.

Remark 4.7. Although we will not need this below, one can mention that strictly standard decompositions can be also characterized by any of the following equivalent minimal properties:

- double pants decompositions with minimal possible number of distinct curves (i.e., with $4g - 3$ curves for $S_{g,0}$);
- double pants decompositions with minimal possible number of intersections of curves (i.e., with g intersections).

To prove transitivity of morphisms of $\mathfrak{ADP}_{g,0}$ on the objects of $\mathfrak{ADP}_{g,0}$ it is sufficient to prove \mathfrak{DP} -equivalence of all standard double pants decompositions (since the objects of $\mathfrak{ADP}_{g,0}$ are admissible ones). Furthermore, any standard double pants decomposition is \mathfrak{DP} -equivalent to some strictly standard one in view of Lemma 4.3. So, it is sufficient to prove the transitivity of morphisms of $\mathfrak{ADP}_{g,0}$ on strictly standard double pants decompositions.

To prove the transitivity of morphisms of $\mathfrak{ADP}_{g,0}$ on strictly standard double pants decompositions we do the following:

- we show that each strictly standard double pants decomposition is compatible with some zipper system (see Proposition 4.8);
- we prove that morphisms of $\mathfrak{ADP}_{g,0}$ act transitively on strictly standard double pants decompositions compatible with a given zipper system (see Lemma 4.16);
- Finally, we show that for two different zipper systems Z and Z' we may find a sequence of zipper systems $Z = Z_1, Z_2, \dots, Z_k = Z'$ in which Z_i differs from Z_{i+1} by a twist along some curve c , $|c \cap Z_i| = 2$. We show (Lemma 4.17) that in this case morphisms of $\mathfrak{ADP}_{g,0}$ are sufficient to change Z_i to Z_{i+1} .

Proposition 4.8. *For any strictly standard double pants decomposition (P_a, P_b) there exists a zipper system Z compatible with (P_a, P_b) .*

Proof. An intersection of the required zipper system $Z = \{z_0, z_1, \dots, z_g\}$ with a handle looks as shown in Fig. 4.1.(a): if a_i and b_i are curves of P_a and P_b contained in a given handle \mathfrak{h}_i , $i = 1, \dots, g$, then \mathfrak{h}_i contains a zipper z_i such that $|z_i \cap a_1| = 1$ and $|z_i \cap b_1| = 1$. The curve z_0 visits each of the handles and goes in \mathfrak{h}_i along z_i .

The condition that (P_a, P_b) is strictly standard leads to the existence of appropriate z_0 outside of handles: to show this we build a *dual graph* of (P_a, P_b) substituting each pair of pants by an Y-shaped figure and each handle by a point (see Fig. 4.1.(b)). Since (P_a, P_b) is strictly standard we obtain a tree. We consider this tree as a graph drawn on a plane Π . Denote by $\tilde{z}_0 \in \Pi$ the curve doing around the tree. Then z_0 is built as any curve on S which projects to \tilde{z}_0 (more precisely, \tilde{z}_0 determines the order in which z_0 visits the handles of (P_a, P_b)). \square




FIGURE 4.1. A zipper system for a given standard double pants decomposition: (a) behavior in a handle; (b) outside of the handles: a dual graph of a strictly standard pants decomposition

Proposition 4.9. *Let (P_a, P_b) be a standard double pants decomposition and Z be a zipper system compatible with (P_a, P_b) . Then*

- *each of the handles of (P_a, P_b) contains exactly one curve of Z ,*
- *if $z_0 \in Z$ does not belong entirely to any handle then z_0 visits each of g handles exactly once.*

Proof. Let c be a principle curve separating a handle \mathfrak{h} in (P_a, P_b) . Then each curve on S intersects c even number of times. By definition of a zipper system compatible with a pants decomposition, c is intersected by exactly one of the curves z_i . To prove that this curve is the same for all handles in (P_a, P_b) , notice that a pairs of pants dissected along a connected curve does not turn into a union of simply-connected components, which implies that there is a curve $z_j \in Z$, $z_j \neq z_i$ which intersects the handle \mathfrak{h} . Since $z_j \cap c = \emptyset$, z_j is contained in \mathfrak{h} . Similarly, each of the other handles contains entirely some of the curves from Z . No of these curves may coincide with z_i (since z_i intersects \mathfrak{h}). Hence, Z consists of g curves contained in the handles of (P_a, P_b) and one additional curve z_i . This proves the first statement of the proposition. The second statement follows from the fact that each principle curve in (P_a, P_b) should be intersected by some of z_i . \square

Definition 4.10 (Principle zipper, cyclic order). Let (P_a, P_b) be a standard double pants decomposition and $Z = \{z_0, z_1, \dots, z_g\}$ be a zipper system compatible with (P_a, P_b) . Suppose that z_0 is the curve visiting all handles of (P_a, P_b) . Then we call z_0 a *principle zipper*.

A *cyclic order* of Z is $[z_1, z_2, \dots, z_g]$ if an orientation of z_0 goes from \mathfrak{h}_i to \mathfrak{h}_{i+1} , where \mathfrak{h}_i is the handle containing z_i and i is considered modulo $g+1$ (more precisely, Z decomposes S into two $(g+1)$ -holed spheres S_+ and S_- , so that we may choose a positive orientation of z_0 as one which goes in positive direction around S_+ ; for a definition of a cyclic order we choose the positive orientation of z_0).

Given two standard double pants decompositions DP and DP' compatible with the same zipper system Z we say that they have *the same cyclic order* if the cyclic order for DP determined by a positive orientation of z_0 coincides with the cyclic order for DP' determined by the same orientation of z_0 .

Remark 4.11. The definition of cyclic order depends on the choice of S_+ among two subsurfaces. However, the definition of *same cyclic order* of two standard double pants decompositions is independent of this choice (provided that the choice of S_+ is the same for both decompositions).

Proposition 4.12. *Let (P_a, P_b) and (P'_a, P'_b) be two standard pants decompositions compatible with the same zipper system $Z = \{z_0, z_1, \dots, z_g\}$. Suppose that z_0 is the principle zipper of Z both for (P_a, P_b) and (P'_a, P'_b) and the cyclic order of Z is $[z_1, z_2, \dots, z_n]$ both for (P_a, P_b) and (P'_a, P'_b) .*

Then the set of principle curves of (P_a, P_b) coincides with the set of principle curves of (P'_a, P'_b) .

Proof. Let c_1, \dots, c_g be the principle curves of (P_a, P_b) and c'_1, \dots, c'_g be the principle curves of (P'_a, P'_b) . Since the cyclic order of Z is the same for both double pants decompositions, we may assume that $z_0 \cap c_i = z_0 \cap c'_i$. Let S^+ and S^- be

the connected components of $S \setminus Z$. Then $c_i \cap S^+$ separates from S^+ an annulus containing z_i as a boundary component. Clearly, the same holds for $c_i \cap S^-$ as well as for $c'_i \cap S^+$ and $c'_i \cap S^-$. This implies that c_i is homotopy equivalent to c'_i (more precisely, there exists an isotopy of c_i to c'_i with the fixed points $c_i \cap z_0 = c'_i \cap z_0$). \square

Corollary 4.13. *In assumptions of Proposition 4.12, (P_a, P_b) may be transformed to (P'_a, P'_b) by morphisms of $\mathfrak{ADP}_{g,0}$ preserving the principle curves of the standard pants decomposition.*

Proof. This follows from Proposition 4.12, Lemma 3.8 and Lemma 4.3. \square

Corollary 4.13 implies that a standard pants decomposition is determined (modulo action of morphisms of $\mathfrak{ADP}_{g,0}$) by the set of principle curves. Thus, it makes sense to consider a set of principle curves itself, independently of remaining curves in the corresponding double pants decomposition.

Definition 4.14. A zipper system Z is *compatible with a set of principle curves* if $Z = \{z_0, z_1, \dots, z_g\}$, where z_0 visits each handle exactly once and each of the handles contain exactly one of z_i , $i = 1, \dots, g$.

In other words, Z is compatible with a set of principle curves if and only if it is compatible with a standard pants decomposition containing this set of principle curves.

Proposition 4.15. *Let (P_a, P_b) and (P'_a, P'_b) be two standard pants decompositions compatible with the same zipper system $Z = \{z_0, z_1, \dots, z_g\}$. Suppose that z_0 is the principle zipper of Z both for (P_a, P_b) and (P'_a, P'_b) . Then (P_a, P_b) is \mathfrak{DP} -equivalent to (P'_a, P'_b) .*

Proof. By Proposition 4.12 together with Corollary 4.13 the proposition is trivial unless the cyclic order of Z is different for the cases of (P_a, P_b) and (P'_a, P'_b) . It is shown in Fig. 4.2 that the transposition of two neighboring zippers z_i and z_{i+1} in the cyclic order of Z may be realized by morphisms of $\mathfrak{ADP}_{g,0}$.

In more details, in Fig. 4.2, upper row, left, we show (a part of) a zipper system Z compatible with a set of principle curves. In Fig. 4.2, lower row, left we show a strictly standard double pants decomposition $(P_a^{(0)}, P_b^{(0)})$ compatible with Z and having principle curves as in the figure above. Applying handle-twists in curves of P_a (one for each handle) we obtain the decomposition $(P_a^{(1)}, P_b^{(1)})$ (we draw only the front, “visible” part of the decomposition, the non-visible part completely repeats it). Applying two flips (one for P_a and one for P_b) we obtain a double pants decomposition $(P_a^{(2)}, P_b^{(2)})$, and then after one more flip we obtain a standard double pants decomposition $(P_a^{(3)}, P_b^{(3)})$. Choosing appropriate curves in the handles of $(P_a^{(3)}, P_b^{(3)})$ we turn it into a standard double pants decomposition $(P_a^{(4)}, P_b^{(4)})$ compatible with Z (this is done in the same way as the transformation from $(P_a^{(0)}, P_b^{(0)})$ to $(P_a^{(1)}, P_b^{(1)})$, i.e., using handle twists in each of the handles). In Fig. 4.2, upper row, right, we show the zipper system Z together with the principle curves of $(P_a^{(4)}, P_b^{(4)})$. Notice that the cyclic order in Z is changed: in the initial

decomposition (an orientation of) the principle zipper z_0 visits first the handle containing z_1 and then the handle containing z_2 , while in the final decomposition the same orientation of z_0 visits the handles in reverse order. So, the transposition of two adjacent (in the cyclic order) handles is realizable by the morphisms of $\mathfrak{A}\mathfrak{D}\mathfrak{P}_{g,0}$.

Since the permutation group is generated by transpositions of adjacent elements, the proposition is proved. \square





FIGURE 4.2. Transposition in a cyclic order is realizable by morphisms of $\mathfrak{A}\mathfrak{D}\mathfrak{P}_{g,0}$: apply handle-twists in each handle, then a flip in each pants decomposition, then a flip in P_b , and again, handle twists in each of two handles

Lemma 4.16. *Let (P_a, P_b) and (P'_a, P'_b) be two standard pants decompositions compatible with the same zipper system Z . Then (P_a, P_b) is $\mathfrak{D}\mathfrak{P}$ -equivalent to (P'_a, P'_b) .*

Proof. Suppose that the principle zipper of Z is different for (P_a, P_b) and (P'_a, P'_b) (otherwise there is nothing to prove in view of Proposition 4.15).

By Proposition 4.15, the lemma is trivial if the principle zipper of Z is the same for (P_a, P_b) and (P'_a, P'_b) . So, we only need to show that the morphisms of $\mathfrak{D}\mathfrak{P}_{g,0}$ allow to change the principle zipper in Z . We will show that it may be done by flips only.

Let Z be a zipper system (see Fig. 4.3) compatible with a set \bar{c} of principle curves. Let (P_a, P_b) be a standard double pants decomposition containing this set of principle curves. Let \bar{c}' be another set of principle curves compatible with Z and such that z_0 is not a principle zipper (see Fig. 4.3, down). We choose a standard double pants decomposition (P'_a, P'_b) with a set of principle curves \bar{c}' , see Fig. 4.3 (to keep the figure readable we do not draw the curves of (P'_a, P'_b) decomposing $S \setminus \cap_{i=1}^g \mathfrak{h}_g$). To prove the lemma it is sufficient to show that P_a is flip-equivalent to P'_a and P_b is flip-equivalent to P'_b .

FIGURE 4.3. Principle zipper may be changed by morphisms of $\mathfrak{AP}_{g,0}$

The fact that P_a is flip-equivalent to P'_a follows from Proposition 1.19, indeed both P_a and P'_a are contractible in the inner handlebody in the embedding shown in Fig. 4.3. Similarly, P_b is flip-equivalent to P'_b , since both of them are contractible in the outer handlebody, so the lemma is proved. \square

4.3. Proof of transitivity in general case

Lemma 4.17. *Let Z be a zipper system, let σ be an involution preserving Z point-wise and let c be a curve satisfying $|Z \cap c| = 2$, $\sigma(c) = c$. Denote by T_c a Dehn twist along c . Then there exist standard pants decompositions (P_a, P_b) and (P'_a, P'_b) compatible with Z and $Z' = T_c(Z)$ respectively and \mathfrak{AP} -equivalent to each other.*

Proof. We will construct the decomposition (P_a, P_b) so that each curve in it is preserved by σ . For simplicity we will draw only one of two connected components S_+ and S_- of $S \setminus Z$, this is sufficient in view of symmetry. We will draw S_+ as a disk with g holes and an outer boundary coming from the principle zipper.

Since $|Z \cap c| = 2$, the curve c either intersects twice the same curve $z_0 \in Z$ or have single intersections with two distinct curves $z_1, z_2 \in Z$. Consider these two cases.

Suppose that $|c \cap z_0| = 2$ and $z_0 \in Z$. We will build (P_a, P_b) so that z_0 will be a principle zipper. Since $\sigma(c) = c$, the curve c decomposes S_+ into two parts, as in Fig. 4.4(a) (each part containing at least one hole, otherwise the intersection $c \cap Z$ is not essential). We build a strictly standard double pants decomposition (P_a, P_b) containing c , compatible with Z and such that z_0 is a principle zipper (see Fig. 4.4(b) for the construction: for each of the holes we draw a segment of the principle curves encircling this hole). We take the second copy of the same pattern as S_- and glue to S_+ along the boundary. Clearly, we obtain a set of principle curves for a standard double pants decomposition (P_a, P_b) compatible

with Z . It is easy to see that the same decomposition (P_a, P_b) is also compatible with $Z' = T_c(Z)$.




FIGURE 4.4. Case $|c \cap z_0| = 2$: (a) c decomposes S_+ ; (b) strictly standard double pants decomposition (only principle curves and c are shown)

Suppose that $|c \cap z_1| = |c \cap z_2| = 1$, $z_1, z_2 \in Z$ (see Fig. 4.5(a)). Choose (P_a, P_b) so that z_1 and z_2 are not principle zippers and z_1 and z_2 are two neighboring curves in the cyclic order (see Fig. 4.5(b)). Fig. 4.6 contains a sequence of morphisms of $\mathfrak{DP}_{g,0}$ taking (P_a, P_b) to a standard double pants decomposition compatible with $Z' = T_c(Z)$. \square




FIGURE 4.5. Case $|c \cap z_1| = |c \cap z_2| = 1$: (a) c in S_+ ; (b) strictly standard double pants decomposition (only principle curves and c are shown)

Applying Lemma 4.17 we obtain the following corollary.

Corollary 4.18. *In assumptions of Lemma 4.17, the statement of Lemma 4.17 holds for Z and $Z'' = T_c^m(Z)$ for any positive integer degree m .*

Definition 4.19 (\mathfrak{DP} -equivalent standard pants decompositions). A standard pants decomposition P_a is \mathfrak{DP} -equivalent to a standard pants decomposition P'_a if there exist standard pants decompositions P_b and P'_b such that (P_a, P_b) and (P'_a, P'_b) are standard and \mathfrak{DP} -equivalent to each other.




FIGURE 4.6. Twist T_c along c as a composition of elementary morphisms of $\mathfrak{DP}_{g,0}$: the steps labeled by “ S ” are double \mathcal{S} -moves, other steps are compositions of two flips, one in P_a another in P_b . The final figure coincides with the initial one twisted around c

Theorem 4.20. *Let $S = S_{g,0}$ a surface without marked points. Then morphisms of $\mathfrak{ADP}_{g,0}$ act transitively on the objects of $\mathfrak{ADP}_{g,0}$.*

Proof. It is clear from the definition of admissible pants decomposition that it is sufficient to prove transitivity for standard pants decompositions only. By Lemmas 3.8 and 4.3 morphisms of $\mathfrak{DP}_{g,0}$ act transitively on standard pants decompositions with the same principle curves. This implies that it is sufficient to show that any two standard pants decompositions P_a and P'_a are \mathfrak{DP} -equivalent.

By Theorem 3.2 there exists a sequence $\{\varphi_i\}$ of flips and \mathcal{S} -moves taking P_a to P'_a . In view of Lemma 4.1 we may assume that in this sequence \mathcal{S} -moves are applied only to the standard double pants decompositions. Lemma 3.8 treats the \mathcal{S} -moves in standard pants decompositions, thus, we may assume that $\{\varphi_i\}$ consists entirely of flips of P_a .

If Z is a zipper system compatible with (P_a, P_b) and all flips in $\{\varphi_i\}$ are zipped flips (with respect to Z) then there is nothing to prove. Our idea is to decompose the sequence $\{\varphi_i\}$ into several subsequences

$$\{\varphi_i\} = \{\{\varphi_i\}_1, \dots, \{\varphi_i\}_k\}$$

such that in j -th subsequence all flips are zipped flips with respect to the same zipper system Z_j . Denote by P_a^j the pants decomposition obtained from P_a after application of the first j subsequences of flips, $P_a^0 = P_a$, $P_a^k = P'_a$. Clearly, P_a^j is compatible both with Z_j and Z_{j+1} . So, we may use zipped flips (with respect to Z_j) to transform P_a^j to some standard pants decomposition $P_a^j(Z_j)$ compatible with Z_j . Similarly, we may transform P_a^j to some standard pants decomposition $P_a^j(Z_{j+1})$ compatible with Z_{j+1} (see Fig. 4.7).




FIGURE 4.7. Proof of transitivity for $g > 2$

Now, for proving \mathfrak{DP} -equivalence of P_a and P'_a we only need to show two facts:

- (1) $P_a^{j-1}(Z_j)$ is \mathfrak{DP} -equivalent to $P_a^j(Z_j)$ for $0 < j \leq k$;
- (2) $P_a^j(Z)$ is \mathfrak{DP} -equivalent to $P_a^j(Z_{j+1})$ for $0 < j < k$.

The first of this facts follows immediately from Lemma 4.16 since both $P_a^{j-1}(Z_j)$ and $P_a^j(Z_j)$ are compatible with the same zipper system Z_j .

So, we are left to prove the second fact. By Lemma 1.13, we may assume $Z_{j+1} = T_c^m(Z_j)$, where T_c is a Dehn twist along a curve $c \in P_a^j$. By Remark 1.6, there exists an involution σ preserving Z pointwise and such that $\sigma(c_i) = c_i$ for each curve $c_i \in P_a$, so we may apply Lemma 4.17. By Lemma 4.17 (together with Corollary 4.18) there exists a pair of \mathfrak{DP} -equivalent double pants decompositions, one compatible with Z_j and another compatible with Z_{j+1} . In view of Lemma 4.16, this implies that $P_a^j(Z)$ is \mathfrak{DP} -equivalent to $P_a^j(Z_{j+1})$. \square

5. SURFACES WITH MARKED POINTS

In this section we generalize Theorem 4.20 and Theorem 6.3 to the case of surfaces with boundary or for surfaces with marked points.

Remark 5.1. We prefer to work with surfaces with holes instead of surfaces with marked points (the latter may be obtained from the former by contracting the boundary components). Since we never consider the boundary and the neighbourhood or the boundary, this makes no difference for our reasoning. In case of marked surfaces one need to extend the definition of a “pair of pants”: for marked surfaces

a pair of pants is a sphere with 3 “features”, each of the features may be either a hole or a marked point.

Let $S_{g,n}$ be a surface of genus g with n holes. Definitions of pants decomposition and double pants decomposition remain the same as in case of $n = 0$.

Definition 5.2 (Standard double pants decomposition of an open surface). A double pants decomposition (P_a, P_b) of $S_{g,n}$ is *standard* if P_a and P_b contain the same set of g handles (where a *handle* is a surface of genus 1 with 1 hole).

A standard double pants decomposition is *strictly standard* if any curve of (P_a, P_b) either belongs to both of P_a and P_b or is contained in some of g handles.

In the same way as in case of $n = 0$ we define: flips and handle twists, admissible double pants decompositions and the category $\mathcal{AP}_{g,n}$ of admissible pants decompositions. We will consider objects of $\mathcal{AP}_{g,n}$ as surfaces with holes, but contracting the boundaries of the surface we obtain the equivalent category whose objects are surfaces with marked points.

Before proving the transitivity of morphisms on the objects of $\mathcal{AP}_{g,n}$, we reprove Theorem 3.2 for the case of open surfaces (the result is contained in [5], but we present the prove here, since the same idea works for double pants decompositions). More precisely, we derive the result for open surfaces from the result for closed ones.

Given a pants decomposition of an open surface, \mathcal{S} -moves are defined in the same way as for closed surfaces.

Lemma 5.3 (A. Hatcher [5]). *Flips and \mathcal{S} -moves act transitively on pants decompositions of $S_{g,n}$.*

Proof. The proof is by induction on the number of holes n . The base ($n = 0$) is Theorem 3.2 for the case of closed surfaces. Suppose that the lemma holds for $n = k$ and consider a surface $S_{g,k+1}$.

We consider simultaneously two surfaces, $S_{g,k+1}$ and $S_{g,k}$, where the latter is thought as a copy of $S_{g,k+1}$ with a disk attached to the boundary of $(k+1)$ -th hole. Each curve on $S_{g,k+1}$ turns into a curve on $S_{g,k}$ (but two distinct curves may became the same). Any pants decomposition of $S_{g,k+1}$ turns into a pants decomposition of $S_{g,k}$ containing one pair of pants less than the initial ones. More precisely, each pants decomposition $S_{g,k+1}$ contains a unique pair of pants one of whose boundary components is $(k+1)$ -th hole. This pair of pants disappears in $S_{g,k}$ (when the hole is removed, two other boundary components turn in the same curve). To go back from a pants decomposition of $P_{g,k}$ of $S_{g,k}$ to a pants decomposition of $S_{g,k+1}$ we need only to choose one of the curves $c \in P_{g,k}$ and attach in the place of c a thin strip containing a hole.

A flip as in Fig. 5.1 allows to change the curve $c \in P(S_{g,k})$ where the holed strip as attached (this flip in the decomposition of $S_{g,k+1}$ does not change the decomposition of $S_{g,k}$). Applying a sequence of flips we may move the strip to any given curve of the pants decomposition of $S_{g,k}$. Furthermore, for any flip or \mathcal{S} -move in the decomposition of $S_{g,k}$ we may apply similar transformation in $S_{g,k+1}$ (we only

need to check in advance that the holed strip is not attached to the curve changed by the transformation, in the latter case, first we need to change the “stripped” curve). So the transitivity of flips and \mathcal{S} -moves on pants decompositions of $S_{g,k+1}$ follows now from transitivity for $S_{g,k}$ and a fact that flips allow as to choose the stripped curve arbitrary. \square




FIGURE 5.1. Flip changing the curve where the holed strip is attached

Definition 5.4 (Simple double pants decomposition). A double pants decomposition (P_a, P_b) is *simple* if $|a_i \cap b_j| \leq 1$ for all curves $a_i \in P_a, b_j \in P_b$.

Theorem 5.5. *Morphisms of $\mathfrak{ADP}_{g,n}$ act transitively on the elements of $\mathfrak{ADP}_{g,n}$.*

Proof. The proof is by induction on the number of holes n . The base ($n = 0$) is proved in Theorem 4.20. Suppose that the theorem holds for $n = k$ and consider a surface $S_{g,k+1}$.

Following the proof of Lemma 5.3, we consider simultaneously double pants decompositions (P_a, P_b) of $S_{g,k+1}$ and $(\tilde{P}_a, \tilde{P}_b)$ of $S_{g,k}$. Each of two pants decompositions of $S_{g,k+1}$ differs from corresponding pants decomposition of $S_{g,k}$ by a holed strip attached in some of curves (so that $(k+1)$ -th hole in $S_{g,k+1}$ is lying in the intersection of two strips). The transitivity for the case of $S_{g,k}$ shows that flips and handle twists are sufficient to transform the double pants decomposition of $S_{g,k+1}$ to one which projects to any given double pants decomposition of $S_{g,k}$. As it is shown in the proof of Lemma 5.3, flips also allow to choose the curves of $(\tilde{P}_a, \tilde{P}_b)$ where the holed strips are attached. This implies transitivity of morphisms of $\mathfrak{ADP}_{g,n}$ on all admissible double pants decomposition of $S_{g,k+1}$ which project to simple double pants decomposition of $S_{g,k}$.

The reasoning above does not work for double pants decomposition of $S_{g,k+1}$ which do not project to simple double pants decompositions of $S_{g,k}$: indeed, in this case we may choose the double pants decomposition $(\tilde{P}_a, \tilde{P}_b)$ of $S_{g,k}$ and the curves a_i and b_j where the strips are attached, but in the case $|a_i \cap b_j| > 1$ we are not able to choose which of the intersections of the strips contains the hole.

To adjust the proof to this case, notice that a strictly standard double pants decomposition of $S_{g,k+1}$ projects to a strictly standard double pants decomposition of $S_{g,k+1}$, which is simple. This implies transitivity on strictly standard pants decompositions, and hence, on standard ones. In view of definition of admissible double pants decomposition (as one which may be obtained from a standard one), we have transitivity for all admissible double pants decompositions of $S_{g,k+1}$.

Thus, given the statement for $n = k$ we have proved it for $n = k + 1$, hence, the theorem holds for any integer $n \geq 0$. \square

Theorem 5.5 completes the proof of the Main Theorem.

6. FLIP-TWIST GROUPOID AND MAPPING CLASS GROUP

In general, the elements of the flip-twist groupoid FT change the topology of the double pants decomposition, so FT is not a group. However, there are some elements which preserve the topology. Clearly, these elements belong to mapping class group $MCG(S)$ of the surface (recall that a *mapping class group* $MCG(S)$ of a surface S is a group of homotopy classes of self-homeomorphisms of S with a composition as a group operation). In fact, all elements of mapping class group occur to belong to FT .

In this section, we consider the curves of a double pants decomposition as *labeled curves*, so that a symmetry of S interchanging the curves would not be trivial.

A double pants decomposition (P_a, P_b) contains a *double curve* c if $c \in P_a \cap P_b$.

Lemma 6.1. *Let $S = S_{g,n}$ be a genus g surface with $n \geq 0$ marked points, where $2g+n > 2$. Let (P_a, P_b) be an admissible double pants decomposition without double curves. Then $g \in MCG(S)$ fixes (P_a, P_b) if and only if $g = id$.*

Proof. Suppose there exists an element $g \in MCG(S)$ such that $g((P_a, P_b)) = (P_a, P_b)$, $g \neq id$. Since $g(P_a) = P_a$, g is a composition of Dehn twists along the curves contained in P_a . Since the curves do not intersect each other, the Dehn twists do commute. Let $a_1 \in P_a$ be any curve whose twist contributes to g . By assumption, $a_1 \notin P_b$. Hence, there exists a curve $b_1 \in P_b$ such that $b_1 \cap a_1 \neq \emptyset$ (otherwise P_b is not a maximal set of non-intersecting curves in S). Then $g(b_1) \neq b_1$, so $g(P_b) \neq P_b$, which contradicts the assumption. The contradiction implies the lemma. \square

An example of such a decomposition on a surface with marked points is shown in Fig. 6.1.




FIGURE 6.1. Double pants decomposition (P_a, P_b) without double curves of a surface S with marked points

Definition 6.2 (Category of topological surfaces). A category $\mathfrak{Top}_{g,n}$ of *topological surfaces* is one whose *objects* are topological surfaces of genus g with n marked points and whose *morphisms* are elements of mapping class group $MCG(S)$.

Theorem 6.3. *For any pair (g, n) such that $2g + n > 2$ the category $\mathfrak{ADP}_{g,n}$ contains a subcategory $\mathfrak{TopDP}_{g,n}$ which is isomorphic to $\mathfrak{Top}_{g,n}$.*

Proof. Consider an admissible double pants decomposition DP without double curves. Consider the orbit $MCG(DP)$ of DP under the action of the mapping class group. It follows from Lemma 6.1 that for $g_1, g_2 \in MCG$, $g_1 \neq g_2$ one has $g_1(DP) \neq g_2(DP)$. Let $\mathfrak{TopDP}_{g,n}$ be a subcategory of $\mathfrak{ADP}_{g,n}$ such that the objects of $\mathfrak{TopDP}_{g,n}$ are elements of the orbit $MCG(DP)$. The assumptions of Lemma 6.1 imply that the objects of $\mathfrak{TopDP}_{g,n}$ are in one-to-one correspondence with the objects of $\mathfrak{Top}_{g,n}$. Furthermore, Theorem 4.20 implies that for each two objects $x, y \in \mathfrak{TopDP}_{g,n}$ there exists a morphism $x \rightarrow y$. So, the morphisms of $\mathfrak{Top}_{g,n}$ and $\mathfrak{TopDP}_{g,n}$ are in one-to-one correspondence and we have an equivalence of two categories. \square

Remark 6.4. The special choice of double pants decomposition without double curves in Theorem 6.3 is indispensable: if $c_1, \dots, c_k \in DP$ are double curves then all Dehn twists along c_1, \dots, c_k preserve DP and the orbit $MCG(DP)$ gives a subcategory of $\mathfrak{ADP}_{g,n}$ isomorphic to some proper subcategory of $\mathfrak{Top}_{g,n}$.

REFERENCES

- [1] J. E. Andersen, A. J. Bene, and R. C. Penner, *Groupoid extensions of mapping class representations for bordered surfaces*, Topology Appl. **156** (2009), no. 17, 2713–2725. MR [2556030](#). Preprint version: [arXiv:0710.2651](#) [math.GT].
- [2] B. Bakalov and A. Kirillov, Jr., *On the lego-Teichmüller game*, Transform. Groups **5** (2000), no. 3, 207–244. MR [1780935](#)
- [3] A. Casson and C. Gordon, *Manifolds with irreducible Heegaard splittings of arbitrarily high genus*, preprint.
- [4] W. J. Harvey, *Boundary structure of the modular group*, Riemann surfaces and related topics: Proceedings of the 1978 Stony Brook Conference (State Univ. New York, Stony Brook, N.Y., 1978), Ann. of Math. Stud., vol. 97, Princeton Univ. Press, Princeton, N.J., 1981, pp. 245–251. MR [624817](#)
- [5] A. Hatcher, *Pants decompositions of surfaces*, Preprint <http://www.math.cornell.edu/~hatcher/Papers/pantsdecomp.pdf>.
- [6] A. Hatcher, *private communication*.
- [7] A. Hatcher and W. Thurston, *A presentation for the mapping class group of a closed orientable surface*, Topology **19** (1980), no. 3, 221–237. MR [579573](#)
- [8] J. Hempel, *3-manifolds as viewed from the curve complex*, Topology **40** (2001), no. 3, 631–657. MR [1838999](#)
- [9] E. Irmak and M. Korkmaz, *Automorphisms of the Hatcher-Thurston complex*, Israel J. Math. **162** (2007), 183–196. MR [2365859](#)
- [10] N. V. Ivanov, *Automorphism of complexes of curves and of Teichmüller spaces*, Internat. Math. Res. Notices (1997), no. 14, 651–666. MR [1460387](#)
- [11] F. Luo, *On Heegaard diagrams*, Math. Res. Lett. **4** (1997), no. 2-3, 365–373. MR [1453066](#)
- [12] D. Margalit, *Automorphisms of the pants complex*, Duke Math. J. **121** (2004), no. 3, 457–479. MR [2040283](#)

INDEPENDENT UNIVERSITY OF MOSCOW

Current address: Jacobs University Bremen

E-mail address: a.felikson@jacobs-university.de

HIGHER SCHOOL OF ECONOMICS, MOSCOW, BELOZERSKY INSTITUTE OF PHYSICO-CHEMICAL BIOLOGY, MOSCOW STATE UNIVERSITY,, INSTITUTE THEORETICAL AND EXPERIMENTAL PHYSICS

E-mail address: natanzons@mail.ru