


Polynomial-time approximation algorithms for the coloring problem in some cases

D. S. Malyshev

**Journal of Combinatorial
Optimization**

ISSN 1382-6905
Volume 33
Number 3

J Comb Optim (2017) 33:809-813
DOI 10.1007/s10878-016-0008-x



 Springer

Your article is protected by copyright and all rights are held exclusively by Springer Science +Business Media New York. This e-offprint is for personal use only and shall not be self-archived in electronic repositories. If you wish to self-archive your article, please use the accepted manuscript version for posting on your own website. You may further deposit the accepted manuscript version in any repository, provided it is only made publicly available 12 months after official publication or later and provided acknowledgement is given to the original source of publication and a link is inserted to the published article on Springer's website. The link must be accompanied by the following text: "The final publication is available at link.springer.com".

Polynomial-time approximation algorithms for the coloring problem in some cases

D. S. Malyshev¹

Published online: 16 March 2016

© Springer Science+Business Media New York 2016

Abstract We consider the coloring problem for hereditary graph classes, i.e. classes of simple unlabeled graphs closed under deletion of vertices. For the family of the hereditary classes of graphs defined by forbidden induced subgraphs with at most four vertices, there are three classes with an open complexity of the problem. For the problem and the open three cases, we present approximation polynomial-time algorithms with performance guarantees.

Keywords Coloring problem · Computational complexity · Approximation algorithm · Performance guarantee

1 Introduction

A graph H is called an *induced subgraph* of G if H is obtained from G by deletion of vertices. A class of graphs is called *hereditary* if it is closed under deletion of vertices. It is well-known that any hereditary (and only hereditary) class \mathcal{X} can be defined by a set of its forbidden induced subgraphs \mathcal{Y} . We write $\mathcal{X} = \text{Free}(\mathcal{Y})$ in this case, and graphs in \mathcal{X} are said to be \mathcal{Y} -free.

An *independent set* in a graph is a subset of its pairwise non-adjacent vertices. The size of a maximum independent set of a graph G is called the *independence number* of G and denoted by $\alpha(G)$. The *independent set problem* is to verify, for a given graph G and a natural number k , whether $\alpha(G) \geq k$ or not. It is a classical NP-complete problem.

✉ D. S. Malyshev
dsmalyshev@rambler.ru; dmalyshev@hse.ru

¹ National Research University Higher School of Economics, 25/12 Bolshaja Pecherskaja Ulitsa, Nizhny Novgorod 603155, Russia

A *coloring* of a graph G is a mapping $c : V(G) \rightarrow \mathbb{N}$, such that $c(u) \neq c(v)$ for any adjacent vertices $u, v \in V(G)$. This notion can be defined in another way via partition into independent sets. A *coloring* of a graph is partitioning the set of its vertices into independent sets called *color classes*. The minimum number of colors in colorings of G is called the *chromatic number of G* and denoted $\chi(G)$. The *chromatic number problem* is to verify, for a given graph G and a natural number k , whether $\chi(G) \leq k$ or not. It is a classical NP-complete problem.

By $\chi^*(G)$ we denote the minimal number of color classes in the colorings of a graph G having at most two vertices in every color class. Clearly, for any n -vertex graph G , we have $\chi^*(G) = n - \pi(\overline{G})$, where $\pi(H)$ is the *matching number of a graph H* , i.e. the maximum number of pairwise non-adjacent edges of H . It is well-known that the matching number of a n -vertex graph can be computed in $O(n^4)$ time (Edmonds 1965). Hence, for any n -vertex graph G , $\chi^*(G)$ can be computed in the same amount of time.

The computational complexity of the coloring problem in the hereditary classes of graphs defined by forbidden induced subgraphs with at most four vertices was studied in the paper Lozin and Malyshev (2015). For all but three classes in this family, it was shown either NP-completeness or polynomial-time solvability of the problem. The three exceptional cases are the classes $Free(\{O_4, C_4\})$, $Free(\{K_{1,3}, O_4\})$, $Free(\{K_{1,3}, O_4, K_2 + 2K_1\})$. Moreover, it was shown in Lozin and Malyshev (2015) that the coloring problem for $Free(\{K_{1,3}, O_4\})$ is polynomially equivalent to the same problem for $Free(\{K_{1,3}, O_4, K_2 + 2K_1\})$. In this paper, we do not find out the complexity of the problem for the two open cases. We present approximation polynomial-time algorithms for the problem with some “asymptotic” performance guarantees. More specific, for a graph G , we present a polynomial-time algorithm for computing a number $p(G)$, such that $\chi(G) \leq p(G) \leq r \cdot \chi(G) + O(1)$, where $r = \frac{3}{2}$ if G is $\{O_4, K_{3,3}\}$ -free and $r = \frac{4}{3}$ if G is $\{K_{1,3}, O_4, K_2 + 2K_1\}$ -free. As $Free(\{O_4, C_4\}) \subseteq Free(\{K_{3,3}, O_4\})$, the performance guarantee for $\{K_{3,3}, O_4\}$ -free graphs also holds for $\{O_4, C_4\}$ -free graphs.

2 Notation

We use the standard notation K_n, O_n, C_n for a complete, an empty graph, a chordless cycle with n vertices, respectively. A graph $K_{p,q}$ is a complete bipartite graph with p vertices in the first part and q in the second. The graph *fork* is obtained from a $K_{1,3}$ by subdividing an arbitrary its edge. The graph $K_4 - e$ is obtained from a K_4 by deleting an arbitrary its edge.

The formula $N(x)$ denotes the neighborhood of a vertex x .

The *sum* $G_1 + G_2$ is the disjoint union of graphs G_1 and G_2 with non-intersected sets of vertices. A graph \overline{G} is the complement of a graph G . A graph kG is the disjoint union of k copies of G . For a graph G and a subset $V' \subseteq V(G)$, $G \setminus V'$ denotes the subgraph of G obtained by deleting all elements of V' .

3 Approximation algorithms

3.1 The case of $\{O_4, K_{3,3}\}$ -free graphs

For a $\{O_4, K_{3,3}\}$ -free graph G , $\chi^*(G)$ computed in polynomial time gives an approximation for $\chi(G)$ with the performance guarantee almost $\frac{3}{2}$, as the following theorem shows.

Theorem 1 *For any $\{O_4, K_{3,3}\}$ -free graph G , we have $\chi(G) \leq \chi^*(G) \leq \frac{3}{2} \cdot \chi(G) + 1$.*

Proof Let us consider an optimal coloring of G . Let k_1 be the number of its three-vertex color classes, k_2 be the number of its color classes with at most two vertices. Since G is O_4 -free, $\chi(G) = k_1 + k_2$. As G is $K_{3,3}$ -free, any two of the k_1 color classes have two non-adjacent vertices in distinct classes. Hence, the union of the k_1 color classes can be partitioned into $\lceil \frac{3k_1}{2} \rceil$ subsets, each of which induces a O_1 or a O_2 . Therefore, G has a coloring with $\lceil \frac{3k_1}{2} \rceil + k_2$ color classes, each of which has at most two elements. Hence, $\chi^*(G) \leq \lceil \frac{3k_1}{2} \rceil + k_2 \leq \frac{3k_1}{2} + k_2 + 1 \leq \frac{3}{2} \cdot (k_1 + k_2) + 1 = \frac{3}{2} \cdot \chi(G) + 1$. Clearly, $\chi^*(G) \geq \chi(G)$. \square

3.2 The case of $\{O_4, K_{1,3}, K_2 + 2K_1\}$ -free graphs

Let G be a graph. By $T(G)$ we denote the graph, whose vertices are the triangles of G and two its vertices are adjacent if and only if the corresponding triangles intersect. Given G , $T(G)$ can be constructed in polynomial time on $|V(G)|$. Let $\alpha^*(G) \triangleq \alpha(T(G))$.


Lemma 1 *If G is $\{K_4, K_4 - e, K_3 + K_1\}$ -free, then the independence number of $T(G)$ can be computed in polynomial time on $|V(G)|$.*

Proof Firstly, we will show that $T(G)$ is $fork + 56K_1$ -free. Any triangle (x, y, z) of G defines a “3-coloring” of $G \setminus V'$ as follows, where $V' = \{x, y, z\}$. We use the term “3-coloring” in a non-classical sense assuming that some adjacent vertices can have the same color. As G is $\{K_4, K_4 - e, K_3 + K_1\}$ -free, each of the elements of $V(G) \setminus V'$ is adjacent to exactly one element of V' . All neighbors of an element $a \in V'$ in $V(G) \setminus V'$ have the same color.

Assume that $T(G)$ contains an induced $fork$. Hence, G contains five pairwise distinct triangles (x_1, x_2, x_3) , (y_1, y_2, y_3) , (z_1, z_2, z_3) , (x_1, y_1, u_1) , (u_1, u_2, u_3) , where $z_1 = u_3$ (see the figure below). Additionally, assume that there is a triangle (a_1, a_2, a_3) of G non-intersecting with each of the five triangles. Let $c : V(G) \setminus \{a_1, a_2, a_3\} \rightarrow \{1, 2, 3\}$ be the “3-coloring” defined above, where $c(w) = i$ for every $w \in V(G) \setminus \{a_1, a_2, a_3\}$ adjacent to a_i . Since G is $\{K_4, K_4 - e\}$ -free, vertices of each of the five triangles have pairwise distinct colors. Hence, there is an element of $\{x_2, x_3\}$, an element of $\{y_2, y_3\}$, an element of $\{z_2, z_3\}$ having the same color as the color of u_1 . Without loss of generality, $c(x_3) = c(y_3) = c(z_3) = c(u_1) = 3$, $c(x_1) = c(u_2) = c(y_2) = c(z_2) = 1$, $c(y_1) = c(x_2) = c(u_3) = 2$ (Fig. 1).

We will show that there is an edge $(v', v'') \in E(G)$, such that v' and v'' belong to some two of the five triangles and $c(v') = c(v'')$. Assume that there is no such an

Fig. 1 The configuration of the triangles



edge. As G is $\{K_4, K_4 - e, K_3 + K_1\}$ -free, $(x_1, z_3) \in E(G)$ and $(z_3, y_2) \in E(G)$. Similarly, $(y_2, z_1) \in E(G)$. We have a contradiction, as y_2 is adjacent to two vertices of the triangle (z_1, z_2, z_3) . So, for any triangle of G non-intersecting with each of the five triangles, there is a triangle of G intersecting with T , whose two vertices belong to $\bigcup_{i=1}^3 \{x_i, y_i, z_i, u_i\}$. The subgraph of G induced by $\bigcup_{i=1}^3 \{x_i, y_i, z_i, u_i\}$ has at most $\frac{11 \cdot (11-1)}{2} = 55$ edges. Hence, by the pigeonhole principle, $T(G)$ must be $fork + 56K_1$ -free, otherwise it contains an induced K_4 or $K_4 - e$.

Clearly, the equality $\alpha(H) = \max(\alpha(H \setminus \{v\}), \alpha(H \setminus N(v)))$ holds for any graph H and its vertex v . Hence, the independent set problem for $fork + 56K_1$ -free graphs can be polynomially reduced to the same problem for $fork$ -free graphs. The independent set problem for $fork$ -free graphs can be solved in polynomial time (Alekseev 2004). Hence, the lemma holds. \square

Lemma 2 For every n -vertex $\{K_{1,3}, O_4, K_2 + 2K_1\}$ -free graph G , the inequalities $0 < \chi^*(G) - \frac{\alpha^*(\overline{G})+1}{2} \leq \chi(G) \leq \min(\chi^*(G), n - 2\alpha^*(\overline{G}))$ hold.

Proof The inequality $\chi(G) \leq \chi^*(G)$ is obvious. Consider a set of $\alpha^*(\overline{G})$ vertex-disjoint triangles of \overline{G} . There are $n - 3\alpha^*(\overline{G})$ vertices of \overline{G} not belonging to the triangles in the set. Hence, there is a coloring of G with $n - 2\alpha^*(\overline{G})$ color classes, $\alpha^*(\overline{G})$ of them have three vertices and $n - 3\alpha^*(\overline{G})$ of them have one element. Hence, $\chi(G) \leq n - 2\alpha^*(\overline{G})$. Consider an arbitrary optimal coloring of G . Every its color class contains at most three vertices, as G is O_4 -free. Let k be the number of the color classes in the coloring having exactly three vertices. As G is $K_{1,3}$ -free, any two of the color classes with three vertices have two non-adjacent vertices in distinct classes. Hence, the union of all k color classes with three vertices can be partitioned into $\lceil \frac{3k}{2} \rceil$ subsets, each of them induces a O_2 or a O_1 . Therefore, there is a coloring of G in at most $\chi(G) + \lceil \frac{3k}{2} \rceil - k$ colors, such that every color class has at most two vertices. Clearly, $\chi(G) + \lceil \frac{3k}{2} \rceil - k \geq \chi^*(G)$, i.e. $\chi(G) \geq \chi^*(G) + k - \lceil \frac{3k}{2} \rceil$. As $k - \lceil \frac{3k}{2} \rceil \geq -\frac{k+1}{2}$ and $\alpha^*(\overline{G}) \geq k$, $\chi(G) \geq \chi^*(G) - \frac{\alpha^*(\overline{G})+1}{2}$. The inequality $\chi^*(G) - \frac{\alpha^*(\overline{G})+1}{2} > 0$ is trivial whenever G is the single-vertex graph. As $\chi^*(G) \geq \frac{n}{2}$ and $\alpha^*(\overline{G}) \leq \frac{n}{3}$, $\chi^*(G) - \frac{\alpha^*(\overline{G})+1}{2} \geq \frac{2n-1}{3} > 0$ whenever $n \geq 2$. \square

Theorem 2 There is an approximation polynomial-time algorithm for the coloring problem for $\{K_{1,3}, O_4, K_2 + 2K_1\}$ -free graphs having the performance guarantee almost $\frac{4}{3}$.

Proof Let G be a n -vertex $\{K_{1,3}, O_4, K_2 + 2K_1\}$ -free graph. The complement of any $\{K_{1,3}, O_4, K_2 + 2K_1\}$ -free graph is $\{K_4, K_4 - e, K_3 + K_1\}$ -free. By Edmonds (1965) and Lemma 1, $\chi^*(G)$ and $\alpha^*(\overline{G})$ can be computed in polynomial time on n . Assume that $\alpha^*(\overline{G}) \leq \frac{n}{4}$. Hence, $\chi^*(G) - \frac{\alpha^*(\overline{G})+1}{2} \geq \chi^*(G) - \frac{\alpha^*(\overline{G})}{2} - \frac{1}{2} \geq \frac{3}{4}\chi^*(G) - \frac{1}{2}$, as $\chi^*(G) \geq \frac{n}{2}$ and $\alpha^*(\overline{G}) \leq \frac{n}{4}$. Therefore, $\chi^*(G)$ approximates $\chi(G)$ with an factor which is asymptotically at most $\frac{4}{3}$, since $\frac{\chi^*(G)}{\chi(G)} \leq \frac{\chi^*(G)}{\chi^*(G) - \frac{\alpha^*(\overline{G})+1}{2}}$ (by Lemma 2) and $\chi^*(G) \geq \chi(G)$. Assume that $\alpha^*(\overline{G}) > \frac{n}{4}$. Hence, $\chi^*(G) \geq \frac{n}{2} > n - 2\alpha^*(\overline{G})$. By Lemma 2, $1 \leq \frac{n-2\alpha^*(\overline{G})}{\chi(G)} \leq \frac{n-2\alpha^*(\overline{G})}{\chi^*(G) - \frac{\alpha^*(\overline{G})+1}{2}}$. Clearly, $\frac{n-2\alpha^*(\overline{G})}{\chi^*(G) - \frac{\alpha^*(\overline{G})+1}{2}} \leq 2\frac{n-2\alpha^*(\overline{G})}{n-\alpha^*(\overline{G})}$, as $\chi^*(G) \geq \frac{n}{2}$. Moreover, $2\frac{n-2\alpha^*(\overline{G})}{n-\alpha^*(\overline{G})}$ is equal to the value of the function $f(x) = 2\frac{1-2x}{1-x}$ at the point $x = \frac{\alpha^*(\overline{G})}{n}$. As $f'(x) = -\frac{2}{(1-x)^2}$, the function $f(x)$ is monotonically decreasing over the segment $[\frac{1}{4}, \frac{1}{3}]$. Hence, its maximum value over the segment is equal to $f(\frac{1}{4}) = \frac{4}{3}$. Therefore, $n - 2\alpha^*(\overline{G})$ approximates $\chi(G)$ with an approximation ratio, which is asymptotically at most $\frac{4}{3}$. \square

4 Concluding remarks

Recall that there are three hereditary classes defined by forbidden induced subgraphs with at most four vertices for which the computational complexity of the coloring problem is open. The cases are $Free(\{O_4, C_4\})$, $Free(\{K_{1,3}, O_4\})$, $Free(\{K_{1,3}, O_4, K_2 + 2K_1\})$. The problem seems to be NP-complete for $Free(\{K_{1,3}, O_4\})$, $Free(\{K_{1,3}, O_4, K_2 + 2K_1\})$ and polynomial-time solvable for $Free(\{O_4, C_4\})$. Clarifying its complexity status in the classes is an interesting problem for future research. For all of the three classes, we presented polynomial-time approximation algorithms with specified performance guarantees. Designing efficient approximation algorithms with better performance guarantees for them is also an interesting problem for future research.

Acknowledgements This article is partially supported by Russian Foundation for Basic Research, Grants 16-31-60008-mol-a-dk and 16-01-00599-a; by RF President Grant MK-4819.2016.1; by LATNA laboratory, National Research University Higher School of Economics.

References

Alekseev V (2004) Polynomial algorithm for finding the largest independent sets in graphs without forks. Discret Appl Math 135(1–3):3–16
 Edmonds J (1965) Paths, trees, and flowers. Can J Math 17:449–467
 Lozin V, Malyshev D (2015) Vertex coloring of graphs with few obstructions. Discret Appl Math. doi:10.1016/j.dam.2015.02.015